

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2007 (09.08.2007)

PCT

(10) International Publication Number
WO 2007/089768 A2

(51) International Patent Classification:

C07D 239/42 (2006.01) **C07D 403/12** (2006.01)
C07D 401/04 (2006.01) **C07D 405/04** (2006.01)
C07D 401/12 (2006.01) **C07D 405/12** (2006.01)
C07D 401/14 (2006.01) **C07D 405/14** (2006.01)

(21) International Application Number:

PCT/US2007/002515

(22) International Filing Date: 30 January 2007 (30.01.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/763,426 30 January 2006 (30.01.2006) US
60/785,239 23 March 2006 (23.03.2006) US
60/840,420 25 August 2006 (25.08.2006) US

(71) Applicant (for all designated States except US): **EX-ELIXIS, INC.** [US/US]; Post Office Box 511, 210 East Grand Avenue, South San Francisco, CA 94083-0511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MANN, Grace** [US/US]; 231 Callippe Court, Brisbane, California 94005 (US). **AAY, Naing** [US/US]; 112 North Rochester Street, San Mateo, CA 94401 (US). **ARCALAS, Arlyn** [US/US]; 434 Brosnan Court, South San Francisco, CA 94080 (US). **BROWN, S. David** [US/US]; 12 Sorrel Lane, San Carlos, California 94070 (US). **CHAN, Wai Ki Vicky** [US/US]; 442 Faxon Avenue, San Francisco, CA 94112 (US). **CHEN, Jeff** [US/US]; 140 South Van Ness Avenue, Apt. 407, San Francisco, California 94103 (US). **DU, Hong-wang** [CN/US]; 400 Richmond Drive, Apt. 4, Millbrae, CA 94030 (US). **EPSHTEYN, Sergey** [US/US]; 2922 Southwycke Ter, Fremont, CA 94536 (US). **FORSYTH, Timothy** [US/US]; 1928 Wingate Way, Hayward, California 94541 (US). **GALAN, Adam A.** [US/US]; 865 Cedar Street, Alameda, California 94501 (US). **HUYNH, Tai Phat** [US/US]; 1634 9th Avenue, Oakland, California 94606 (US). **IBRAHIM, Mohamed Abdulkader** [US/US]; 3380 Lubich Drive, Mountain View, CA 94040 (US). **JOHNSON, Henry William Beecroft** [US/US]; 441 Linden Avenue, San Bruno, CA 94066 (US). **KANE, Brian** [US/US]; 23 Clinton Street, Apt. A, Redwood City, CA 94062 (US). **KEARNEY, Patrick** [US/US]; 195 Bocana Street, San Francisco, CA 94110 (US). **KIM,**

Byung Gyu [KR/US]; 10 De Sabla Road, Apt. 801, San Mateo, California 94402 (US). **KOLTUN, Elena** [RU/US]; 1075 Grebe Street, Foster City, California 94404 (US). **LEAHY, James William** [US/US]; 1185 Camellia Court, San Leandro, CA 94577 (US). **LEE, Matthew Sangyup** [US/US]; 8311 Shelter Creek Lane, San Bruno, California 94066 (US). **LEWIS, Gary L.** [US/US]; 412 Noe Street, #5, San Francisco, California 94110 (US). **MEYR, Lisa E.** [US/US]; 13203 3rd Avenue Nw, Seattle, Washington 98177 (US). **NOGUCHI, Robin Tammie** [US/US]; 5213 Shelter Creek Lane, San Bruno, California 94066 (US). **PACK, Michael** [US/US]; 592 South Van Ness, San Francisco, CA 94110 (US). **RIDGWAY, Brian Hugh** [US/US]; 2403 Palmer Avenue, Belmont, California 94002 (US). **SHI, Xian** [CN/US]; 4222 Shelter Creek Lane, San Bruno, California 94066 (US). **WOOLFREY, John** [US/US]; 25961 Vinedo Lane, Los Altos, California 94022 (US). **ZHOU, Peiwen** [CN/US]; 765 San Antonio Road, #68, Palo Alto, CA 94303 (US).

(74) Agents: **BERNSTEIN, Robert L.** et al.; EXELIXIS, Inc., 210 East Grand Avenue Post Office Box 511, South San Francisco, CA 94083-0511 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-ARYL-2-AMINO-PYRIMIDINES OR 4-ARYL-2-AMINOALKYL-PYRIMIDINES AS JAK-2 MODULATORS AND METHODS OF USE

(57) Abstract: This invention relates to certain pyrimidine derivative inhibitors of JAK-2, having Formula (I): wherein D, E, L, Z, R¹, R², R²⁵, and n1 are as defined in the specification, pharmaceutically acceptable salts thereof, pharmaceutical compositions thereof, and methods of use thereof.

WO 2007/089768 A2

**4-ARYL-2-AMINO-PYRIMIDINES OR 4-ARYL-2-AMINOALKYL-PYRIMIDINES
AS JAK-2 MODULATORS AND METHODS OF USE**

10

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Application Ser. No. 60/840,420, filed on August 25, 2006, U.S. Provisional Application Ser. No. 60/785,239, filed on March 23, 2006, and U.S. Provisional Application Ser. No. 60/763,426, filed on January 30, 2006.

15

FIELD OF THE INVENTION

[0001] This invention relates to certain 4-aryl-2-amino-pyridines and 4-aryl-2-aminoalkyl-pyridines as inhibitors of protein tyrosine kinases. In particular, the invention relates to inhibitors of JAK-2 that involve the cytokine receptor signaling pathways.

20

BACKGROUND OF THE INVENTION

[0002] Janus kinases (JAKs) are protein tyrosine kinases ubiquitously expressed in cells. JAKs are involved in membrane signalling events which are triggered by a variety of extracellular factors that interact with cell surface receptors. JAKs initiate the cytoplasmic signal transduction cascades of cytokine receptors that lack a protein tyrosine kinase domain.

25

The signal transduction cascades are initiated after oligomerisation of surface receptors due to ligand binding. Cytoplasmic receptor-associated JAKs are then activated which subsequently phosphorylate tyrosine residues along the receptor chains. These phosphotyrosine residues are targets for a variety of SH2 domain-containing transducer proteins, such as the signal transducers and activators of transcription (STAT) proteins. After STAT binds to receptor chains, they are phosphorylated by the JAK proteins, dimerise and translocate into the nucleus. In the nucleus, STAT alter the expression of cytokine-regulated genes.

30

[0003] Mammalian JAK-2 belongs to a kinase family that include JAK-1, JAK-3 and TYK-2. JAK-1, JAK-2, and TYK-2 are ubiquitously expressed, while JAK-3 is predominantly expressed in hematopoietic cells. These kinases consist of approximately 1150 amino acids, with molecular weights of about 120 kDa to 130 kDa. The amino acid sequences of the JAK kinase family are characterised by the presence of highly conserved domains. These domains include the JAK homology (JH) domains, C-terminal domain (JH1)

35

5 responsible for the tyrosine kinase function, the tyrosine kinase-like domain (JH2) that shows high similarity to functional kinases but does not possess any catalytic activity, and the N-terminal domain that spans JH7 to JH3) that is important for receptor association and non-catalytic activity. Although the function of the N-terminal domain is not well established, there is some evidence for a regulatory role on the JH1 domain, thus modulating catalytic
10 activity.

[0004] The down-stream substrates of the JAK family of kinases include the signal transducer and activator of transcription (STAT) proteins. JAK/STAT signaling has been implicated in the mediation of many abnormal immune responses such as allergies, asthma, autoimmune diseases such as transplant rejection, rheumatoid arthritis, amyotrophic lateral
15 sclerosis and multiple sclerosis as well as in solid and hematologic malignancies such as leukemias and lymphomas.

[0005] Signal transducer and activator of transcription (STAT) proteins are activated by JAK family kinases. Recent studies suggested the possibility of modulating the JAK/STAT signaling pathway by targeting JAK family kinases with specific inhibitors for the treatment
20 of leukemia (see Sudbeck, et al., Clin. Cancer Res. 5: 1569-1582 (1999)). In animal models, TEL/JAK-2 fusion proteins induced myeloproliferative disorders (Schwaller, et al., EMBO J. 17: 5321-5333 (1998)). In hematopoietic cell lines, introduction of TEL/JAK2 resulted in activation of STAT1, STAT3, STAT5, and cytokine-independent growth (Schwaller, et al., EMBO J. 17: 5321-5333 (1998)).

5 [0006] The JAK/STAT pathway is involved in abnormal cell growth (Yu, et al., J. Immunol. 159: 5206-5210 (1997)). STAT3, STAT5, JAK1 and JAK2 are constitutively activated in mouse T cell lymphoma characterized initially by LCK over-expression. In addition, IL-6-mediated STAT3 activation was blocked by inhibition of JAK, leading to sensitization of myeloma cells to apoptosis (Catlett-Falcone, et al., Immunity 10:105-115
0 (1999)).

[0007] One particularly attractive target for small-molecule modulation, with respect to antiproliferative and antiangiogenic activity, is JAK-2. Accumulating evidence shows that constitutive activation of JAK/STAT pathway promotes cell growth, survival, differentiation, neoplastic transformation, and angiogenesis in response to growth factors, cytokines, and
5 hormones. JAK-2 is also activated in a wide variety of human cancers such as prostate, colon, ovarian, breast cancers, melanoma, leukemia and other haematopoietic malignancies. In addition, somatic point mutation of the JAK-2 gene has been identified to be highly

5 associated with classic myeloproliferative disorders (MPD) and infrequently in other myeloid disorders. Constitutive activation of JAK-2 activity is also caused by chromosomal translocation in hematopoietic malignancies, such as in TEL-JAK-2 which is primarily associated with T-ALL, and in PCM1-JAK-2 which is associated with B-ALL and CML. It has been shown that inhibition of the JAK/STAT pathway, and in particular inhibition of JAK-2 activity, results in anti-proliferative and pro-apoptotic effects largely due to inhibition of phosphorylation of STAT. Furthermore, inhibition of JAK-2 activity by pharmacological agents or by expression of dominant negative JAK-2 effectively block tumor growth and induce apoptosis by reducing the STAT phosphorylation in cell culture and human tumor xenografts *in vivo*. Therefore, the JAK/STAT signal transduction pathway is a well-validated target pathway for therapeutic intervention.

5 [0008] Accordingly, the identification of small-molecule compounds that specifically inhibit, regulate and/or modulate the signal transduction of kinases, particularly JAK-2, is desirable as a means to treat or prevent diseases and conditions associated with cancers. Thus, an object of this invention is the identification of JAK-2 inhibitors as new therapeutic agents to treat human diseases.


SUMMARY OF THE INVENTION

5 [0009] The invention relates to compounds for inhibiting JAK-2 and pharmaceutical compositions of the compounds for inhibiting JAK-2. The invention is also related to methods of treating a disease mediated at least in part by JAK-2 using a compound of the invention, or a pharmaceutical composition thereof, as well as in combination with other therapies.

0 [0010] The foregoing only summarizes certain aspects of the invention and is not intended to be limiting in nature. These aspects and other aspects and embodiments are described more fully below. All references any sort referred to in this specification are hereby incorporated by reference in their entirety. In the event of a discrepancy between the express disclosure of this specification and the references incorporated by reference, the express disclosure of this specification shall control.

DETAILED DESCRIPTION OF THE INVENTION

[0011] A first aspect of the invention relates to a compound of Formula I:



I

or a pharmaceutically acceptable salt thereof, wherein

D is hydrogen, halo, -CF₃, heterocycloalkyl or alkyl;

E is hydrogen, halo, -CF₃, heterocycloalkyl or alkyl; or

- 10 D and E, together with the carbon atoms to which they are attached, form a 5-7 membered heteroaryl or a 5-7 membered heterocycloalkyl, wherein the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl are each fused to the pyrimidinyl moiety to which D and E are attached;

L is a bond, -O- or -N(H)-;

- 15 Z is selected from alkoxy, cycloalkyl, heteroaryl optionally substituted with alkyl, halo, -C(O)OR²⁶, -C(=N-OH)alkyl, -C(O)R⁸, -C(O)NR³⁰R^{30a}, -CH₂R², -(CH₂)_{n5}NR²⁶R^{26a}, -CF₃, -CN, -SO₂R¹², -S-R^{12a}, -OR^{32a},

- Z and R²⁵, together with the carbon atoms to which they are attached, join to form a 5 or 6 membered heterocycloalkyl, a 5 or 6 membered heteroaryl, or a 5 or 6 membered cycloalkyl ring, wherein the 5 or 6 membered heterocycloalkyl, 5 or 6 membered heteroaryl, and 5 or 6 membered cycloalkyl ring are fused to the phenyl moiety to which Z and R²⁵ are attached, and wherein the 5 or 6 membered heterocycloalkyl, 5 or 6 membered heteroaryl, or 5 or 6 membered cycloalkyl ring are each optionally substituted with 1, 2, or 3 groups independently selected from oxo, alkyl, alkoxy and halo;

n₁ is 0, 1, 2, 3, or 4, and each n₁ is independently selected when more than one n₁ is present;

n₂ is 0, 1, 2, 3, or 4, and each n₂ is independently selected when more than one n₂ is present;

n₃ is 0, 1, 2, or 3, and each n₃ is independently selected when more than one n₃ is present;

n₄ is 0, 1, 2, 3 or 4, and each n₄ is independently selected when more than one n₄ is present;






















- 0 n₅ is 0, 1, 2, 3 or 4, and each n₅ is independently selected when more than one n₅ is present;

p is 0-3;

r is 1-3;

R¹ is hydrogen;

5 R^2 is a group of formula:



5

R^7 , $R^{7'}$, R^9 , R^{10} , R^{12} and R^{15} are each independently hydrogen, alkyl, alkoxy, or alkoxyalkyl; R^8 is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxylamino, hydroxyalkyl, alkoxyalkyl, dihydroxyalkyl, alkylamino, dialkylamino, aminoalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, alkylaminoalkyl, dialkylaminoalkyl, $-(CH_2)_r-C(O)OR^7$, $-(CH_2)_r-C(O)NR^7R^{7'}$, aryl, heteroaryl, cycloalkyl, arylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with one, two, three, four or five groups independently selected from alkyl, alkenyl, lower alkynyl, halo, hydroxy, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclylalkoxy, oxo and haloalkyl;

10

15

each R^{11} , when R^{11} is present, is independently selected from alkyl, alkenyl, lower alkynyl, $-CF_3$, alkoxy, halo, haloalkoxy, haloalkyl, aminoalkyl, aminoalkoxy, alkylaminoalkyl, alkylaminoalkoxy, dialkylaminoalkyl, dialkylaminoalkoxy, oxo, thioalkyl, alkylthioalkyl, $-(CH_2)_p-OR^{17}$, $-CN$, $-O-CH_2-C(O)-R^{17}$, $-C(O)R^{16}$, $-(CH_2)_p-C(O)OR^{17}$, $-S(O)_2R^{17}$, $-S(O)_2NR^{15}R^{17}$, aryl, heteroaryl, cycloalkyl, arylalkyl, arylalkoxy, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at any ring position with 1, 2, 3 or 4 R^{21} ;

20

25

R^{12} is hydrogen or alkyl;

R^{12a} is hydrogen or alkyl;

R^{13} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxylamino,

30

haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, $-(CH_2)_r-C(O)OR^7$, $-(CH_2)_r-C(O)NR^7R^{7'}$, aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with 1, 2, 3, 4 or 5 groups independently selected from alkyl, alkenyl, lower alkynyl,

35

- 5 halo, hydroxy, hydroxyalkyl, alkoxycarbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl; and wherein the alkyl of cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, and heteroarylalkyl are independently optionally substituted with 1, 2, 3, 4, or 5 groups selected from halo and hydroxy;
- 10 R^{14} is a bond, heterocycloalkyl or cycloalkyl;
 R^{16} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxyamino, haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, dialkylaminoalkyl,
- 15 $-(CH_2)_r-C(O)OR^7$, aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with one, two, three, four or five groups
- 20 independently selected from alkyl, alkenyl, lower alkynyl, halo, hydroxy, hydroxyalkyl, alkoxycarbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl; and wherein the alkyl of cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, and heteroarylalkyl is optionally substituted with 1, 2, 3, 4, or 5 groups selected from halo and hydroxy;
- 25 R^{17} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxyamino, haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, dialkylaminoalkyl,
- 30 $-(CH_2)_r-C(O)OR^7$, $-(CH_2)_r-C(O)NR^7R^{7'}$, aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with one, two, three, four or five groups independently selected from alkyl, alkenyl,
- 35 lower alkynyl, halo, hydroxy, hydroxyalkyl, alkoxycarbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl; and wherein the alkyl of cycloalkylalkyl, heterocycloalkylalkyl,

- 5 arylalkyl, and heteroarylalkyl is optionally substituted with 1, 2, 3, 4, or 5 groups selected from halo and hydroxy;
- each R^{21} , when R^{21} is present, is independently selected from alkyl, alkenyl, lower alkynyl, cyano, halo, haloalkoxy, haloalkyl, hydroxyalkyl, amino, alkylamino, dialkylamino, dialkylaminoalkyl, dialkylaminoalkoxy, haloalkyl, oxo, $-OR^{13}$, $-NHS(O)_2R^{17}$,
 10 $-S(O)_2R^{17}$, $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)NR^{15}R^{17}$, $-NR^{15}C(O)R^{17}$, aryl, arylalkyl, heteroarylalkyl, aryloxy, and heteroaryl; wherein each of the aryl, arylalkyl, heteroarylalkyl, aryloxy, and heteroaryl within R^{21} are optionally substituted at any ring position with 1, 2, or 3 groups selected from alkyl, lower alkoxy halo, phenyl, heteroaryl and alkylheteroalkyl;
- 15 R^{25} is selected from alkyl, alkenyl, lower alkyl, halo, haloalkyl, haloalkoxy, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, $-OR^{12}$, cyano, $-C(O)R^8$, $-CH_2NHC(O)OR^7$, $-CH_2NHC(O)R^7$, $-SR^7$, $-S(O)_2R^7$, $-S(O)_2NR^7R^8$, $-C(O)OR^8$, $-C(O)NR^7R^8$, cycloalkyl, heterocycloalkyl, aryl and heteroaryl; wherein the cycloalkyl, heterocycloalkyl, aryl and heteroaryl are each optionally substituted with
 20 one, two or three groups independently selected from alkyl, alkenyl, halo, haloalkoxy, haloalkyl, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, $-OR^8$, $-NHS(O)_2R^8$, cyano, $-C(O)R^8$, $-CH_2NHC(O)OR^7$, $-CH_2NHC(O)R^7$, $-SR^7$, $-S(O)_2R^7$, $-S(O)_2NR^7R^8$, $-C(O)OR^8$, $-C(O)NR^7R^8$, $-NR^7C(O)-CHR^3-OR^8$, $-NR^7C(O)-CHR^3-NR^7-R^8$, and $-NR^7C(O)R^8$;
- 25 R^{26} is hydrogen, $-C(O)$ -phenyl or alkyl, wherein the $-C(O)$ -phenyl is optionally substituted at any ring position with 1, 2 or 3 halo;
- R^{26a} is hydrogen, alkyl, heteroaryl, $-C(O)R^{32}$, $-C(O)NHR^{32a}$, $-S(O)_2R^9$, $-SR^9$, $-C(O)OR^{32}$, or $-C(O)NR^{32a}R^{32}$;
- R^{27} and R^{28} are each independently selected from alkyl, alkenyl, hydroxy, alkoxy, and
 30 alkoxyalkyl;
- R^{27a} and R^{28a} are independently selected from hydrogen, alkyl, alkenyl, alkoxyalkyl, alkoxyalkylalkyl, hydroxyalkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, dialkylaminoalkyl, arylcarbonylalkyl, aryloxyalkyl, dialkylaminoalkyl, alkyl-O- $C(O)$ heterocycloalkyl, $-(CH_2)_{n4}$ heterocycloalkyl, heterocycloalkylalkyl, heteroaryl, heteroarylalkyl, $-(CH_2)_{n4}-C(O)R^{29}$, $-(CH_2)_{n4}NR^{28}R^{28a}$, $-(CH_2)_{n4}NHR^{28a}$, $-CH(phenyl)_2$,
 35 $-S(O)_2R^{29}$, $-C(O)R^{29}$, $-C(O)OR^{29}$, and $-C(O)NR^{29a}R^{29}$, wherein the aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and


- 5 heterocycloalkylalkyl groups within R^{27a} and R^{28a} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, alkoxy, alkylcarbonyl, phenyl, phenoxy, arylcarbonyl, $-CF_3$, oxo, $-OCF_3$, alkoxyphenyl, and heteroaryl optionally substituted with alkyl or halo;
- or R^{27} and R^{27a} , together with the nitrogen to which they are attached, form
- 10 heterocycloalkylamino, heterocycloalkyl or heteroaryl, wherein the heterocycloalkylamino and heteroaryl are each independently optionally substituted with 1, 2, 3, 4, or 5 R^{31} ;
- or R^{28} and R^{28a} together with the nitrogen to which they are attached form heterocycloalkyl or heteroaryl, wherein the heterocycloalkyl and heteroaryl are each optionally substituted
- 15 with 1, 2, 3, 4, or 5 R^{31} ;
- R^{29a} is hydrogen or alkyl;
- R^{29} is selected from alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl,
- 20 and heterocycloalkylalkyl groups within R^{29} are each optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, alkoxy, alkylcarbonyl, phenyl, phenoxy, arylcarbonyl, $-CF_3$, oxo, $-OCF_3$, alkoxyphenyl, and heteroaryl optionally substituted with alkyl or halo;
- R^{30a} is hydrogen or alkyl;
- 25 R^{30} is selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkoxycarbonylalkyl, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, aryl, arylalkyl, phenoxyalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, arylheteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, arylalkyl, phenoxyalkyl, cycloalkyl,
- 30 arylheteroarylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl groups within R^{30} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, alkoxy, alkoxyalkyl, $-C(O)OCH_3$, $-CF_3$, $-OCF_3$, alkylcarbonyl, phenyl, phenoxy, alkylphenoxy, dialkylaminoalkoxy and heteroaryl;
- 35 R^{31} is selected from alkyl, hydroxyalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, $-C(O)R^{30}$, $-C(O)NR^{30}R^{30a}$, $-C(O)OR^{30}$, $-S(O)_2R^{30}$, amino, dihydroxyalkyl, arylcarbonyl, alkylcarbonylamino, alkoxyphenyl, phenylalkoxyalkyl, arylheteroarylalkyl, alkylamino,

- 5 -O-dialkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, dialkylaminoalkoxy, oxo, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, spirocyclic cycloalkyl, spirocyclic heterocycloalkyl, and heterocycloalkylalkyl, wherein the aryl, arylalkyl, cycloalkyl, arylheteroarylalkyl, arylalkoxyalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and
- 10 heterocycloalkylalkyl groups within R^{31} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, $-CF_3$, $-OCF_3$, cyano, alkoxy, alkoxyalkyl, $-C(O)OCH_3$, alkylcarbonyl, phenyl optionally substituted at any ring position with halo, phenoxy, alkylphenoxy, arylalkoxyalkyl, dialkylaminoalkoxy and heteroaryl;
- 15 R^{32a} is hydrogen, $-OCF_3$, $-CF_3$, or alkyl;
 R^{32} is selected from aryl, arylalkyl, arylalkoxy, arylcycloalkyl, alkoxycarbonylalkoxy, cycloalkyl, cycloalkylalkyl, cycloalkylhydroxyalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl, wherein the aryl, arylalkyl, cycloalkyl, arylcycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and
- 20 heterocycloalkylalkyl are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from hydroxy, oxo, alkyl, alkoxy, amino, hydroxyalkyl, alkylcarbonyl, alkoxycarbonyl, halo, $-CF_3$, $-OCF_3$, aminoalkyl, alkylaminoalkyl, aryl and dialkylaminoalkyl, and wherein the alkyl portion of the heteroarylalkyl can be substituted with amino;
- 25 or R^{32} is alkyl optionally substituted with 1, 2, 3, 4, or 5 groups independently selected from hydroxy, alkoxycarbonyl, alkoxy, $-CF_3$, halo, , aminocarbonyl, alkylaminocarbonyl, alkoxycarbonylalkylamino, dialkylaminocarbonyl, $-NR^{34}R^{34a}$ and phenyl optionally substituted with 1, 2, or 3 halo;
- or R^{32} is alkylamino or arylalkylamino;
- 30 R^{34} is hydrogen or alkyl;
 R^{34a} is selected from hydrogen, alkyl, heteroaryl, aryl, aminoalkyl, aminocarbonylalkyl, heteroarylalkyl, arylalkoxy and arylalkyloxycarbonylalkyl; wherein the heteroaryl, aryl, heteroarylalkyl, arylalkoxy or arylalkyloxycarbonylalkyl are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from
- 5 hydroxy, oxo, alkyl, amino, hydroxyalkyl, alkylcarbonyl, alkoxycarbonyl, halo, aminoalkyl, alkylaminoalkyl, and dialkylaminoalkyl; and

- 5 R^{35} is selected from halo, $-(CH_2)_pC(O)OR_{17}$, cycloalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the heterocycloalkyl and heterocycloalkylalkyl are each optionally substituted with 1, 2, 3, 4, or 5 groups each independently selected from alkyl, alkoxy, and halo.

[0012] Another embodiment of the first aspect of the invention relates to a compound of


10 Formula II:



wherein E, D, L, Z, R^1 , R^2 and R^{25} are as defined above for the compound of Formula I.

[0013] Another embodiment of the first aspect of the invention relates to a compound of


5 Formula III:



wherein E, D, L, Z, R^1 , R^2 and R^{25} are as defined above for the compound of Formula I.

[0014] Another embodiment of the first aspect of the invention relates to a compound of


10 Formula IV:



wherein D, E, R^{25} and R^{32} are as defined above for Formula I, and R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the


- 5 heterocycloalkyl is optionally substituted with one or two R^{31} , and wherein R^{31} is as defined above in Formula I.

[0015] Another embodiment of the first aspect of the invention relates to a compound of Formula V:



10 wherein D, E, R^{25} and R^{32} are as defined above for Formula I, and R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or two R^{31} , and wherein R^{31} is as defined above in Formula I.

- 15 [0016] Another embodiment of the first aspect of the invention relates to a compound of Formula VI:




20 wherein D, E, R^{25} and R^{32} are as defined above for Formula I, and R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or two R^{31} , and wherein R^{31} is as defined above in Formula I.

[0017] In other embodiments of the first aspect of the invention, D, E and R^{25} for Formula IV, Formula V or Formula VI are each hydrogen.

- 25 [0018] In other embodiments of the first aspect of the invention, R^{32} for Formula IV, Formula V or Formula VI is heterocycloalkyl.


- 5 [0019] In other embodiments of the first aspect of the invention, R^{32} for Formula IV, Formula V or Formula VI is alkyl optionally substituted with alkoxy, hydroxy, amino, alkylamino, or dialkylamino.

[0020] I In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




10 wherein R^{27a} , R^{11} and $n2$ are as defined above for the compound of Formula I.

[0021] In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




- 15 wherein R^{28} , R^{11} and $n2$ are as defined above for the compound of Formula I, and R^{28a} is arylalkyl or heteroarylalkyl, wherein the arylalkyl or heteroarylalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents selected from halo or lower alkyl.

[0022] In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




20 wherein R^{28} , R^{28a} , R^{11} and $n2$ are as defined above for the compound of Formula I.

[0023] In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




- 5 wherein R^{28} , R^{11} and n_2 are as defined above for the compound of Formula I, and R^{28a} is selected from lower alkyl, dialkylaminoalkyl, alkoxyalkyl, arylalkyl, heteroarylalkyl, and heterocycloalkylalkyl.

[0024] In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




wherein R^{11} and n_2 are as defined above for the compound of Formula I, and R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, join together to form a ring structure selected from thiazolidinyl, piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyrimidinyl, and pyridinyl, wherein the ring structure is optionally substituted with 1, 2, 3, 4 or 5 substituents selected from halo, lower alkyl or alkoxy.

[0025] In other embodiments of the first aspect of the invention, R^2 in Formula I, II or III is




wherein R^{27a} , R^{11} and n_2 are as defined above for the compound of Formula I.

[0026] Other embodiments of the first aspect of the invention relate to a compound of




Formula I, II or III, wherein L is a bond, and Z is

[0027] Other embodiments of the first aspect of the invention relate to a compound of



Formula I, II or III, wherein Z is and R^{25} is hydrogen.

[0028] Other embodiment of the first aspect of the invention relate to a compound of



Formula I, II or III, wherein Z is, R^{25} is hydrogen and E and D are hydrogen.

- 5 [0029] Other embodiments of the first aspect of the invention relate to a compound of in Formula I, II or III, wherein R^{25} is on the 3 position.

[0030] Other embodiment of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, and R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$.

[0031] Other embodiments of the first aspect of the invention relate to compound of

- 10 Formula I, II or III, wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$, and R^{32} is selected from lower alkyl, cycloalkyl, diaminoalkyl, aminoalkyl, arylalkyl, heterocycloalkyl, alkoxyalkyl, alkylamino, and hydroxyalkyl optionally substituted with amino.

[0032] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$, and R^{32} is cycloalkyl.

- 15 [0033] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$, and R^{32} is lower alkyl.


[0034] Other embodiments of the first aspect of the invention relate to a compound of

- Formula I, II or III wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$, R^{26} is hydrogen, wherein R^{32} selected from aryl, arylalkyl, cycloalkyl, alkoxyalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl, wherein R^{32} optionally substituted with 1, 2, 3, 4 or 5 groups selected from hydroxyl, oxo, alkyl, alkoxy, amino, hydroxyalkyl or halo.
- 20


[0035] Other embodiments of the first aspect of the invention relate to a compound of

- Formula I, II or III, wherein Z is $\begin{array}{c} \text{S} \text{---} \text{N} \text{---} \text{R}^{26} \\ | \\ \text{R}^{26a} \end{array}$, R^{26a} is $-\text{C}(\text{O})\text{R}^{32}$, R^{26} is hydrogen, wherein R^{32} selected from tetrahydrofuran, pyrrolidinyl or pyrimidinyl, wherein R^{32} optionally substituted with 1, 2, 3, 4 or 5 groups selected from hydroxyl, oxo, alkyl, alkoxy, amino, hydroxyalkyl or halo.
- 25

- 5 [0036] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein Z is , R^{26a} is $-C(O)R^{32}$, R^{26} is hydrogen, wherein R^{32} is lower alkyl optionally substituted with 1, 2, 3, 4 or 5 groups selected from dialkylaminocarbonyl, hydroxy and $-NR^{34}R^{34a}$, wherein R^{34} and R^{34a} are as defined above for Formula I.

- 10 [0037] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein R^2 is .

[0038] In another embodiment of the first aspect of the invention, R^{32} is methyl.

[0039] In another embodiment of the first aspect of the invention, R^{32} is alkyl substituted with $-NR^{34}R^{34a}$.

- 15 [0040] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, wherein R^{32} is U or $-CH_2-U$, wherein U is selected from pyrrolidinyl, thiazolidinyl, morpholinyl, azetidyl, cyclobutyl, cyclopropyl, tetrahydrofuranyl, pyrazinyl, imidazolyl, piperazinyl, thienyl, thienylmethyl, furanyl, phenyl, prolinamidyl, pyridinyl, tetrahydronaphthalene, tetrazolyl, isoindolinyl, pyranyl, cyclopentyl, and octahydro-1H-indolyl.


[0041] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, wherein R^{11} , when present, is halo or lower alkyl.

[0042] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, wherein R^{11} , when present, is lower alkyl.


- 25 [0043] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, wherein R^{35} is heterocycloalkylalkyl, wherein the heterocycloalkyl is selected from piperazinyl, piperidinyl, morpholinyl and dioxanyl.

[0044] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, wherein n_2 is 0.

- 30 [0045] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein R^2 is .

- 5 [0046] Other embodiments of the first aspect of the invention relate to a compound of

Formula I, II or III, wherein R^2 is , and wherein R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl.

[0047] Other embodiments of the first aspect of the invention relate to a compound of Formula I, II or III, IV or V, wherein R^{25} is hydrogen.

- 10 [0048] A second aspect of the invention relates to a pharmaceutical composition comprising a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, excipient, or diluent.

[0049] A third aspect of the invention relates to a method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a compound of
15 Formula I, II, III, IV, V or VI, or a pharmaceutically salt thereof.

[0050] A fourth aspect of the invention relates to a method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a pharmaceutical composition comprising a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof.

- 20 [0051] A fifth aspect of the invention relates to a method for treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a therapeutically effective amount of a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof.

[0052] A sixth aspect of the invention relates to a method for treating a disease, disorder,
25 or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof.

[0053] The disease being treated in these aspects of the invention can be a
30 myeloproliferative disorder, cancer, cardiovascular disease, and/or hematopoietic abnormality where hyperactivation of JAK-STAT signaling is present. Nonlimiting examples of myeloproliferative disorders that are contemplated as being treatable by the compounds of the invention include myelofibrosis, thrombocythemia, polycythemia vera (PV), essential thrombocythemia (ET), agnogenic myeloid metaplasia (AMM), also referred to as idiopathic myelofibrosis (IMF), and chronic myelogenous leukemia (CML). Nonlimiting examples of
35 cancers that are contemplated as being treatable by the compounds of the invention include

- 5 leukemias, lymphomas, multiple myeloma, prostate cancers, lung cancers, breast cancers, and ovarian cancers. Nonlimiting examples of cardiovascular diseases that are contemplated as being treatable by the compounds of the invention include congestive heart failure and hypertension. Nonlimiting examples of hematopoietic abnormalities that are contemplated as being treatable by the compounds of the invention include thrombocytosis.
- 10 **[0054]** A seventh aspect of the invention relates to a method of treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof, in combination with one or more treatment(s) selected from surgery, one or more therapeutic agent(s), plateletpheresis, and radiation.
- 15 **[0055]** An eighth aspect of the invention relates to a method of treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I, II, III, IV, V or VI, or a pharmaceutically acceptable salt thereof, in combination with one or more treatment(s) selected from surgery, 20 one or more therapeutic agent(s), plateletpheresis, and radiation.
- [0056]** When treating myeloproliferative disorders, the compound of Formula I, II, III, IV, V or VI can also be administered with one or more additional treatment(s) selected from plateletpheresis and one or more therapeutic agent(s) selected from interferon- α ; aspirin; a platelet-decreasing drug, such as anagrelide; and a myelosuppressive agent (such as a 25 radiophosphorus and alkylating agents). Non-limiting examples of the myelosuppressive agent include hydroxyurea, melphalan, and busulfan.
- [0057]** When treating cancer, the compound of Formula I, II, III, IV, V or VI can be administered in combination with one or more chemotherapeutic agent(s) selected from fludarabine, vinblastine, adriamycin and cisplatin.

10 **Abbreviations and Definitions**

[0058] The following abbreviations and terms have the indicated meanings throughout:

Abbreviation	Meaning
Ac	Acetyl
Br	Broad
°C	degrees Celsius
c-	Cyclo

Abbreviation	Meaning
CBZ	CarboBenZoxy = benzyloxycarbonyl
D	doublet
Dd	doublet of doublet
Dt	doublet of triplet
DIPEA	<i>N,N</i> -diisopropylethylamine
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethyl sulfoxide
EI	Electron Impact ionization
Et	Ethyl
G	gram(s)
GC	gas chromatography
h or hr	hour(s)
HOAc	acetic acid
HOBt	Hydroxybenzotriazole
HPLC	high pressure liquid chromatography
L	liter(s)
M	molar or molarity
M	Multiplet
Me	Methyl
Mesyl	Methanesulfonyl
Mg	milligram(s)
MHz	megahertz (frequency)
Min	minute(s)
mL	milliliter(s)
mM	Millimolar
Mmol	millimole(s)
Mol	mole(s)
MS	mass spectral analysis
MTBE	methyl t-butyl ether
N	normal or normality

Abbreviation	Meaning
NBS	<i>N</i> -bromosuccinimide
NCS	<i>N</i> -chlorosuccinimide
nM	Nanomolar
NMO	<i>N</i> -methylmorpholine oxide
NMR	nuclear magnetic resonance spectroscopy
PEG	polyethylene glycol
pEY	poly-glutamine, tyrosine
Ph	Phenyl
PhOH	Phenol
PfP	Pentafluorophenol
PfPy	Pentafluoropyridine
PPTS	Pyridinium <i>p</i> -toluenesulfonate
Py	Pyridine
PyBroP	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
Q	Quartet
RT	Room temperature
Sat'd	Saturated
S	Singlet
s-	Secondary
t-	Tertiary
t or tr	Triplet
TBDMS	<i>t</i> -butyldimethylsilyl
TES	Triethylsilyl
TFA	trifluoroacetic acid
THF	Tetrahydrofuran
TMOF	trimethyl orthoformate
TMS	Trimethylsilyl
Tosyl	<i>p</i> -toluenesulfonyl
Trt	triphenylmethyl

Abbreviation	Meaning
uL	microliter(s)
uM	Micromole(s) or micromolar


uL	microliter(s)
uM	Micromole(s) or micromolar

5


[0059] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise or they are expressly defined to mean something different.

10 [0060] The symbol “-” means a single bond, “=” means a double bond, “≡” means a triple bond, “——” means a single or double bond. When a group is depicted removed from its parent formula, the “~” symbol will be used at the end of the bond which was theoretically cleaved in order to separate the group from its parent structural formula.

15 [0061] When chemical structures are depicted or described, unless explicitly stated otherwise, all carbons are assumed to have hydrogen substitution to conform to a valence of four. For example, in the structure on the left-hand side of the schematic below there are nine hydrogens implied. The nine hydrogens are depicted in the right-hand structure. Sometimes a particular atom in a structure is described in textual formula as having a hydrogen or hydrogens as substitution (expressly defined hydrogen), for example, -CH₂CH₂-. It is
 20 understood by one of ordinary skill in the art that the aforementioned descriptive techniques are common in the chemical arts to provide brevity and simplicity to description of otherwise complex structures.




25 [0062] If a group “R” is depicted as “floating” on a ring system, as for example in the formula:




then, unless otherwise defined, a substituent “R” can reside on any atom of the ring system, assuming replacement of a depicted, implied, or expressly defined hydrogen from one of the
 30 ring atoms, so long as a stable structure is formed.

- 5 **[0063]** If a group “R” is depicted as floating on a fused ring system, as for example in the formulae:




- 10 then, unless otherwise defined, a substituent “R” can reside on any atom of the fused ring system, assuming replacement of a depicted hydrogen (for example the -NH- in the formula above), implied hydrogen (for example as in the formula above, where the hydrogens are not shown but understood to be present), or expressly defined hydrogen (for example where in the formula above, “X” equals =CH-) from one of the ring atoms, so long as a stable structure is formed. In the example depicted, the “R” group can reside on either the 5-membered or the 6-membered ring of the fused ring system. In the formula depicted above, when y is 2 for example, then the two “R’s” can reside on any two atoms of the ring system, again assuming
- 15 each replaces a depicted, implied, or expressly defined hydrogen on the ring.

[0064] When a group “R” is depicted as existing on a ring system containing saturated carbons, as for example in the formula:



- 20 where, in this example, “y” can be more than one, assuming each replaces a currently depicted, implied, or expressly defined hydrogen on the ring; then, unless otherwise defined, where the resulting structure is stable, two “R’s” can reside on the same carbon. A simple example is when R is a methyl group; there can exist a geminal dimethyl on a carbon of the depicted ring (an “annular” carbon). In another example, two R’s on the same carbon, including that carbon, can form a ring, thus creating a spirocyclic ring (a “spirocyclyl” group)
- 25 structure with the depicted ring as for example in the formula:



- [0065]** “Administration” and variants thereof (e.g., “administering” a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., surgery, radiation, and chemotherapy, etc.), “administration” and its variants are each
- 30

5 understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

[0066] "Alkyl" is intended to include C₁-C₂₀, more typically, C₁-C₁₂ linear or branched structures and combinations thereof, inclusively. "Lower alkyl" is intended to include C₁-C₆ linear or branched structures and combinations thereof, inclusively. For example, "C₆ alkyl" can refer to an *n*-hexyl, *iso*-hexyl, cyclobutylethyl, and the like. Examples of lower alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *s*-butyl, *t*-butyl, isobutyl, pentyl, hexyl and the like. Higher alkyl refers to alkyl groups containing more than six carbon atoms. In this application, alkyl refers to alkanyl, alkenyl, and alkynyl residues (and combinations thereof); it is intended to include cyclohexylmethyl, vinyl, allyl, isoprenyl, and the like. Thus when an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, either "butyl" or "C₄ alkyl" is meant to include *n*-butyl, *sec*-butyl, isobutyl, *t*-butyl, isobutenyl and but-2-ynyl groups; and for example, "propyl" or "C₃ alkyl" each include *n*-propyl, propenyl, and isopropyl.

20 [0067] "Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 14 carbon atoms, 5 to 10 carbon atoms, or 5 to about 7 ring atoms. Non-limiting examples of monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of multicyclic cycloalkyls include 1-decalin, norbornyl, adamantyl and the like. Cycloalkyls can be fused or bridge ring systems or spirocyclic systems.

[0068] "Alkyl substituted with halo and hydroxy" means an alkyl group substituted with 1, 2, or 3 hydroxy and 1, 2, 3, 4, or 5 halo.

[0069] "Alkylene" refers to straight or branched chain divalent group consisting solely of carbon and hydrogen atoms, containing no unsaturation and having from one to ten carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene and the like. Alkylene is a subset of alkyl, referring to the same residues as alkyl, but having two points of attachment and, specifically, fully saturated. Examples of alkylene include ethylene (-CH₂CH₂-), propylene (-CH₂CH₂CH₂-), dimethylpropylene (-CH₂C(CH₃)₂CH₂-), and cyclohexylpropylene (-CH₂CH₂CH(C₆H₁₁)).

5 [0070] "Alkylidene" refers to a straight or branched chain unsaturated divalent group consisting solely of carbon and hydrogen atoms, having from two to ten carbon atoms, for example, ethylidene, propylidene, *n*-butylidene, and the like. Alkylidene is a subset of alkyl,

5 referring to the same residues as alkyl, but having two points of attachment and, specifically, double bond unsaturation. The unsaturation present includes at least one double bond.

[0071] "Alkylidyne" refers to a straight or branched chain unsaturated divalent group consisting solely of carbon and hydrogen atoms having from two to ten carbon atoms, for example, propylid-2-ynyl, *n*-butylid-1-ynyl, and the like. Alkylidyne is a subset of alkyl,
10 referring to the same residues as alkyl, but having two points of attachment and, specifically, triple bond unsaturation. The unsaturation present includes at least one triple bond.

[0072] Any of the above groups, "alkylene," "alkylidene" and "alkylidyne," when optionally substituted, can contain alkyl substitution which itself contains unsaturation. For example, 2-(2-phenylethynyl-but-3-enyl)-naphthalene (IUPAC name) contains an *n*-butylid-
15 3-ynyl group with a vinyl substituent at the 2-position of said group.

[0073] "Alkoxy" or "alkoxyl" refers to the group -O-alkyl, wherein the term "alkyl" is as defined hereinabove. Examples include methoxy, ethoxy, propoxy, isopropoxy, and the like. Lower alkoxy refers to groups containing one to six carbons.

[0074] "Substituted alkoxy" refers to the group -O-(substituted alkyl), the substitution on
20 the alkyl group generally containing more than only carbon (as defined by alkoxy). Another exemplary substituted alkoxy group is hydroxyalkoxy or -O-alkyl-OH.

[0075] "Aryl" means a monovalent five- to fourteen-membered mono- or multicyclic ring, wherein the monocyclic ring is aromatic and at least one of the rings in the multicyclic ring is aromatic. An aryl can also be five- to ten membered, or six membered. Representative non-
25 limiting examples of aryl include phenyl, naphthyl, and the like.

[0076] "Arylalkyl" means a residue in which an aryl moiety, as defined above, is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne group. Examples include benzyl, phenethyl, phenylvinyl, phenylallyl and the like. The "alkyl" portion of the group can be one to ten carbons, and in another embodiment, one to six carbons; the latter
30 can also be referred to as C₁₋₆ arylalkyl. When a group is referred to as or "-(C₁-C₆)alkylaryl," an aryl moiety is attached to a parent structure via an alkylene group. Examples include benzyl, phenethyl, and the like.

[0077] In some examples, as appreciated by one of ordinary skill in the art, two adjacent groups on an aromatic system can be fused together to form a ring structure. The fused ring
35 structure can contain heteroatoms and can be optionally substituted with one or more groups. It should additionally be noted that saturated carbons of such fused groups (*i.e.* saturated ring structures) can contain two substitution groups.

5 [0078] "Fused-polycyclic" or "fused ring system" refers to a polycyclic ring system that contains bridged or fused rings; that is, where two rings have more than one shared atom in their ring structures. In this application, fused-polycyclics and fused ring systems includes non-aromatic and aromatic systems. Typically, but not necessarily, fused-polycyclics share a vicinal set of atoms, for example naphthalene or 1,2,3,4-tetrahydro-naphthalene. A spiro ring
10 system is not a fused-polycyclic by this definition, but fused polycyclic ring systems of the invention can themselves have spiro rings attached thereto via a single ring atom of the fused-polycyclic.

[0079] "Halogen" or "halo" refers to fluorine, chlorine, bromine or iodine. "Haloalkyl" and "haloaryl" refer generically to alkyl and aryl groups that are substituted with one or more
15 halogens, respectively. Non-limiting examples of "haloalkyl" include -CH₂F, -CHCl₂ or -CF₃.

[0080] "Heteroatom" refers to O, S, N, or P.

[0081] "Heterocyclyl" refers to a stable three- to fifteen-membered ring substituent that consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, phosphorus, oxygen and sulfur. For purposes of this invention, the heterocyclyl
20 substituent can be a monocyclic, bicyclic or tricyclic ring system, which can include fused or bridged ring systems as well as spirocyclic systems. The terms "heterocycloalkyl" and "heteroaryl" are groups that are encompassed by the broader term "heterocyclyl." The nitrogen, phosphorus, carbon or sulfur atoms in the heterocyclyl group can be optionally oxidized to various oxidation states. In a specific example, the group -S(O)₀₋₂-, refers to -S-
25 (sulfide), -S(O)- (sulfoxide), and -SO₂- (sulfone) respectively. For convenience, nitrogens, particularly but not exclusively, those defined as annular aromatic nitrogens, are meant to include their corresponding *N*-oxide form, although not explicitly defined as such in a particular example. Thus, for a compound of the invention having, for example, a pyridyl ring; the corresponding pyridyl-*N*-oxide is meant to be included as another compound of the
30 invention. In addition, annular nitrogen atoms can be optionally quaternized; and the ring substituent can be partially or fully saturated or aromatic. Examples of heterocyclyl groups include, but are not limited to, azetidiny, acridiny, benzodioxolyl, benzodioxanyl, benzofuranyl, carbazoyl, cinnoliny, dioxolanyl, indoliziny, naphthyridiny, perhydroazepiny, phenaziny, phenothiaziny, phenoxaziny, phthalaziny, pteridiny,
35 puriny, quinazoliny, quinoxaliny, quinoliny, isoquinoliny, tetrazoyl, tetrahydroisoquinolyl, piperidiny, piperaziny, 2-oxopiperaziny, 2-oxopiperidiny, 2-oxopyrrolidiny, 2-oxoazepiny, azepiny, pyrroly, 4-piperidonyl, pyrrolidiny, pyrazoly,

- 5 pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, dihydropyridinyl, tetrahydropyridinyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazoliny, oxazolidinyl, triazolyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolinyl, thiazolidinyl, isothiazolyl, quinuclidinyl, isothiazolidinyl, indolyl, isoindolyl, indolinyl, isoindolinyl, octahydroindolyl, octahydroisoindolyl, quinolyl, isoquinolyl,
- 10 decahydroisoquinolyl, benzimidazolyl, thiadiazolyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, dioxaphospholanyl, oxadiazolyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, and tetrahydroquinolinyl.

[0082] "Heterocycloalkyl" refers to a stable 4-12 membered monocyclic, bicyclic or

15 tricyclic ring containing one or more heteroatoms.

[0083] "Heterocycloalkylalkyl" refers to a heterocycloalkyl, as defined herein, attached to the parent moiety through an "alkyl," as defined herein. One non-limiting example of heterocycloalkyl includes piperadinyl. Another non-limiting example of heterocycloalkyl includes piperadinyl. Another non-limiting example of heterocycloalkyl includes

20 piperazinyl. Another non-limiting example of heterocycloalkyl includes furanyl. Another non-limiting example of heterocycloalkyl includes pyrrolidinyl. Another non-limiting example of heterocycloalkyl includes morpholinyl.

[0084] "amino" refers to -NH_2 .

[0085] "alkylamino" refers to -NH(alkyl) , wherein "alkyl" is as defined above, and

25 wherein the the parent moiety is attached to the nitrogen atom.

[0086] "dialkylamino" refers to -N(alkyl)_2 , wherein "alkyl" is as defined above, and wherein the parent moiety is attached to the nitrogen atom.

[0087] "dialkylaminoalkyl" refers to $\text{-(alkyl)N(alkyl)}_2$, wherein "alkyl" is as defined above. One such non-limiting example of "dialkylaminoalkyl" includes -

30 $\text{CH}_2\text{C(CH}_3)_2\text{CH}_2\text{N(CH}_3)_2$.

[0088] "aminoalkyl" refers to -(alkyl)NH , wherein "alkyl" is as defined above, and wherein the the parent moiety is attached to the alkyl group.

[0089] "aminoalkyl" refers to -(alkyl)NH_2 , wherein "alkyl" is as defined above, and wherein the the parent moiety is attached to the alkyl group. The amino group can be attached

35 at any point along the alkyl group. Non-limiting examples of aminoalkyl include - $\text{CH(NH}_2\text{)CH}_3$,

5 [0090] "Phenoxy" refers to a -alkyl-O-phenyl group, wherein "alkyl" is as defined above, and the parent moiety is attached to the alkyl group.

[0091] "Heteroaryl" means a 5- to 12-membered, monocyclic aromatic heterocyclyl (where heterocyclyl is defined herein) or bicyclic heterocyclyl ring system (where at least one of the rings in the bicyclic system is aromatic) where the monocyclic ring and at least one of
10 the rings in the bicyclic ring system contains one, two, three, four, or five heteroatom(s) selected from nitrogen, oxygen, phosphorous, and sulfur. The ring containing the heteroatom can be aromatic or non-aromatic. Representative examples include pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinoliny, isoquinoliny, indolyl, benzimidazolyl,
15 benzdioxolyl, benzofuranyl, cinnoliny, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridiny, and furopyridiny. Fused, bridged, and spiro moieties are also included within the scope of this definition.

20 [0092] "Carbonyl" refers to the group "-C(O)-", which is bivalent.

[0093] "Aminocarbonyl" refers to the group "-C(O)-NH₂," wherein the parent moiety is attached to the amino group.

[0094] "Alkoxycarbonyl" refers to the group "-C(O)alkoxy," wherein alkoxy is as defined above, and the parent moiety is attached to the carbonyl. A non-limiting example
25 includes -C(O)-OC(CH₃)₃.

[0095] When a group is referred to as "-C₁-C₆ alkyl heterocyclyl" the heterocyclyl is attached to a parent structure via one of an alkylene, alkylidene, or alkylidyne group. Examples include (4-methylpiperazin-1-yl) methyl, (morpholin-4-yl) methyl, (pyridine-4-yl) methyl, 2-(oxazolin-2-yl) ethyl, 4-(4-methylpiperazin-1-yl)-2-butenyl, and the like. Both the
30 heterocyclyl and the corresponding alkylene, alkylidene, or alkylidyne portion of a heterocyclylalkyl group can be optionally substituted.


[0096] "Hydroxyalkyl" means -alkyl-OH, wherein alkyl is as defined hereinabove.

[0097] "Optional" or "optionally" means that the subsequently described event or circumstance can or can not occur, and that the description includes instances where said
35 event or circumstance occurs and instances in which it does not. One of ordinary skill in the art would understand that with respect to any molecule described as containing one or more optional substituents, only sterically practical and/or synthetically feasible compounds are

5 meant to be included. "Optionally substituted" refers to all subsequent modifiers in a term. So, for example, in the term "optionally substituted arylalkyl," both the "alkyl" portion and the "aryl" portion of the molecule can or can not be substituted. A list of exemplary optional substitutions is presented below in the definition of "substituted."

10 [0098] "Saturated bridged ring system" refers to a bicyclic or polycyclic ring system that is not aromatic. Such a system can contain isolated or conjugated unsaturation, but not aromatic or heteroaromatic rings in its core structure (but can have aromatic substitution thereon). For example, hexahydro-furo[3,2-b]furan, 2,3,3a,4,7,7a-hexahydro-1H-indene, 7-aza-bicyclo[2.2.1]-heptane, and 1,2,3,4,4a,5,8,8a-octahydro-naphthalene are all included in the class "saturated bridged ring system."

15 [0099] "Spirocyclyl" or "spirocyclic ring" refers to a ring originating from a particular annular carbon of another ring. For example, as depicted below, a ring atom of a saturated bridged ring system (rings B and B'), but not a bridgehead atom, can be a shared atom between the saturated bridged ring system and a spirocyclyl (ring A) attached thereto. A spirocyclyl can be carbocyclic or heteroalicyclic.



20

[0100] "Substituted" alkyl, aryl, and heterocyclyl, refer respectively to alkyl, aryl, and heterocyclyl, one or more (for example up to about five, in another example, up to about three) hydrogen atoms are replaced by a substituent independently selected from: alkyl (for example, fluoromethyl), aryl (for example, 4-hydroxyphenyl), arylalkyl (for example, 1-phenyl-ethyl), heterocyclylalkyl (for example, 1-pyridin-3-yl-ethyl), heterocyclyl (for example, 5-chloro-pyridin-3-yl or 1-methyl-piperidin-4-yl), alkoxy, alkylenedioxy (for example methylenedioxy), amino (for example, alkylamino and dialkylamino), amidino, aryloxy (for example, phenoxy), arylalkyloxy (for example, benzyloxy), carboxy (-CO₂H), carboalkoxy (that is, acyloxy or -OC(=O)R), carboxyalkyl (that is, esters or -CO₂R),
25 carboxamido, benzyloxycarbonylamino (CBZ-amino), cyano, acyl, halogen, hydroxy, nitro, sulfanyl, sulfinyl, sulfonyl, thiol, halogen, hydroxy, oxo, carbamyl, acylamino, and sulfonamido. And each substituent of a substituted group is optionally substituted, but these optional substituents themselves are not further substituted. Thus, an optionally substituted moiety is one that can or can not have one or more substituents, and each of the substituents
30

5 can or can not have one or more substituents. But, the substituents of the substituents can not be substituted.

[0101] Some of the compounds of the invention can have imino, amino, oxo or hydroxy substituents off aromatic heterocyclyl systems. For purposes of this disclosure, it is understood that such imino, amino, oxo or hydroxy substituents can exist in their
10 corresponding tautomeric form, *i.e.*, amino, imino, hydroxy or oxo, respectively.

[0102] "Patient" for the purposes of the present invention includes humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In a preferred embodiment the patient is a mammal, and in a most preferred embodiment the patient is human.

15 [0103] "Kinase-dependent diseases or conditions" refer to pathologic conditions that depend on the activity of one or more protein kinases. Kinases either directly or indirectly participate in the signal transduction pathways of a variety of cellular activities including proliferation, adhesion, migration, differentiation and invasion. Diseases associated with kinase activities include tumor growth, the pathologic neovascularization that supports solid
20 tumor growth, and associated with other diseases where excessive local vascularization is involved such as ocular diseases (diabetic retinopathy, age-related macular degeneration, and the like) and inflammation (psoriasis, rheumatoid arthritis, and the like).

[0104] While not wishing to be bound to theory, phosphatases can also play a role in "kinase-dependent diseases or conditions" as cognates of kinases; that is, kinases
25 phosphorylate and phosphatases dephosphorylate, for example protein substrates. Therefore compounds of the invention, while modulating kinase activity as described herein, can also modulate, either directly or indirectly, phosphatase activity. This additional modulation, if present, can be synergistic (or not) to activity of compounds of the invention toward a related or otherwise interdependent kinase or kinase family. In any case, as stated previously, the
30 compounds of the invention are useful for treating diseases characterized in part by abnormal levels of cell proliferation (*i.e.* tumor growth), programmed cell death (apoptosis), cell migration and invasion and angiogenesis associated with tumor growth.

[0105] "Therapeutically effective amount" is an amount of a compound of the invention, that when administered to a patient, ameliorates a symptom of the disease. The amount of a
35 compound of the invention which constitutes a "therapeutically effective amount" will vary depending on the compound, the disease state and its severity, the age of the patient to be

5 treated, and the like. The therapeutically effective amount can be determined routinely by one of ordinary skill in the art having regard to their knowledge and to this disclosure.

[0106] "Cancer" refers to cellular-proliferative disease states, including but not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocyoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiforme, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell

5 carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma]; Skin: malignant melanoma, basal cell carcinoma, squamous cell
10 carcinoma, Karposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

[0107] A "pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the
15 parent compound. It is understood that the pharmaceutically acceptable salts are non-toxic. Additional information on suitable pharmaceutically acceptable salts can be found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, which is incorporated herein by reference or S. M. Berge, et al., "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977;66:1-19 both of which are incorporated herein by reference.

20 [0108] Examples of pharmaceutically acceptable acid addition salts include those formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; as well as organic acids such as acetic acid, trifluoroacetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric
25 acid, benzoic acid, cinnamic acid, 3-(4-hydroxybenzoyl)benzoic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid,
30 trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, p-toluenesulfonic acid, and salicylic acid and the like.

[0109] Examples of a pharmaceutically acceptable base addition salts include those formed when an acidic proton present in the parent compound is replaced by a metal ion,
35 such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferable salts are the ammonium, potassium, sodium, calcium, and magnesium salts. Salts derived from pharmaceutically acceptable

5 organic non-toxic bases include, but are not limited to, salts of primary, secondary, and
tertiary amines, substituted amines including naturally occurring substituted amines, cyclic
amines and basic ion exchange resins. Examples of organic bases include isopropylamine,
trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-
dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine,
10 caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine,
methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine,
tromethamine, N-methylglucamine, polyamine resins, and the like. Exemplary organic bases
are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine,
choline, and caffeine.

15 [0110] "Prodrug" refers to compounds that are transformed (typically rapidly) *in vivo* to
yield the parent compound of the above formulae, for example, by hydrolysis in blood.
Common examples include, but are not limited to, ester and amide forms of a compound
having an active form bearing a carboxylic acid moiety. Examples of pharmaceutically
acceptable esters of the compounds of this invention include, but are not limited to, alkyl
20 esters (for example with between about one and about six carbons) the alkyl group is a
straight or branched chain. Acceptable esters also include cycloalkyl esters and arylalkyl
esters such as, but not limited to benzyl. Examples of pharmaceutically acceptable amides of
the compounds of this invention include, but are not limited to, primary amides, and
secondary and tertiary alkyl amides (for example with between about one and about six
25 carbons). Amides and esters of the compounds of the present invention can be prepared
according to conventional methods. A thorough discussion of prodrugs is provided in T.
Higuchi and V. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S.
Symposium Series, and in Bioreversible Carriers in Drug Design, ed. Edward B. Roche,
American Pharmaceutical Association and Pergamon Press, 1987, both of which are
30 incorporated herein by reference for all purposes.

[0111] "Metabolite" refers to the break-down or end product of a compound or its salt
produced by metabolism or biotransformation in the animal or human body; for example,
biotransformation to a more polar molecule such as by oxidation, reduction, or hydrolysis, or
to a conjugate (see Goodman and Gilman, "The Pharmacological Basis of Therapeutics"
35 8.sup.th Ed., Pergamon Press, Gilman et al. (eds), 1990 for a discussion of
biotransformation). As used herein, the metabolite of a compound of the invention or its salt
can be the biologically active form of the compound in the body. In one example, a prodrug

5 can be used such that the biologically active form, a metabolite, is released *in vivo*. In another example, a biologically active metabolite is discovered serendipitously, that is, no prodrug design *per se* was undertaken. An assay for activity of a metabolite of a compound of the present invention is known to one of skill in the art in light of the present disclosure.

[0112] The present invention also includes N-oxide derivatives and protected derivatives
10 of compounds of Formula I. For example, when compounds of Formula I contain an oxidizable nitrogen atom, the nitrogen atom can be converted to an N-oxide by methods well known in the art. When compounds of Formula I contain groups such as hydroxy, carboxy, thiol or any group containing a nitrogen atom(s), these groups can be protected with a suitable "protecting group" or "protective group". A comprehensive list of suitable protective
15 groups can be found in T.W. Greene, *Protective Groups in Organic Synthesis*, John Wiley & Sons, Inc. 1991, the disclosure of which is incorporated herein by reference in its entirety. The protected derivatives of compounds of Formula I can be prepared by methods well known in the art.

[0113] "Treating" or "treatment" of a disease, disorder, or syndrome, as used herein,
20 includes (i) preventing the disease, disorder, or syndrome from occurring in a human, i.e. causing the clinical symptoms of the disease, disorder, or syndrome not to develop in an animal that can be exposed to or predisposed to the disease, disorder, or syndrome but does not yet experience or display symptoms of the disease, disorder, or syndrome; (ii) inhibiting the disease, disorder, or syndrome, i.e., arresting its development; and (iii) relieving the
25 disease, disorder, or syndrome, i.e., causing regression of the disease, disorder, or syndrome. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition can be necessary, and will be ascertainable with routine experimentation by one of ordinary skill in the art.

0 [0114] One of ordinary skill in the art would understand that certain crystallized, protein-ligand complexes, in particular JAK-2-ligand complexes, and their corresponding x-ray structure coordinates can be used to reveal new structural information useful for understanding the biological activity of kinases as described herein. As well, the key structural features of the aforementioned proteins, particularly, the shape of the ligand
5 binding site, are useful in methods for designing or identifying selective modulators of kinases and in solving the structures of other proteins with similar features. Such protein-

5 ligand complexes, having compounds of the invention as their ligand component, are an aspect of the invention.

[0115] As well, one of ordinary skill in the art would appreciate that such suitable x-ray quality crystals can be used as part of a method of identifying a candidate agent capable of binding to and modulating the activity of kinases. Such methods can be characterized by the following aspects: a) introducing into a suitable computer program, information defining a ligand binding domain of a kinase in a conformation (*e.g.* as defined by x-ray structure coordinates obtained from suitable x-ray quality crystals as described above) wherein the computer program creates a model of the three dimensional structures of the ligand binding domain, b) introducing a model of the three dimensional structure of a candidate agent in the computer program, c) superimposing the model of the candidate agent on the model of the ligand binding domain, and d) assessing whether the candidate agent model fits spatially into the ligand binding domain. Aspects a-d are not necessarily carried out in the aforementioned order. Such methods can further entail: performing rational drug design with the model of the three-dimensional structure, and selecting a potential candidate agent in conjunction with computer modeling.

[0116] Additionally, one skilled in the art would appreciate that such methods can further entail: employing a candidate agent, so-determined to fit spatially into the ligand binding domain, in a biological activity assay for kinase modulation, and determining whether said candidate agent modulates kinase activity in the assay. Such methods can also include administering the candidate agent, determined to modulate kinase activity, to a mammal suffering from a condition treatable by kinase modulation, such as those described above.





[0117] Also, one skilled in the art would appreciate that compounds of the invention can be used in a method of evaluating the ability of a test agent to associate with a molecule or molecular complex comprising a ligand binding domain of a kinase. Such a method can be characterized by the following aspects: a) creating a computer model of a kinase binding pocket using structure coordinates obtained from suitable x-ray quality crystals of the kinase, b) employing computational algorithms to perform a fitting operation between the test agent and the computer model of the binding pocket, and c) analyzing the results of the fitting operation to quantify the association between the test agent and the computer model of the binding pocket.








[0118] In addition to the preferred embodiments recited hereinabove, also preferred are embodiments comprising combinations of preferred embodiments.








- 5 [0119] Representative compounds of Formula I and/or II are depicted below. The examples are merely illustrative and do not limit the scope of the invention in any way. The compounds in Table 1(Part A and Part B) below can be prepared using art recognized methods.








10








Table 1 (Part A)

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
1	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide		27
2	2,6-dichloro- <i>N</i> -(3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)propyl)benzamide		27
3	2,6-dichloro- <i>N</i> -(3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl)benzamide		27
4	2,6-dichloro- <i>N</i> -(3-({4-(2,3-dihydro-1-benzofuran-6-yl)pyrimidin-2-yl}amino)propyl)benzamide		27








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
5	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-fluoro-6-iodobenzamide		27
6	<i>N</i> -(3-({4-[4-aminophenyl]pyrimidin-2-yl}amino)propyl)-2,6-dichlorobenzamide		27
7	<i>N</i> -[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		2
8	<i>N</i> -{4-[2-({3-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		29
9	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(dimethylamino)ethyl]benzamide		29
10	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluorobenzamide		39
11	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluoro-6-iodobenzamide		39

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
12	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide		39
13	<i>N</i> -(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
14	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyridine-4-carboxamide		39
15	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,3,4,5,6-pentafluorobenzamide		39
16	4-(4-chlorophenyl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
17	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		39
18	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
19	4-(2,4-dichlorophenyl)-N-{3-[(2-piperidin-1-ylethyl)oxy]phenyl}pyrimidin-2-amine		33
20	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chlorobenzamide		39
21	N-(4-{2-[(3-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3
22	N-(4-{2-[(3-piperidin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3
23	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-bromobenzamide		39
24	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3-fluorobenzamide		39
25	N-[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		39

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
26	<i>N</i> -(4-{2-[(3-{[(2,6-dichlorophenyl)sulfonyl]amino}phenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide		36
27	2,6-dichloro- <i>N</i> -(3-{[4-(1 <i>H</i> -indol-5-yl)pyrimidin-2-yl]amino}phenyl)benzamide		39
28	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]-5-fluoropyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		36
29	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-methylbenzamide		39
30	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,4-dichlorobenzamide		39
31	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,3-dichlorobenzamide		39
32	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,5-dichlorobenzamide		39








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
33	<i>N</i> -[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		3
34	<i>N</i> -(4-{2-[(4-piperidin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide		34
35	<i>N</i> -(4-{2-[(2-methyl-4-piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide		34
36	<i>N</i> -(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-carboxamide		1
37	<i>N</i> -(4-{5-methyl-2-[(3-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide		3
38	<i>N</i> -(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)-2-(phenyloxy)acetamide		1
39	<i>N</i> -(4-{6-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide		5








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
40	<i>N</i> -(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)-2-morpholin-4-ylacetamide		1
41	<i>N</i> -[4-(2-{3-(methoxy)phenyl}amino)pyrimidin-4-yl]phenylacetamide		
42	<i>N</i> -[3-({4-[4-(acetylamino)-2-chlorophenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		39
43	2,6-dichloro- <i>N</i> -{3-[(4-phenylpyrimidin-2-yl)amino]phenyl}benzamide		39
44	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-difluorobenzamide		39
45	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,4,5-trifluorobenzamide		39
46	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]benzamide		39








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
47	<i>N</i> -(4-{6-morpholin-4-yl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		37
48	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3,5-difluorobenzamide		39
49	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-fluoro-3-(methyloxy)benzamide		39
50	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-fluoro-4-methylbenzamide		39
51	<i>N</i> -(4-{2-[(3-{[(2,6-dimethylphenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
52	4-(2,4-dichlorophenyl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
53	4-(2,4-dichlorophenyl)- <i>N</i> -{3-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}pyrimidin-2-amine		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
54	<i>N</i> -(3-([4-(4-aminophenyl)pyrimidin-2-yl]amino)phenyl)-2,6-dichlorobenzamide		39
55	4-(4-aminophenyl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		8
56	4-[4-(ethylamino)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		49
57	<i>N</i> -[4-(2-{[3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
58	<i>N</i> -(4-{2-[(4-aminophenyl)amino]pyrimidin-4-yl}phenyl)acetamide		1
59	<i>N</i> -[4-(2-{[4-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
60	<i>N</i> -[4-(2-{[4-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
61	<i>N</i> -{4-[2-({3-[4-(pyridin-4-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
62	<i>N</i> -[5-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-2-morpholin-4-ylphenyl]-2,6-dichlorobenzamide		30
63	<i>N</i> -(4-{5-fluoro-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		5
64	<i>N</i> -(4-{2-[(4-{[2-(4-ethylpiperazin-1-yl)-2-oxoethyl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		33
65	<i>N</i> -[4-(2-{[3-(morpholin-4-ylcarbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		29
66	<i>N</i> -{3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-5-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}-2,6-dichlorobenzamide		31
67	4-[4-(dimethylamino)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
68	2,6-dichloro- <i>N</i> -(3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}phenyl)benzamide		39
69	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}phenyl)acetamide		5
70	<i>N</i> -(3-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-carboxamide		1
71	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-1-methylpiperidine-4-carboxamide		39
72	<i>N</i> -{4-[2-({3-[(phenylmethyl)amino]phenyl}amino)pyrimidin-4-yl}phenyl}acetamide		28
73	<i>N</i> -(4-{2-[(3-aminophenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide		1
74	<i>N</i> -(4-{2-[(3-aminophenyl)amino]-5-fluoropyrimidin-4-yl}phenyl)acetamide		5








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
75	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-ethylphenyl)benzamide		29
76	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(phenylmethyl)benzamide		29
77	N-{4-[2-({3-[(4-cyclopentylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		29
78	N-{4-[2-({3-[(4-phenylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		29
79	N-(3-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		9
80	N-(2-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		9
81	N-{4-[2-({3-[(4-pyrazin-2-yl)piperazin-1-yl]carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		29

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
82	<i>N</i> -(4-{2-[(3-{[4-(3-chlorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		46
83	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(1-methyl-1 <i>H</i> -benzimidazol-2-yl)methyl]benzamide		46
84	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -propylbenzamide		29
85	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -cyclopropylbenzamide		29
86	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(3-fluorophenyl)methyl]benzamide		29
87	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(naphthalen-1-ylmethyl)benzamide		29
88	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(dimethylamino)ethyl]- <i>N</i> -methylbenzamide		29

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
89	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-methylphenyl)methyl]benzamide		29
90	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(3-chlorophenyl)methyl]benzamide		29
91	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-phenylethyl)benzamide		29
92	N-{4-[2-({3-[(4-methylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		29
93	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(tetrahydrofuran-2-ylmethyl)benzamide		29
94	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[3-(2-oxopyrrolidin-1-yl)propyl]benzamide		29
95	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(3s,5s,7s)-tricyclo[3.3.1.1~3,7~]dec-1-yl]benzamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
96	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(methyloxy)ethyl]benzamide		29
97	N-[4-(2-({3-(1,3-thiazolidin-3-yl)carbonyl}phenyl)amino)pyrimidin-4-yl]phenyl]acetamide		29
98	N-{4-[2-({3-[(4-pyridin-2-yl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl]acetamide		29
99	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{2-(methyloxy)phenyl}methyl}benzamide		29
100	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{3-(methyloxy)phenyl}methyl}benzamide		29
101	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-fluorophenyl)methyl]benzamide		29
102	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(4-fluorophenyl)methyl]benzamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
103	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(3,3-dimethylbutyl)benzamide		29
104	N-[4-(2-{{3-(thiomorpholin-4-yl)carbonyl}phenyl}amino)pyrimidin-4-yl]phenylacetamide		29
105	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-thienylmethyl)benzamide		29
106	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[3-(dimethylamino)propyl]benzamide		29
107	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{2-(trifluoromethyl)phenyl}methyl}benzamide		29
108	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{3-(trifluoromethyl)phenyl}methyl}benzamide		29
109	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{4-(trifluoromethyl)phenyl}methyl}benzamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
110	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -[(2,4-difluorophenyl)methyl]benzamide		29
111	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -ethyl- <i>N</i> -methylbenzamide		29
112	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -({4-[(trifluoromethyl)oxy]phenyl}methyl)benzamide		29
113	<i>N</i> -{4-[2-({3-[(4-acetyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		29
114	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -(cyclopropylmethyl)benzamide		29
115	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -[2-(2-fluorophenyl)ethyl]benzamide		29
116	<i>N</i> -[4-(2-({3-(pyrrolidin-1-yl)carbonyl}phenyl)amino)pyrimidin-4-yl]phenyl}acetamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
117	<i>N</i> -{4-[2-({3-[(4-pyrimidin-2-yl)piperazin-1-yl]carbonyl}phenyl)amino)pyrimidin-4-yl}phenyl}acetamide		29
118	<i>N</i> -methyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		32
119	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)benzamide		9
120	<i>N</i> -[4-(2-{[3-(1,3-dioxan-2-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
121	<i>N</i> -[4-(2-{[3-(morpholin-4-yl)methyl]phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		22
122	<i>N</i> -{4-[2-({3-[(4-ethylpiperazin-1-yl)methyl]phenyl}amino)pyrimidin-4-yl}phenyl}acetamide		29
123	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3-[(2-morpholin-4-ylethyl)oxy]benzamide		39








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
124	4-[4-(methylamino)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		29
125	N-[4-(2-{{4-(4-acetylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		7
126	N-(4-{2-[(3-amino-2,4,5,6-tetrafluorophenyl)amino]pyrimidin-4-yl}phenyl)acetamide		1
127	N-(4-{2-[(3-{[4-(2-fluorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		
128	3-{{4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino}-N-[2-(phenyloxy)ethyl]benzamide		29
129	methyl 1-{{3-{{4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino}phenyl}carbonyl}piperidine-4-carboxylate		29
130	N-[4-(2-{{3-{{4-[3-(methyloxy)phenyl]piperazin-1-yl}carbonyl}phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
131	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -{2-[2-(methyloxy)phenyl]ethyl} benzamide		29
132	<i>N</i> -[4-(2-{[3-(1,3-dihydro-2 <i>H</i> -isoindol-2-ylcarbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		29
133	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -(biphenyl-4-ylmethyl)benzamide		29
134	<i>N</i> -(4-{2-[(3-{[4-(phenylcarbonyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		29
135	<i>N</i> -[4-(2-{[3-({4-[4-(methyloxy)phenyl]piperazin-1-yl}carbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		29
136	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -methyl- <i>N</i> -{[2-(methyloxy)phenyl]methyl} benzamide		29
137	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -[(2-fluorophenyl)methyl]- <i>N</i> -methylbenzamide		29

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
138	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(diphenylmethyl)benzamide		29
139	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(2-pyridin-2-ylethyl)benzamide		29
140	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(pyridin-2-ylmethyl)benzamide		29
141	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(2-chlorophenyl)ethyl]benzamide		29
142	<i>N</i> -[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}-5-fluoropyrimidin-4-yl)phenyl]acetamide		5
143	<i>N</i> ² -[3-(1 <i>H</i> -imidazol-1-yl)propyl]- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide		23
144	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)- <i>N</i> ² -(2-pyridin-3-ylethyl)glycinamide		23

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
145	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(pyridin-3-ylmethyl)benzamide		29
146	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(pyridin-4-ylmethyl)benzamide		29
147	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-methyl-N-(phenylmethyl)benzamide		29
148	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-cyclopentylbenzamide		29
149	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-chlorophenyl)methyl]benzamide		29
150	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(4-chlorophenyl)methyl]benzamide		29
151	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(furan-2-ylmethyl)benzamide		29

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
152	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)-N-{{4-(methyloxy)phenyl}methyl} benzamide		29
153	N-[4-(2-{{3-({4-[2-(methyloxy)phenyl]piperazin-1-yl} carbonyl)phenyl} amino} pyrimidin-4-yl)phenyl]acetamide		29
154	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)-N-[3-(methyloxy)propyl]benzamide		29
155	N-(4-{2-{{3-({[(2R,6S)-2,6-dimethylmorpholin-4-yl] carbonyl)phenyl} amino} pyrimidin-4-yl}phenyl)acetamide		29
156	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)-N-[(6-chloropyridin-3-yl)methyl]benzamide		29
157	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)-N-butylbenzamide		29
158	N-(4-{2-{{3-{{4-(2-chlorophenyl)piperazin-1-yl} carbonyl)phenyl} amino} pyrimidin-4-yl}phenyl)acetamide		29








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
159	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -ethyl- <i>N</i> -(2-(methyloxy)ethyl)benzamide		29
160	<i>N</i> -(4-(2-(3-(3-morpholinopropoxy)phenylamino)pyrimidin-4-yl)phenyl)acetamide		35
161	<i>N</i> -(4-(2-(3-(2-(dimethylamino)ethoxy)phenylamino)pyrimidin-4-yl)phenyl)acetamide		33
162	<i>N</i> -[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide		39
163	<i>N</i> -[4-(2-{[4-(phenyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
164	4-(4-aminophenyl)- <i>N</i> -[4-(phenyloxy)phenyl]pyrimidin-2-amine		8
165	<i>N</i> -[4-[2-({4-[(phenylmethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl]acetamide		3








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
166	4-(4-aminophenyl)-N-[3-(morpholin-4-ylsulfonyl)phenyl]pyrimidin-2-amine		8
167	N-(4-{2-[(3,5-dimorpholin-4-yl)phenyl]amino}-5-fluoropyrimidin-4-yl)phenyl)acetamide		6
168	N-(4-{2-[(4-{4-(phenylmethyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
169	N-(4-{2-[(4-{4-[(5-methyl-3-phenylisoxazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
170	N-(4-{2-[(4-{4-[(5-methyl-1-phenyl-1H-pyrazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
171	N-(4-{2-[(4-{4-[(2-phenyl-1,3-thiazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
172	N-[4-(2-{[4-(4-{[6-(phenyloxy)pyridin-3-yl]methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl)phenyl)acetamide		49








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
173	<i>N</i> -{4-[2-({4-[4-(cyclohexylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
174	<i>N</i> -(4-{2-[(4-{4-[(1 <i>S</i> ,4 <i>S</i>)-bicyclo[2.2.1]hept-5-en-2-ylmethyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
175	<i>N</i> -[4-(2-{[4-(4-pentylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		49
176	<i>N</i> -(4-{2-[(4-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
177	<i>N</i> -[4-(2-{[4-(4-{[3,5-bis(methoxy)phenyl]methyl]piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49
178	<i>N</i> -(4-{2-[(4-{4-[(4-fluorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
179	<i>N</i> -(4-{2-[(4-{4-[(1-methyl-1 <i>H</i> -pyrrol-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
180	<i>N</i> -(4-{2-[(4-{4-(2,4-dichlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
181	<i>N</i> -(4-{2-[(4-{4-(9 <i>H</i> -fluoren-2-ylmethyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
182	<i>N</i> -(4-{2-[(4-{4-[(3-methyl-2-thienyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
183	<i>N</i> -(4-{2-[(4-{4-[(5-ethylfuran-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
184	<i>N</i> -(4-{2-[(4-{4-[(3-{4-(1,1-dimethylethyl)phenoxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
185	<i>N</i> -(4-{2-[(4-{4-(3-thienylmethyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
186	methyl 4-({4-[4-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazin-1-yl}methyl)benzoate		49








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
187	<i>N</i> -(4-{2-[(4-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
188	<i>N</i> -(4-{2-[(4-{4-[(4-{[3-(dimethylamino)propyl]oxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
189	<i>N</i> -(4-(2-{[4-(4-{2-[(phenylmethyl)oxy]ethyl]piperazin-1-yl}phenyl)amino}pyrimidin-4-yl}phenyl)acetamide		49
190	<i>N</i> -(4-{2-[(4-{4-[(2-chloroquinolin-3-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
191	<i>N</i> -(4-{2-[(4-{4-[(4-chloro-2,6-dimethylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
192	<i>N</i> -(1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyrrolidin-3-yl)acetamide		3
193	<i>N</i> ² -[3-(4-methylpiperazin-1-yl)propyl]- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide		23








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
194	<i>N</i> ² -(1-methylpiperidin-4-yl)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide		23
195	<i>N</i> -(4-{2-[(4-{[(pyridin-4-yl)methyl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		33
196	<i>N</i> -(4-{2-[(4-{[2-(methyloxy)ethyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
197	2-(dimethylamino)- <i>N</i> -(4-{2-(4-morpholinophenylamino)pyrimidin-4-yl}phenyl)acetamide		12
198	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)furan-2-carboxamide		12
199	2-(methyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		12
200	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)cyclobutanecarboxamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
201	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-3-carboxamide		13
202	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-2-carboxamide		13
203	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-3-carboxamide		13
204	<i>N</i> -[4-(2-{[4-(dimethylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
205	<i>N</i> -(4-{2-[(4-chlorophenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3
206	<i>N</i> -(4-{2-[(3-{[(2-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
207	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-4-carboxamide		13

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
208	2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14
209	<i>N</i> -(4-(2-[(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl)phenyl)glycinamide		14
210	<i>N</i> -(4-(2-[(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl)phenyl)morpholine-2-carboxamide		13
211	<i>N</i> ² -methyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide		13
212	<i>N</i> -(4-(2-[(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl)phenyl)-beta-alaninamide		13
213	<i>N</i> -(4-(2-[(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl)phenyl)phenylalaninamide		13
214	<i>N</i> -[4-(2-{[4-(3-oxopiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		34








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
215	<i>N</i> -[4-(2-{{4-(4-{{5-(3-chlorophenyl)furan-2-yl)methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49
216	<i>N</i> -[4-(2-{{4-(4-{{4-fluoro-2-(trifluoromethyl)phenyl}methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49
217	<i>N</i> -[4-(2-{{4-(4-{{4-(1 <i>H</i> -imidazol-1-yl)phenyl)methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49
218	<i>N</i> -[4-(2-{{4-(4-{{[2,5-bis(trifluoromethyl)phenyl]methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49
219	<i>N</i> -(4-{2-[[4-{4-[(2,6-dimethylphenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		49
220	<i>N</i> -(4-{2-[[4-{4-[(2,3-dimethylphenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		49
221	<i>N</i> -[4-(2-{{4-(4-{{[2,4-bis(ethoxy)phenyl]methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		49

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
222	<i>N</i> -[4-(2-{{4-(4-{{3-(ethyloxy)phenyl}methyl}piperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		49
223	<i>N</i> -{4-[2-({4-[4-(2,2'-bithien-5-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
224	<i>N</i> -[4-(2-{{4-(4-{{4-(2-thienyl)phenyl}methyl}piperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		49
225	<i>N</i> -(4-{2-[(4-{4-[(4-cyanophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
226	<i>N</i> -[4-(2-{{4-(4-{{2,5-bis(methyloxy)phenyl}methyl}piperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		49
227	<i>N</i> -{4-[2-({4-[4-(2,2-diphenylethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
228	<i>N</i> -{4-[2-({4-[4-(1 <i>H</i> -pyrrol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
229	<i>N</i> -{4-[2-(1 <i>H</i> -indazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide		3
230	<i>N</i> -{4-[2-(1 <i>H</i> -indol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide		3
231	<i>N</i> -[4-(2-{[4-(morpholin-4-ylmethyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		22
232	<i>N</i> -(4-{2-[(3-{[(3-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
233	<i>N</i> -(4-{2-[(3-{[(4-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
234	4-[4-(4-{4-(acetylamino)phenyl}pyrimidin-2-yl)amino)phenyl]- <i>N</i> -ethylpiperazine-1-carboxamide		7
235	<i>N</i> -(4-[2-({4-[4-(ethylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl)acetamide		36








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
236	<i>N</i> -{4-[2-(1 <i>H</i> -indazol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide		3
237	<i>N</i> -[4-(2-{[4-(4-propylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		49
238	<i>N</i> -[4-(2-{[4-(4-butylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		49
239	<i>N</i> -{4-[2-({4-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
240	4-[4-(methylsulfonyl)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		16
241	ethyl <i>N</i> -[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]- <i>N</i> -methylglycinate		3
242	4-[3-(methylsulfonyl)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
243	4-[4-(methylthio)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
244	N-(4-{2-[(4-cyclohexylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3
245	N-{4-[2-({4-[(tetrahydrofuran-2-yl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		28
246	N-{4-[2-({4-[(phenylmethyl)amino]phenyl}amino)pyrimidin-4-yl}phenyl}acetamide		28
247	N-[4-(2-{[4-(acetylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
248	methyl (4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)carbamate		17
249	1-ethyl-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea		17








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
250	ethyl 1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-3-carboxylate		3
251	ethyl [4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]acetate		3
252	4-[4-(methyloxy)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
253	4-[3-(methyloxy)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
254	4-(1H-indol-5-yl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
255	2-[(2-amino-2-oxoethyl)amino]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		23
256	2-morpholin-4-yl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
257	2,6-dichloro- <i>N</i> -{3-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}benzamide		2
258	<i>N</i> ² -(2-aminoethyl)- <i>N</i> -2-methyl- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide		1
259	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)- <i>N</i> ² -1 <i>H</i> -pyrazol-5-ylglycinamide		23
260	phenylmethyl <i>N</i> -(2-{[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]-2-oxoethyl}- <i>L</i> -alaninate		23
261	4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzamide		9
262	1,1-dimethylethyl [(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)methyl]carbamate		9
263	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
264	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylacetamide		12
265	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylpropanamide		12
266	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-2-carboxamide		12
267	5-methyl- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrazine-2-carboxamide		12
268	2-(ethyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		23
269	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-(phenyloxy)acetamide		23
270	<i>N</i> -[4-(2-{[4-(1 <i>H</i> -pyrrol-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
271	<i>N</i> -[4-(2-{[4-(2,6-dimethylmorpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
272	ethyl 1-[4-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-4-carboxylate		3
273	2-cyclopentyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		12
274	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}-3-pyridin-3-yl)propanamide		12
275	6-(methyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)pyridine-3-carboxamide		12
276	methyl 4-[(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]amino]-4-oxobutanoate		12
277	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)butanamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
278	<i>N</i> -(4-{2-[(4-{bis[2-(methoxy)ethyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3
279	<i>N</i> -[4-(2-{[4-(morpholin-4-ylsulfonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
280	4-(4-(aminomethyl)phenyl)- <i>N</i> -(4-morpholinophenyl)pyrimidin-2-amine		47
281	<i>N</i> -[4-(2-{[4-(morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)methyl]acetamide		48
282	<i>N</i> -(4-morpholin-4-ylphenyl)-4-{4-[(propylamino)methyl]phenyl}pyrimidin-2-amine		47
283	<i>N</i> -(4-{2-[(4-piperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		3
284	<i>N</i> -(4-{2-[(3,5-dimorpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		6








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
285	2-(2-methylphenyl)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
286	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopentanecarboxamide		12
287	<i>N,N</i> -dimethyl- <i>N'</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanediamide		12
288	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)- <i>N</i> -(2-pyrimidin-4-yl)glycinamide		12
289	3-chloro- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide		12
290	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-piperidin-1-ylacetamide		12
291	<i>N</i> ² -ethyl- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
292	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyrrolidin-1-ylacetamide		11
293	2-(1 <i>H</i> -imidazol-1-yl)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		11
294	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-piperazin-1-ylacetamide		11
295	<i>N</i> -[4-(2-{[4-(4-phenylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
296	<i>N</i> -(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		15
297	<i>N</i> -(4-{2-[(4-piperazin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		34
298	2,6-dichloro- <i>N</i> -(3-{[4-(4-fluorophenyl)pyrimidin-2-yl]amino}propyl)benzamide		27








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
299	<i>N</i> -[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyridin-2-yl]-2,6-dichlorobenzamide		38
300	<i>N</i> -[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyrimidin-4-yl]-2,6-dichlorobenzamide		38
301	<i>N</i> -(4-{2-[(6-aminopyridin-2-yl)amino]pyrimidin-4-yl}phenyl)acetamide		38
302	<i>N</i> -(4-{2-[(6-aminopyrimidin-4-yl)amino]pyrimidin-4-yl}phenyl)acetamide		38
303	5-fluoro- <i>N</i> ⁴ -[2-(methyloxy)phenyl]- <i>N</i> ² -[3-(methyloxy)phenyl]pyrimidine-2,4-diamine		50
304	2,6-dichloro- <i>N</i> -{3-[(4-{[3-chloro-4-(methyloxy)phenyl]oxy}pyrimidin-2-yl)amino]phenyl}benzamide		41
305	4-{[2-chloro-4-(methyloxy)phenyl]oxy}- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		41








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
306	<i>N</i> -[4-({2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}amino)phenyl]acetamide		35
307	<i>N</i> -[4-({2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}oxy)phenyl]acetamide		41
308	<i>N</i> -{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
309	<i>N</i> -{4-[2-({4-[(2 <i>R</i> ,6 <i>S</i>)-2,6-dimethylmorpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		3
310	<i>N</i> -(4-{2-[(4-{4-[(2-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
311	<i>N</i> -{4-[2-({4-[4-(1 <i>H</i> -pyrazol-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
312	<i>N</i> -(4-{2-[(3-piperazin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		34








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
313	<i>N</i> -[4-(2-{[3-(4-{[2-(methoxy)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		49
314	<i>N</i> -(4-[2-(3-[4-(1,3-thiazol-2-ylmethyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl)phenyl]acetamide		49
315	<i>N</i> -(4-[2-(3-bromo-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl)phenyl]acetamide		3
316	<i>N</i> -(4-[2-([4-{[2-(diethylamino)ethyl]oxy}-3-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
317	4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzoic acid		9
318	4-(4-furan-2-ylphenyl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
319	4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine		10








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
320	4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl} benzonitrile		9
321	methyl 4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl} benzoate		9
322	4-(4-fluorophenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
323	N-[3-(2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl)amino]phenyl]acetamide		35
324	N-(4-morpholin-4-ylphenyl)-4-[4-(pyridin-3-ylamino)phenyl]pyrimidin-2-amine		9
325	N-(4-morpholin-4-ylphenyl)-4-[4-(pyridin-2-ylamino)phenyl]pyrimidin-2-amine		9
326	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl} phenyl)methanesulfonamide		10







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
327	1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-(phenylmethyl)urea		17
328	4-(2,3-dihydro-1,4-benzodioxin-6-yl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
329	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}phenyl)acetamide		43
330	N-(4-morpholin-4-ylphenyl)-4-quinolin-6-ylpyrimidin-2-amine		9
331	4-[4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9
332	N-(4-morpholin-4-ylphenyl)-4-(4-pyrimidin-5-ylphenyl)pyrimidin-2-amine		9
333	N-(4-morpholin-4-ylphenyl)-4-quinoxalin-6-ylpyrimidin-2-amine		9







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
334	2-chloro- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12
335	2-(2-fluorophenyl)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
336	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrimidine-5-carboxamide		12
337	(2 <i>S</i>)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-2-carboxamide		45
338	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)- <i>N</i> -(2-phenylglycinamide)		23
339	<i>N</i> -(4-{2-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-ylphenyl}acetamide		29
340	<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide		14








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
341	<i>N</i> -(4-(2-(3-methoxy-4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)acetamide		15
342	<i>N</i> -(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)-pyrimidin-4-yl)phenyl)acetamide		7
343	<i>N</i> -(4-(2-(4-(4-(3-methylbutanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
344	<i>N</i> -(4-(2-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
345	<i>N</i> -(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
346	<i>N</i> -(4-(2-(4-(4-(cyclopentanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
347	<i>N</i> -(4-(2-(4-(4-(2-methoxybenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
348	<i>N</i> -(4-(2-(4-(4-pentanoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
349	<i>N</i> -(4-(2-(4-(4-picolinoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
350	<i>N</i> -(4-(2-(4-(4-isonicotinoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
351	<i>N</i> -(4-(2-(4-(4-(1-acetyl)piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
352	<i>N</i> -(4-(2-(4-(4-(2-cyclopropylacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
353	<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
354	<i>N</i> -(4-(2-(4-(4-(2-(2-methoxyethoxy)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
355	<i>N</i> -(4-(2-(4-(4-(2-(pyridin-3-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
356	<i>N</i> -(4-(2-(4-(4-(3-(pyridin-3-yl)propanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
357	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		12
358	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyridin-3-yl)acetamide		12
359	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)isonicotinamide		12
360	<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-prolinamide		45
361	<i>N</i> -[4-(2-{[3-(methoxy)-4-morpholin-4-yl]phenyl}amino)pyrimidin-4-yl)phenyl]-D-prolinamide		45

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
362	O-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-serinamide		14
363	(S)-3-hydroxy-N-(4-(2-(4-morpholino-phenylamino)-pyrimidin-4-yl)-phenyl)-butanamide		18
364	(R)-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		20
365	(R)-2-amino-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		21
366	2-Hydroxy-2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		11
367	2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		44







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
368	<i>(R)</i> - <i>N</i> -(4-(2-(4-((<i>R</i>)-3-(dimethylamino)pyrrolidin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
369	4-amino-1,1-dioxo- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydro-2 <i>H</i> -thiopyran-4-carboxamide		44
370	<i>(R)</i> -4-(4-aminophenyl)- <i>N</i> -(4-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)pyrimidin-2-amine		8
371	<i>(R)</i> - <i>N</i> -(4-(2-(4-(3-(dimethylamino)pyrrolidin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide		12
372	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperazine-2-carboxamide		12
373	2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1,2,3,4-tetrahydronaphthalene-2-carboxamide		44

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
374	4-(4-(1,1-dioxo-isothiazolidin-2-yl)phenyl)- <i>N</i> -(4-morpholinophenyl)-pyrimidin-2-amine		25
375	4-(4-(1 <i>H</i> -tetrazol-1-yl)phenyl)- <i>N</i> -(4-morpholinophenyl)-pyrimidin-2-amine		25
376	<i>(R)</i> - <i>N</i> -(4-(2-(3-(benzyloxy)-4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide		45
377	<i>(S)</i> -2-amino-3-hydroxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		21
378	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(1 <i>H</i> -tetrazol-1-yl)acetamide		24
379	<i>(R)</i> - <i>N</i> -(4-(2-(3-ethoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide		45
380	<i>(R)</i> - <i>N</i> -(4-(2-(1,2,3,4-tetrahydroquinolin-6-ylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide		45







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
381	3-hydroxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-3-methylbutanamide		18
382	(3 <i>S</i> ,7 <i>S</i>)-7-(hydroxymethyl)- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)quinuclidine-3-carboxamide		18
383	1-hydroxy- <i>N</i> -(4-(2-(4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		18
384	(S)-2-amino- <i>N</i> -(4-(2-(3-methyl-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14
385	(R)- <i>N</i> -(4-(2-(3-methyl-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
386	(R)- <i>N</i> -(4-(2-(4-morpholino-3-(trifluoromethyl)-phenylamino)pyrimidin-4-yl)-phenyl)-pyrrolidine-2-carboxamide		45








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
387	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-((<i>S</i>)-tetrahydrofuran-2-carbonyl)-piperazin-1-yl)-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide		45
388	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-((<i>R</i>)-tetrahydrofuran-2-carbonyl)-piperazin-1-yl)-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide		45
389	4-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperazine-1-carboxamide		12
390	3-methoxy- <i>N</i> -(4-(2-(4-morpholino-3-(trifluoromethyl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		12
391	3-methoxy- <i>N</i> -(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-propane-1-sulfonamide		16
392	2-methoxy- <i>N</i> -(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-ethanesulfonamide		16








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
393	(<i>S</i>)-3-hydroxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		18
394	(<i>R</i>)-3-hydroxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		20
395	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2,5-dihydro-1 <i>H</i> -pyrrole-2-carboxamide		13
396	1-(3-(dimethylamino)propyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea		17
397	(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-((<i>S</i>)-pyrrolidin-2-ylmethyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
398	(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)-5-methylpyrimidin-4-yl)phenyl)propanamide		14
399	1-(3-methoxypropyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea		17








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
400	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)-5-methylpyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
401	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)-3-fluorophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide		45
402	<i>(R)</i> - <i>N</i> -(4-(2-(3-chloro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
403	1-(2-morpholinoethyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea		17
404	1-(2-(dimethylamino)ethyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea		17
405	<i>(S)</i> - <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyrrolidin-2-yl)acetamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
406	2,3-dihydroxy- <i>N</i> -(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-propanamide		18
407	(<i>S</i>)-2-amino-4-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide		14
408	(<i>R</i>)-2-amino-4-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide		14
409	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)isindoline-1-carboxamide		13
410	<i>N</i> -ethyl-4-(4-(4-(4-(tetrahydrofuran-2-carboxamido)phenyl)pyrimidin-2-ylamino)phenyl)piperazine-1-carboxamide		12
411	<i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12
412	(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-((<i>R</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
413	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-((<i>S</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
414	<i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		12
415	3-methoxy- <i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		12
416	<i>N</i> -(4-(2-(4-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		12
417	<i>(R)</i> - <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1-methylpyrrolidine-2-carboxamide		12
418	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
419	<i>(R)</i> -4-(4-(4-(4-(2-aminopropanamido)phenyl)-pyrimidin-2-ylamino)phenyl)- <i>N</i> -ethylpiperazine-1-carboxamide		14
420	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-((<i>R</i>)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
421	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-((<i>S</i>)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
422	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-((<i>S</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
423	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
424	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
425	<i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
426	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
427	1-ethyl-3-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea		17
428	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
429	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide		12
430	<i>(S)</i> -3-methoxy- <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		12
431	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		1
432	<i>N</i> -(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
433	<i>N</i> -(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
434	<i>N</i> -(4-(2-(4-(1-butyryl-1,2,4-triazin-4-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		11
435	1-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea		17
436	<i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide		11
437	<i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
438	<i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		11
439	1-ethyl-3-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		17

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
440	1-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea		17
441	1-ethyl-3-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		17
442	<i>N</i> -ethyl-4-(4-(4-(4-(3-ethylureido)phenyl)pyrimidin-2-ylamino)phenyl)piperazine-1-carboxamide		17
443	(S)-1-ethyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
444	(R)-1-(2-hydroxyethyl)- <i>N</i> -(4-(2-(4-morpholinobenzyl)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		47
445	(R)-1-isopropyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		42
446	(S)-2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
447	1-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea		17
448	(R)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
449	4-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydro-2H-pyran-4-carboxamide		44
450	(R)-2-amino-N-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
451	(R)-2-amino-N-(4-(2-(4-(4-((R)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
452	(R)-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
453	(R)-2-amino-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
454	<i>(R)</i> -2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12
455	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
456	<i>(S)</i> - <i>N</i> -(4-(5-methyl-2-(4-(4-((1-methyl-1 <i>H</i> -imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
457	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
458	<i>(2R)</i> - <i>N</i> -(4-(2-(4-(4-(tetrahydrofuran-3-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
459	<i>(S)</i> -1-ethyl-3-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		17
460	<i>(S)</i> -1-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea		17

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
461	<i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		11
462	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		11
463	3-methoxy- <i>N</i> -(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		12
464	<i>(R)</i> -2-amino- <i>N</i> -(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14
465	2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12
466	1-ethyl-3-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		17
467	3-methoxy- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
468	<i>(R)</i> - <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide		45
469	<i>(S)</i> - <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide		45
470	<i>(S)</i> - <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide		42
471	<i>(2R,3S)</i> -2-amino-3-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		18
472	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14
473	<i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
474	<i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butyramide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
475	<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		12
476	<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		12
477	(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
478	(2 <i>S</i> ,3 <i>R</i>)-2-amino-3-hydroxy- <i>N</i> -(4-(2-(4-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		14
479	(<i>R</i>)- <i>N</i> -(4-(2-(4-((2 <i>S</i> ,6 <i>R</i>)-2,6-dimethylmorpholino)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
480	(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(3-hydroxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
481	1-ethyl-3-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
482	<i>(R)</i> -1-ethyl-3-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		12
483	3,3,3-trifluoro-2-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		18
484	<i>(R)</i> -1-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea		17
485	2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12
486	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		20
487	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
488	<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
489	<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide		12
490	<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		12
491	<i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		11
492	<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		11
493	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		11
494	2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12
495	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
496	3-methoxy- <i>N</i> -(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		12
497	1-ethyl-3-(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea		17
498	<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropanoyl)piperazin-1-yl)benzyl)pyrimidin-4-yl)phenyl)acetamide		49
499	(S)- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12
500	(R)- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12
501	(4S)-4-hydroxy- <i>N</i> -(4-(2-(4-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		21
502	(R)- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
503	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12
504	<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		11
505	<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		11
506	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide		11
507	<i>(2R)</i> - <i>N</i> -(4-(2-(4-(4-(tetrahydrofuran-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
508	<i>(R)</i> - <i>N</i> -(4-(5-chloro-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
509	<i>N</i> -(4-(2-(4-(4-(3-(diethylamino)propanoyl)piperazin-1-yl)benzyl)pyrimidin-4-yl)phenyl)acetamide		7

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
510	(S)-1-(2-hydroxyethyl)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide		42
511	(S)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanediamide		14
512	(R)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanediamide		18
513	(R)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)succinamide		20
514	(R)-N-(4-(2-(4-(4-chloro-1-methyl-1H-pyrazol-3-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
515	(S)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide		42
516	(R)-N-(4-(2-(4-(4-ethoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
517	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-(pyrrolidin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
518	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(2-(morpholinoacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
519	<i>N</i> -(4-(2-(4-(morpholinophenylamino)pyrimidin-4-yl)phenyl)-1 <i>H</i> -imidazole-4-carboxamide		12
520	2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		12
521	<i>(S)</i> - <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide		12
522	<i>(R)</i> -2-hydroxy-2-methyl- <i>N</i> -(4-(2-(4-(morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		18
523	<i>(S)</i> -2-hydroxy-2-methyl- <i>N</i> -(4-(2-(4-(morpholinobenzyl)pyrimidin-4-yl)phenyl)butanamide		18








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
524	<i>(R)</i> -2-methoxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		11
525	<i>(S)</i> -2-methoxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		11
526	<i>(R)</i> - <i>N</i> -(4-(2-(4-(2-methoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		49
527	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-acetylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		49
528	<i>(S)</i> -2-amino- <i>N</i> -(4-(5-methyl-2-(4-(4-((1-methyl-1 <i>H</i> -imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide		14
529	<i>N</i> -(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide		12
530	<i>(2R,4S)</i> -4-hydroxy- <i>N</i> -(4-(2-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		21








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
531	1-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopentanecarboxamide		44
532	<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-formylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
533	<i>(R)</i> -1-(2-hydroxyethyl)- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide		42
534	1-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		12
535	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1 <i>H</i> -pyrrole-2-carboxamide		12
536	<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1 <i>H</i> -imidazole-2-carboxamide		12
537	<i>(S)</i> -2-hydroxy-3,3-dimethyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		20








Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
538	(R)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide		18
539	(S)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide		20
540	(S)-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		20
541	1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclobutanecarboxamide		14
542	(R)-N-(4-(2-(6-morpholinopyridin-3-ylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
543	(S)-N-(4-(2-(3-chloro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
544	(2R,3R)-2-amino-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide		14



Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
545	1-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopentanecarboxamide		18
546	(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(dimethylamino)butanoyl)pyrazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
547	(<i>R</i>)- <i>N</i> -(4-(2-(4-(2-methoxyethyl)-3,4-dihydro-2 <i>H</i> -benzo[<i>b</i>][1,4]oxazin-7-ylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
548	(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide		14
549	(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide		14
550	(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)hexanamide		14
551	(<i>R</i>)-2-amino-3-methoxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
552	(2 <i>S</i> ,3 <i>R</i>)-2-amino-3-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide		14
553	(R)- <i>N</i> -(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
554	<i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(1 <i>H</i> -tetrazol-1-yl)acetamide		24
555	(S)- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)indoline-2-carboxamide		45
556	(R)-tert-butyl 2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylcarbamoyl)pyrrolidine-1-carboxylate		45
557	1-acetyl-4-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperidine-4-carboxamide		44
558	(R)-2-amino-3-methoxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
559	(S)-N-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
560	(R)-2-amino-N-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide		14
561	2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide		18
562	(R)-N-(4-(2-(4-(4-(2-hydroxyethyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
563	(R)-N-(4-(2-(4-(4-((1-methyl-1H-pyrrol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
564	(R)-N-(4-(2-(4-(4-((R)-pyrrolidin-2-ylmethyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide		45
565	(2S,3aS,7aS)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)octahydro-1H-indole-2-carboxamide		45



Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
566	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopropanecarboxamide		12
567	N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopropanecarboxamide		12
568	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)valinamide		14
569	N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
570	N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide		6
571	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide		14
572	N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
573	2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylacetamide		4
574	N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		5
575	3-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		12
576	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide		13
577	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide		14
578	N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
579	N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
580	N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)-1,3-thiazole-4-carboxamide		12
581	2,6-dichloro-N-(3-{[4-(2,3-dihydro-1,4-benzodioxin-6-yl)pyrimidin-2-yl]amino}propyl)benzamide		27





5





Table 1 (Part B)





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
582	N-(4-{2-[(2-methyl-4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		3
583	N-(4-{2-[(4-pyrrolidin-1-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
584	N-[4-(2-([4-(diethylamino)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide		3
585	N-(4-{2-[(4-azepan-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
586	N-{4-[2-({4-[methyl(2-phenylethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		3
587	N-[4-(2-{[4-(1,4-dioxaspiro[4.5]dec-8-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
588	N-[4-(2-{[4-(2-oxopiperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
589	N-[4-(2-{[4-(2-methylpiperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		3
590	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-valinamide	<p>Chiral</p> 	45
591	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-valinamide	<p>Chiral</p> 	45
592	2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)alaninamide		14





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
593	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tryptophanamide		14
594	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1,2,3,4-tetrahydroisoquinoline-1-carboxamide		14
595	O-(1,1-dimethylethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide	<p>Chiral</p> 	45
596	3-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-carboxamide		14





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
597	bis(1,1-dimethylethyl) (2R)-2-[[[4-{2-[(4-morpholin-4-yl)phenyl]amino]pyrimidin-4-yl}phenyl]amino]carbon-yl]piperazine-1,4-dicarboxylate	 <p>Chiral</p>	14
598	N-(4-{2-[(4-{4-[2-(2-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
599	N-(4-{2-[(4-{4-[2-(2-methylphenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
600	N-(4-{2-[(4-{4-[2-(3-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
601	N-{4-[2-({4-[4-(3-thienylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
602	N-(4-{2-[(4-{4-[(6-chloropyridin-3-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
603	N-(4-{2-[(4-{4-[(3-methylfuran-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
604	N-(4-(2-(4-(4-(3-fluoro-2-methylbenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
605	N-(4-(2-(4-(4-(1H-imidazole-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
606	N-(4-(2-(4-(4-(2-methoxynicotinoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7
607	N-(4-(2-(4-(4-(4-fluoro-3-methylbenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
608	N-{4-[2-({4-[4-(naphthalen-2-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36
609	N-{4-[2-({4-[4-(quinolin-8-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36
610	N-{4-[2-({4-[4-({4-(1,1-dimethylethyl)phenyl)sulfonyl]piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
611	N-[4-(2-[[4-(4-{5-bromo-2-(methyloxy)phenyl}sulfonyl)piperazin-1-yl]phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		36
612	N-(4-{2-[(4-{4-[(phenylmethyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
613	N-[4-(2-[[4-(4-{3-(trifluoromethyl)phenyl}sulfonyl)piperazin-1-yl]phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		36
614	N-(4-{2-[(4-{4-[(2-methylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
615	N-(4-{2-[(4-{(3-fluorophenyl)sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		36
616	N-(4-{2-[(4-{(2,4-difluorophenyl)sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		36
617	N-{4-[2-[(3-{4-[(4-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		49
618	N-(4-{2-[(3-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		49





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
619	N-{4-[2-({3-[4-({2-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		49
620	N-(4-{2-[(3-{4-[(3-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
621	N-{4-[2-({3-[4-(2,3-dihydroxypropyl)]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		49
622	N-{4-[2-({3-[4-(1,3-benzodioxol-5-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		49
623	N-{4-[2-({3-[4-(pyridin-2-ylmethyl)]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		49

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
624	N-{4-[2-({3-[4-(pyridin-3-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
625	N-{4-[2-({3-[4-(1H-pyrrol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
626	4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-(phenylmethyl)piperazine-1-carboxamide		53
627	N-[4-(2-{[3-(4-{[2-(methyloxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		7




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
628	N-{4-[2-({3-[4-(1H-pyrazol-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
629	N-{4-[2-({3-[4-(3-pyridin-3-ylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
630	N-{4-[2-({3-[4-(3-(methyloxy)propanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
631	N-[4-(2-({3-(4-(2-((4-fluorophenyl)oxy)acetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl]acetamide		7




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
632	N-{4-[2-({3-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
633	N-{4-[2-({3-[4-(pyridin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
634	N-{4-[2-({3-[4-(pyridin-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
635	N-{4-[2-({3-[4-(2-methylphenylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
636	N-{4-[2-({3-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
637	N-{4-[2-({3-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
638	N-{4-[2-({3-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
639	N-[4-(2-({3-(methyloxy)-4-morpholin-4-ylphenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		12
640	(2R)-N-[4-(2-({3-(methyloxy)-4-morpholin-4-ylphenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
641	(2S)-N-[4-(2-{{3-(methoxy)-4-morpholin-4-yl}phenyl)amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		12
642	N-(4-{2-[(4-{(2-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
643	N-(4-{2-[(3-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
644	ethyl 3-[(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]amino]-3-oxopropanoate		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
645	N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide		7
646	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide		7
647	N-ethyl-4-{4-[(4-{4-[(tetrahydrofuran-3-ylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}piperazine-1-carboxamide		53





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
648	N-[4-(2-{{4-(4-D-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		7
649	N-[4-(2-{{4-(4-L-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		7
650	N-[4-(2-{{4-(4-D-prolyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
651	N-[4-(2-([4-(4-L-prolylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		7
652	N-{4-[2-(1H-benzimidazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide		5
653	4-(4-furan-2-ylphenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine		9




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
654	N-(4-morpholin-4-ylphenyl)-4-[4-(pyrimidin-2-ylamino)phenyl]pyrimidin-2-amine		9
655	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}-5-methylpyrimidin-4-yl)phenyl]cyclopropanecarboxamide		51
656	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]cyclopropanecarboxamide		51

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
657	N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-methylpyrimidin-4-yl}phenyl)-N,N-dimethylglycinamide		6
658	N ² ,N ² -dimethyl-N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide		12
659	N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide		45
660	N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
661	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide		52
662	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide		52
663	N-ethyl-4-[4-({4-[4-(D-prolylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazine-1-carboxamide		53
664	N-[4-(2-({4-[4-(D-prolylpiperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl)-D-prolinamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
665	N-[4-(2-[[4-(4-L-prolylpiperazin-1-yl)phenyl]amino]pyrimidin-4-yl)phenyl]-D-prolinamide		52
666	1-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)piperidine-2-carboxamide		12
667	N-{4-[2-({4-[4-(piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		7
668	1-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-prolinamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
669	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-4-ylacetamide		12
670	2-(3-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
671	3-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		12
672	2-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
673	2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylpropanamide		12
674	(1R,2R)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylcyclopropanecarboxamide		12
675	2-(4-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
676	3-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		12
677	3-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		12
678	3-(2-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide		678
679	Nalpha,Nalpha-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-phenylalaninamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
680	2-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
681	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-2-ylacetamide		12
682	2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
683	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-{4-[(trifluoromethyl)oxy]phenyl}acetamide		12
684	2-[2-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
685	2-[3-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
686	2-[4-(methoxy)phenyl]-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide		12
687	N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-alaninamide		51
688	N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
689	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-3-(methoxy)propanamide		51
690	(2R)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide		45
691	N ² ,N ² -dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide		12




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
692	1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide		12
693	N ² ,N ² -dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide		12
694	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1-phenylcyclopropanecarboxamide		12
695	2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
696	(2S)-1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-2-carboxamide		12
697	2,4,6-trichloro-N-[3-{[4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino}propyl]benzamide		27
698	N-[3-{[4-[3,4-bis(methoxy)phenyl]pyrimidin-2-yl]amino}propyl]-2,6-dichlorobenzamide		27
699	2,6-dichloro-N-[3-{[4-(4-morpholin-4-ylphenyl)amino]pyrimidin-2-yl]amino}propyl]benzamide		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
700	2,6-dichloro-N-(3-([4-(2,3-dihydro-1,4-benzodioxin-6-yl)-5-fluoropyrimidin-2-yl]amino)propyl)benzamide		27
701	2,6-dichloro-N-{3-[(4-{[(dimethylamino)methyl]phenyl}pyrimidin-2-yl)amino]propyl}benzamide		27
702	2,6-dichloro-N-[3-({4-[3-(1-methylethyl)phenyl]pyrimidin-2-yl}amino)propyl]benzamide		27
703	2,6-dichloro-N-{3-[(4-{4-[(1-methylethyl)oxy]phenyl}pyrimidin-2-yl)amino]propyl}benzamide		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
704	N-[3-({4-[3-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide		27
705	2,6-dichloro-N-[3-({4-[(E)-2-phenylethenyl]pyrimidin-2-yl}amino)propyl]benzamide		27
706	phenyl (4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)carbamate		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
707	phenylmethyl (4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)carbamate		17
708	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide		7
709	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
710	4-{4-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide		7
711	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide		7
712	3-(methyloxy)-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}propanamide		7







Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
713	N-ethyl-4-(4-{[4-(4-{[3-(methoxy)propanoyl]amino}phenyl)pyrimidin-2-yl]amino}phenyl)piperazine-1-carboxamide		53
714	N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-2-phenylacetamide		51
715	1-(4-(2-[(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl)phenyl)pyrrolidin-2-one		25





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
716	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide		52
717	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide		52
718	(2S)-2-hydroxy-3-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide		18




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
719	(2R)-2-hydroxy-3-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide		18
720	N-{4-[2-({4-[4-(cyclopropylcarbonyl)pyrazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide		52
721	(2S)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide		45




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
722	2-amino-2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		14
723	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)morpholine-3-carboxamide		14
724	1-ethyl-3-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methoxy)phenyl]amino}pyrimidin-4-yl)phenyl]urea		53




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
725	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)-3-(methoxy)phenyl}amino}pyrimidin-4-yl)phenyl]-D-prolinamide		24
726	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)-3-(methoxy)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		3
727	1-(2,6-dichlorophenyl)-3-(3-{{4-(4-methyl-2-thienyl)pyrimidin-2-yl}amino}propyl)urea		27
728	1-[2-fluoro-5-(trifluoromethyl)phenyl]-3-(3-{{4-(4-methyl-2-thienyl)pyrimidin-2-yl}amino}propyl)urea		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
729	2,6-dichloro-N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]benzenesulfonamide		36
730	N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-difluorobenzenesulfonamide		36
731	N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]naphthalene-2-sulfonamide		36
732	N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-3,4-bis(methoxy)benzenesulfonamide		36
733	3-chloro-N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]propane-1-sulfonamide		36
734	N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]propane-1-sulfonamide		17




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
735	methyl (3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}propyl)carbamate		17
736	1-methylethyl (3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}propyl)carbamate		17
737	phenylmethyl (3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}propyl)carbamate		17
738	N-{4-[2-({[3-(3-chlorophenyl)isoxazol-5-yl]methyl}amino)pyrimidin-4-yl]phenyl}acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
739	ethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)piperidine-1-carboxylate		17
740	1,1-dimethylethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)piperidine-1-carboxylate		1
741	N-(4-{2-[(4-cyanophenyl)amino]pyrimidin-4-yl}phenyl)acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
742	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyridin-4-yl}phenyl)acetamide		51
743	1,1-dimethylethyl {1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidin-4-yl} carbamate		1
744	N-{4-[2-({4-[4-(cyclopropylcarbonyl)pyrazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
745	N-{1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidin-4-yl}acetamide		1
746	4-(4-aminophenyl)-N-[4-(4-aminopiperidin-1-yl)phenyl]pyrimidin-2-amine		1
747	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}-5-methylpyrimidin-4-yl)phenyl]-3-(methyloxy)propanamide		51



Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
748	N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide		52
749	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide		52
750	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
751	N-cyclopropyl-4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzamide		9
752	N-[2-(methyloxy)ethyl]-4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzamide		9
753	2,6-dichloro-N-{3-[(4-pyridin-3-ylpyrimidin-2-yl)amino]propyl}benzamide		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
754	2,6-dichloro-N-(3-{[4-(4-methyl-3,4-dihydro-2H-1,4-benzoxazin-7-yl)pyrimidin-2-yl]amino}propyl)benzamide		27
755	2,6-dichloro-N-(3-{[4-(2,3-dihydro-1,4-benzodioxin-6-yl)-6-methylpyrimidin-2-yl]amino}propyl)benzamide		27
756	N-(4-{2-[(3-{[(2,6-dichlorophenyl)carbonyl]amino}propyl)amino]pyrimidin-4-yl}phenyl)morpholine-4-carboxamide		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
757	2,6-dichloro-N-{3-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]propyl}benzamide		27
758	N-(4-{2-[(3-{[(2,6-dichlorophenyl)carbonyl]amino}propyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-carboxamide		27
759	2,6-dichloro-N-(3-{[4-(4-{[N-(2-morpholin-4-ylethyl)glycyl]amino}phenyl)pyrimidin-2-yl]amino}propyl)benzamide		27




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
760	1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)ethanone		9
761	(1E)-1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)ethanone oxime		9
762	N-{4-[2-({4-[4-(cyclopropylcarbonyl)pyrazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-2-phenylacetamide		7




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
763	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-bromobenzamide		27
764	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-fluorobenzamide		27





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
765	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-chlorobenzamide		27
766	N-[4-(2-({3-(morpholin-4-ylsulfonyl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide		36
767	N-{4-[2-({3-[(cyclohexylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl} acetamide		28




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
768	N-(4-{2-[(3-[(5-bromo-2-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
769	N-(4-{2-[(3-[(2,5-dimethylphenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		28
770	N-(4-{2-[(3,4-dimorpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		6




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
771	N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide		52
772	N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide		52
773	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
774	N-{4-[2-({4-[4-(cyclobutylcarbonyl)pipe razin-1-yl]phenyl} amino)pyrimi din-4-yl]phenyl} butanamide		52
775	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-2-carboxamide		12
776	2-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
777	3-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12
778	4-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12
779	4-chloro-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
780	(2R)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		51
781	(2S)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		51
782	1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-prolinamide		42
783	N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)thiophene-2-carboxamide		12





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
784	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide		51
785	2-phenyl-N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		52
786	3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(diphenylmethyl)benzamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
787	N-[4-(2-{{4-(4-methylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide		51
788	N-{4-[2-({4-[4-(phenylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
789	N-{4-[2-({4-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
790	N-{4-[2-({4-[4-(cyclohexylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
791	N-(4-{2-[(4-{4-[(2-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
792	N-(4-{2-[(4-{4-[(3-fluorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
793	N-(4-{2-[(4-{4-[(3-fluoro-4-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
794	N-(4-{2-[(4-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
795	N-(4-{2-[(4-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
796	N-[4-(2-[[4-(4-{[3-(methyloxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		7
797	N-(4-{2-[(4-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
798	N-(4-{2-[(4-{4-[(4-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
799	N-(4-{2-[(4-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
800	N-{4-[2-({4-[4-(furan-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
801	N-[4-(2-{[4-(4-{2-[(4-fluorophenyl)oxy]acetyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		7
802	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)furan-3-carboxamide		12
803	N-{4-[2-({4-[4-(phenylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
804	N-{4-[2-({4-[4-(2-thienylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36
805	N-(4-{2-[(4-{4-[(4-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
806	N-[4-(2-{[4-(4-{4-(methyloxy)phenyl]sulfonyl]piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide		36




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
807	N-(4-{2-[(4-{4-[(4-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
808	N-(4-{2-[(4-{4-[(3-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
809	N-{4-[2-({4-[4-(biphenyl-4-yl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		36





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
810	N-{4-[2-({4-[4-(naphthalen-1-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		36
811	N-(4-{2-[(3-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
812	N-[4-(2-{{3-(4-{{3-(methyloxy)phenyl}methyl}piperazin-1-yl}phenyl)amino}pyrimidin-4-yl}phenyl]acetamide		49
813	N-{4-[2-({3-[4-(3-methylbutyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
814	N-{4-[2-({3-[4-(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
815	N-{4-[2-({3-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49
816	N-(4-{2-[(3-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
817	N-(4-{2-[(3-{4-[(4-{[3-(dimethylamino)propyl]oxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		49
818	N-{4-[2-({3-[4-({3-[(trifluoromethyl)oxy]phenyl)methyl}piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		49





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
819	4-[4-((4-[4-(acetylamino)phenyl]pyrimidin-2-yl)amino)phenyl]-N-phenylpiperazine-1-carboxamide		53
820	N-[4-(2-((3-(4-propanoylpiperazin-1-yl)phenyl)amino)pyrimidin-4-yl)phenyl]acetamide		7
821	N-{4-[2-((3-[4-(phenylcarbonyl)piperazin-1-yl]phenyl)amino)pyrimidin-4-yl]phenyl}acetamide		7
822	N-{4-[2-((3-[4-(2-phenylacetyl)piperazin-1-yl]phenyl)amino)pyrimidin-4-yl]phenyl}acetamide		7
823	N-{4-[2-((3-[4-(cyclopentylcarbonyl)piperazin-1-yl]phenyl)amino)pyrimidin-4-yl]phenyl}acetamide		7





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
824	N-{4-[2-({3-[4-(2-pyridin-3-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
825	N-{4-[2-({3-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
826	N-(4-{2-[(3-{4-[(2-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
827	N-(4-{2-[(3-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
828	N-(4-{2-[(3-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
829	N-(4-{2-[(3-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		7
830	N ² ,N ² -dimethyl-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]glycinamide		24




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
831	3-(methyloxy)-N-[4-(2-{{[3-(methyloxy)-4-morpholin-4-yl]phenyl}amino}pyrimidin-4-yl)phenyl]propanamide		24
832	N-(4-{2-[(4-{[(2-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide		36
833	N-{4-[2-(3-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		7
834	N-{4-[2-(3-[4-(2-cyclopropylacetyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl}acetamide		7

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
835	N-[4-(2-{[3-(4-{[3-(methoxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide		7
836	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7
837	1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-3-carboxylic acid		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
838	1,1-dimethylethyl methyl {2-[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]-2-oxoethyl}carbamate		14
839	1,1-dimethylethyl [4-(2-[[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]carbamate		51
840	N-[4-(2-[[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-N ² ,N ² -dimethylglycinamide		51
841	4-(4-aminophenyl)-N-[4-(4-ethylpiperazin-1-yl)phenyl]pyrimidin-2-amine		51



Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
842	Nalpha-{[(1,1-dimethylethyl)oxy]carbonyl}-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-phenylalaninamide	<p>Chiral</p> 	45
843	Nalpha-{[(1,1-dimethylethyl)oxy]carbonyl}-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-phenylalaninamide	<p>Chiral</p> 	45
844	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-phenylalaninamide	<p>Chiral</p> 	45
845	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-phenylalaninamide	<p>Chiral</p> 	45




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
846	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-L-valinamide	<p>Chiral</p> 	45
847	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-D-valinamide	<p>Chiral</p> 	45
848	1-ethyl-3-[4-(2-{{3-(methoxy)-4-morpholin-4-ylphenyl}amino}pyrimidin-4-yl)phenyl]urea		53
849	(2R)-N-(4-(2-{{4-(4-morpholin-4-ylphenyl)amino}pyrimidin-4-yl)phenyl)piperidine-2-carboxamide	<p>Chiral</p> 	45
850	N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}-5-methylpyrimidin-4-yl)phenyl]acetamide		51




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
851	4-{4-[(4-{4-[(N,N-dimethylglycyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide		53
852	N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N ² ,N ² -dimethylglycinamide		52
853	N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N ² ,N ² -dimethylglycinamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
854	N ² ,N ² -dimethyl-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}glycinamide		52
855	N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N ² ,N ² -dimethylglycinamide		52
856	N-[4-(2-({4-(4-D-alanyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl]-N ² ,N ² -dimethylglycinamide		52




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
857	N-[4-(2-([4-(4-L-alanyl)piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-N ² ,N ² -dimethylglycinamide		52
858	N-(4-{2-[(1-[(2,6-dichlorophenyl)carbonyl]azetidin-3-yl)methyl]amino}pyrimidin-4-yl)phenyl)acetamide		41
859	N-(4-{2-[(3-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
860	N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)cyclohexyl]-2,6-dichlorobenzamide		2
861	N-{4-[2-({[4-(4-methylpiperazin-1-yl)phenyl]methyl}amino)pyrimidin-4-yl]phenyl}acetamide		3




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
862	N-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide		2
863	N-{4-[2-(piperidin-4-ylamino)pyrimidin-4-yl]phenyl}acetamide		1
864	N-{4-[2-({1-[(2,6-dichlorophenyl)carbonyl]piperidin-4-yl}amino)pyrimidin-4-yl]phenyl}acetamide		2





Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
865	N-{4-[2-({4-[(2-hydroxyethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		33
866	1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylurea		53
867	N-[5-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)-2-(4-ethylpiperazin-1-yl)phenyl]-2,6-dichlorobenzamide		30






Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
868	1-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-3-(phenylmethyl)urea		53
869	N ² ,N ² -dimethyl-N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}glycinamide		52
870	N-(3-fluoro-4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)cyclopropanecarboxamide		9




Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
871	N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)cyclopropanecarboxamide		52
872	N-[4-(2-{[4-(4-L-alanyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl]acetamide		52
873	N-[4-(2-{[4-(4-L-prolyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl]acetamide		7

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
874	N-[4-(2-([4-(4-D-alanylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide		7
875	N-[4-(2-([4-(4-D-prolylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide		7
876	N-{4-[2-({4-[4-(2-piperazin-1-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide		7

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
877	N-[4-(2-{{4-(4-L-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		52
878	N-[4-(2-{{4-(4-L-prolyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		52
879	N-[4-(2-{{4-(4-D-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		52

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
880	N-[4-(2-{{4-(4-D-prolylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide		52
881	1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1H-pyrrole-2-carboxamide		12
882	3-fluoro-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide		12
883	6-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
884	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridazine-4-carboxamide		12
885	2-cyclopropyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide		12
886	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)isoxazole-5-carboxamide		12
887	N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide		12
888	4-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide		12

Cmpd No.	Name	Structure	Can Be Made According To Synthetic Example #:
889	N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-prolinamide		52
890	N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]butanamide		51
891	1-ethyl-3-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]urea		53

5

[0120] In another embodiment, the compound of the invention is selected from a compound from Table 2:

Table 2

10 N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide;

- 5 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopropanecarboxamide;
N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopropane-
carboxamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)valinamide;
N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-
10 yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide;
N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide;
N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-
acetamide;
15 2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-
phenylacetamide;
N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide;
3-(methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-
yl}phenyl)propanamide;
20 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide;
N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-
yl}phenyl)amino]pyrimidin-
4-yl}phenyl)acetamide;
25 N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-
yl}phenyl)-acetamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide;
N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]-D-
prolinamide;
30 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-
carboxamide;
O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide;
1-ethyl-3-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-
yl]phenyl}urea;
35 N-ethyl-4-(4-{[4-(4-{[(ethylamino)carbonyl]amino}phenyl)pyrimidin-2-yl]amino}phenyl)-
piperazine-1-carboxamide;

- 5 N-2~,N-2~-dimethyl-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)glycinamide;
 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
 (3R)-3-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide;
 10 (3S)-3-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide;
 N-{4-[2-[(4-{4-(tetrahydrofuran-2-ylcarbonyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}-D-prolinamide;
 2-hydroxy-2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide;
 15 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide;
 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide;
 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide;
 O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide;
 20 O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
 O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
 and
 N-(4-{2-[(3-fluoro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide.

[0121] In another embodiment, the invention relates to a pharmaceutical composition
 25 comprising a compound according to Table 1 or Table 2, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, excipient, or diluent.

[0122] In another embodiment, the invention relates to a method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a compound according to Table 1 or Table 2, or a pharmaceutically salt thereof.

30 [0123] In another embodiment, the invention relates to a method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a pharmaceutical composition comprising a compound according to Table 1 or Table 2, or a pharmaceutically acceptable salt thereof.

[0124] In another embodiment, the invention relates to a method for treating a disease,
 35 disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a therapeutically effective

5 amount of a compound according to Table 1 or Table 2, or a pharmaceutically acceptable salt thereof.

[0125] In another embodiment, the invention relates to a method for treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a pharmaceutical composition
10 comprising a therapeutically effective amount of a compound according to Table 1 or Table 2, or a pharmaceutically acceptable salt thereof.

General Administration

[0126] In certain other preferred embodiments, administration can preferably be by the
15 oral route. Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally (intravenous, intramuscular, or subcutaneous), topically, transdermally, intravaginally, intravesically,
20 intracisternally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages.

[0127] The compositions will include a conventional pharmaceutical carrier or excipient
25 and a compound of the invention as the/an active agent, and, in addition, can include carriers and adjuvants, etc.

[0128] Adjuvants include preserving, wetting, suspending, sweetening, flavoring, perfuming, emulsifying, and dispensing agents. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, parabens,
30 chlorobutanol, phenol, sorbic acid, and the like. It can also be desirable to include isotonic agents, for example sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0129] If desired, a pharmaceutical composition of the invention can also contain minor
35 amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, etc.

5 [0130] The choice of formulation depends on various factors such as the mode of drug administration (e.g., for oral administration, formulations in the form of tablets, pills or capsules are preferred) and the bioavailability of the drug substance. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

[0131] Compositions suitable for parenteral injection can comprise physiologically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, and sterile powders for reconstitution into sterile injectable solutions or dispersions.

20 Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

[0132] One preferable route of administration is oral, using a convenient daily dosage regimen that can be adjusted according to the degree of severity of the disease-state to be treated.

[0133] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is admixed with at least one inert customary excipient (or carrier) such as sodium citrate or dicalcium phosphate or (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, cellulose derivatives, starch, alginates, gelatin, polyvinylpyrrolidone, sucrose, and gum acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, croscarmellose sodium, complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary

5 ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, magnesium stearate and the like (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms can also comprise buffering agents.

10 [0134] Solid dosage forms, as described above, can be prepared with coatings and shells, such as enteric coatings and others well known in the art. They can contain pacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedded compositions that can be used are polymeric substances and waxes. The active compounds can also be in
15 microencapsulated form, if appropriate, with one or more of the above-mentioned excipients.

[0135] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. Such dosage forms are prepared, for example, by dissolving, dispersing, etc., a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and optional pharmaceutical adjuvants in a carrier,
20 such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like; solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butylene glycol, dimethylformamide; oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol,
25 polyethyleneglycols and fatty acid esters of sorbitan; or mixtures of these substances, and the like, to thereby form a solution or suspension.

[0136] Suspensions, in addition to the active compounds, can contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or
30 mixtures of these substances, and the like.

[0137] Compositions for rectal administrations are, for example, suppositories that can be prepared by mixing the compounds of the present invention with for example suitable non-irritating excipients or carriers such as cocoa butter, polyethyleneglycol or a suppository wax, which are solid at ordinary temperatures but liquid at body temperature and therefore, melt
35 while in a suitable body cavity and release the active component therein.

[0138] Dosage forms for topical administration of a compound of this invention include ointments, powders, sprays, and inhalants. The active component is admixed under sterile

5 conditions with a physiologically acceptable carrier and any preservatives, buffers, or propellants as can be required. Ophthalmic formulations, eye ointments, powders, and solutions are also contemplated as being within the scope of this invention.

[0139] Compressed gases can be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

10 [0140] Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. In one example, the composition will be between about 5% and about 75% by weight of a compound(s) of the invention, or a
15 pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

[0141] Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, 18th Ed., (Mack Publishing Company, Easton, Pa., 1990). The composition to be administered will, in
20 any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state in accordance with the teachings of this invention.

[0142] The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount which will vary depending upon a variety
25 of factors including the activity of the specific compound employed, the metabolic stability and length of action of the compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular disease-states, and the host undergoing therapy. The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per
30 day. For a normal human adult having a body weight of about 70 kilograms, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is an example. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum
35 dosages for a particular patient is well known to one of ordinary skill in the art.

[0143] The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, can include other

5 medicinal agents and pharmaceutical agents. Compositions of the invention can be used in combination with anticancer and/or other agents that are generally administered to a patient being treated for cancer, e.g. surgery, radiation and/or chemotherapeutic agent(s).

Chemotherapeutic agents that can be useful for administration in combination with compounds of Formula I in treating cancer include alkylating agents, platinum containing
10 agents.

[0144] If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described above and the other pharmaceutically active agent(s) within its approved dosage range. Compounds of the instant invention can alternatively be used sequentially with known pharmaceutically acceptable agent(s) when a
15 combination formulation is inappropriate.

[0145] Representative pharmaceutical formulations containing a compound of Formula I are described below.

UTILITY

[0146] The compounds of this invention are JAK-2 inhibitors. As such the compounds of
20 Formula I are useful for treating diseases, particularly myeloproliferative disorders, for example, myelofibrosis, thrombocythemia, polycythemia vera (PV), essential thrombocythemia (ET), agnogenic myeloid metaplasia (AMM), also referred to as idiopathic myelofibrosis (IMF), and chronic myelogenous leukemia (CML); and cancer, for example, ovarian cancer, cervical cancer, breast cancer, colorectal cancer, glioblastomas, prostate,
25 colon, melanoma, leukemia and haematopoietic malignancies, as described above, in which JAK-2 activity contributes to the pathology and/or symptomatology of the disease.

[0147] Suitable *in vitro* assays for measuring JAK-2 activity and the inhibition thereof by compounds are known. For further details of an *in vitro* assay for measuring JAK-2 activity see Biological Examples, Example 1 *infra*.

30 [0148] Assays for measurement of efficacy in treatment of various cancers are described in Biological Examples, Example 3, 5, and 6, *infra*.

[0149] Suitable *in vivo* models of various cancers are known to those of ordinary skill in the art. For further details of *in vivo* assays see Biological Examples 2 and 4, *infra*.

Utility of compounds of the invention as screening agents

35 [0150] To employ the compounds of the invention in a method of screening for candidate agents that bind to, for example JAK-2, the protein is bound to a support, and a compound of

5 the invention is added to the assay. Alternatively, the compound of the invention is bound to the support and the protein is added. Classes of candidate agents among which novel binding agents can be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays can be
10 used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[0151] The determination of the binding of the candidate agent to, for example, JAK-2 can be done in a number of ways. In one example, the candidate agent (the compound of the
15 invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this can be done by attaching all or a portion of the JAK-2 protein to a solid support, adding a labeled agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support.
20 Various blocking and washing steps can be utilized as is known in the art.

[0152] The term "labeled" as used herein is meant to include both direct and indirect labeling with a compound that provides a detectable signal, for example, radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, and the like. Specific binding molecules include pairs,
25 such as biotin and streptavidin, digoxin and antidigoxin, and the like. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[0153] In some embodiments, only one of the components is labeled. For example, JAK-2
30 protein can be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component can be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the candidate agents.

[0154] The compounds of the invention can also be used as competitors to screen for additional drug candidates. The terms "candidate bioactive agent" or "drug candidate" or
35 grammatical equivalents as used herein describe any molecule, *e.g.*, protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They can be capable of directly or indirectly altering the cellular proliferation phenotype or

5 the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. In the case where protein binding or activity is screened, some embodiments exclude molecules already known to bind to that particular protein. Exemplary
10 embodiments of assays described herein include candidate agents, which do not bind the target protein in its endogenous native state, termed herein as "exogenous" agents. In one example, exogenous agents further exclude antibodies to JAK-2.

[0155] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules having a molecular weight of more than about 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural
15 interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, for example at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclyl structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules
20 including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs, or combinations thereof.

[0156] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including
25 expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents can be subjected to directed or random chemical modifications, such as acylation,
30 alkylation, esterification, amidification to produce structural analogs.

[0157] In one example, the binding of the candidate agent is determined through the use of competitive binding assays. In this example, the competitor is a binding moiety known to bind to IGF1R, such as an antibody, peptide, binding partner, ligand, etc. Under certain
35 circumstances, there can be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[0158] In some embodiments, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to JAK-2 protein for a time sufficient to allow binding,

5 if present. Incubations can be performed at any temperature that facilitates optimal activity, typically between 4°C and 40°C.

[0159] Incubation periods are selected for optimum activity, but can also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is
10 then added, and the presence or absence of the labeled component is followed, to indicate binding.

[0160] In one example, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to JAK-2 and thus is capable of binding to, and potentially modulating the activity of the JAK-2. In this
15 embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[0161] In an alternative embodiment, the candidate agent is added first, with incubation
20 and washing, followed by the competitor. The absence of binding by the competitor can indicate the candidate agent is bound to JAK-2 with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, can indicate the candidate agent is capable of binding to JAK-2.

[0162] It can be of value to identify the binding site of JAK-2. This can be done in a
25 variety of ways. In one embodiment, once JAK-2 is identified as binding to the candidate agent, the JAK-2 is fragmented or modified and the assays repeated to identify the necessary components for binding.

[0163] Modulation is tested by screening for candidate agents capable of modulating the activity of JAK-2 comprising the steps of combining a candidate agent with JAK-2, as above,
30 and determining an alteration in the biological activity of the JAK-2. Thus, in this embodiment, the candidate agent should both bind to (although this can not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both *in vitro* screening methods and *in vivo* screening of cells for alterations in cell viability, morphology, and the like.

35 [0164] Alternatively, differential screening can be used to identify drug candidates that bind to native JAK-2, but cannot bind to modified JAK-2.

5 [0165] Positive controls and negative controls can be used in the assays. For example, all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of samples is for a time sufficient for the binding of the agent to the protein. Following incubation, samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a
10 radiolabel is employed, the samples can be counted in a scintillation counter to determine the amount of bound compound.

[0166] A variety of other reagents can be included in the screening assays. These include reagents like salts, neutral proteins, *e.g.*, albumin, detergents, etc which can be used to facilitate optimal protein-protein binding and/or reduce non-specific or background
15 interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., can be used. The mixture of components can be added in any order that provides for the requisite binding.

[0167] One of ordinary skill in the art would understand that certain crystallized, protein-ligand complexes, in particular JAK-2-ligand complexes, and their corresponding x-ray
20 structure coordinates can be used to reveal new structural information useful for understanding the biological activity of JAK-2 kinase's as described herein. As well, the key structural features of the aforementioned proteins, particularly, the shape of the ligand binding site, are useful in methods for designing or identifying selective modulators of JAK-2 kinase's and in solving the structures of other proteins with similar features. Ligands of such
25 complexes can include compounds of the invention as described herein.

[0168] As well, one of ordinary skill in the art would appreciate that such suitable x-ray quality crystals can be used as part of a method of identifying a candidate agent capable of binding to and modulating the activity of JAK-2 kinases. Such methods can be characterized by the following aspects: a) introducing into a suitable computer program, information
30 defining a ligand binding domain of a JAK-2 kinase in a conformation (*e.g.* as defined by x-ray structure coordinates obtained from suitable x-ray quality crystals as described above) the computer program creates a model of the three dimensional structures of the ligand binding domain, b) introducing a model of the three dimensional structure of a candidate agent in the computer program, c) superimposing the model of the candidate agent on the model of the
35 ligand binding domain, and d) assessing whether the candidate agent model fits spatially into the ligand binding domain. Aspects a-d are not necessarily carried out in the aforementioned order. Such methods can further entail: performing rational drug design with the model of the

5 three-dimensional structure, and selecting a potential candidate agent in conjunction with computer modeling.

[0169] Additionally, one skilled in the art would appreciate that such methods can further entail: employing a candidate agent, so-determined to fit spatially into the ligand binding domain, in a biological activity assay for JAK-2 kinase modulation, and determining whether
10 said candidate agent modulates JAK-2 kinase activity in the assay. Such methods can also include administering the candidate agent, determined to modulate JAK-2 kinase activity, to a mammal suffering from a condition treatable by JAK-2 kinase modulation, such as those described above.

[0170] Also, one skilled in the art would appreciate that compounds of the invention can
15 be used in a method of evaluating the ability of a test agent to associate with a molecule or molecular complex comprising a ligand binding domain of a JAK-2 kinase. Such a method can be characterized by the following aspects: a) creating a computer model of a JAK-2 kinase binding pocket using structure coordinates obtained from suitable x-ray quality crystals of the JAK-2 kinase, b) employing computational algorithms to perform a fitting
20 operation between the test agent and the computer model of the binding pocket, and c) analyzing the results of the fitting operation to quantify the association between the test agent and the computer model of the binding pocket.

[0171] The disclosures in this application of all articles and references, including patents, are incorporated herein by reference.

25 **Synthetic Procedures**

[0172] The compounds of the invention, or their pharmaceutically acceptable salts, can have asymmetric carbon atoms, oxidized sulfur atoms or quaternized nitrogen atoms in their structure.

[0173] The compounds of the invention and their pharmaceutically acceptable salts can
30 exist as single stereoisomers, racemates, and as mixtures of enantiomers and diastereomers. The compounds can also exist as geometric isomers. All such single stereoisomers, racemates and mixtures thereof, and geometric isomers are intended to be within the scope of this invention.

[0174] It is assumed that when considering generic descriptions of compounds of the
35 invention for the purpose of constructing a compound, such construction results in the creation of a stable structure. That is, one of ordinary skill in the art would recognize that

5 theoretically some constructs which would not normally be considered as stable compounds (that is, sterically practical and/or synthetically feasible, *supra*).

[0175] Methods for the preparation and/or separation and isolation of single stereoisomers from racemic mixtures or non-racemic mixtures of stereoisomers are well known in the art. For example, optically active (R)- and (S)- isomers can be prepared using chiral synthons or
10 chiral reagents, or resolved using conventional techniques. Enantiomers (R- and S-isomers) can be resolved by methods known to one of ordinary skill in the art, for example by: formation of diastereoisomeric salts or complexes which can be separated, for example, by crystallization; via formation of diastereoisomeric derivatives which can be separated, for example, by crystallization, selective reaction of one enantiomer with an enantiomer-specific
15 reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where a desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a
20 further step can be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer can be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents or by converting one enantiomer to the other by asymmetric transformation. For a mixture of enantiomers, enriched in a particular enantiomer, the major component enantiomer can be further enriched (with concomitant loss in yield) by
25 recrystallization.


[0176] In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

30 [0177] In addition, it is intended that the present invention cover compounds made either using standard organic synthetic techniques, including combinatorial chemistry or by biological methods, such as bacterial digestion, metabolism, enzymatic conversion, and the like.

[0178] Scheme 1 depicts the general synthetic procedure for the compounds of the
35 invention. Synthesis of the compounds of the invention is not limited by the procedure of Scheme 1. One skilled in the art will know that other procedures can be used to synthesize the compounds of the invention, and that the procedure described in Scheme 1 is only one such

5 procedure. In the descriptions below, one of ordinary skill in the art would recognize that specific reaction conditions, added reagents, solvents, and reaction temperatures can be modified for the synthesis of specific compounds of the invention. Thus, the general synthetic procedure depicted in Scheme 1 in conjunction with the specific examples that follow provide sufficient information and guidance to allow one of ordinary skill in the art to
 10 synthesize compounds of the invention.

[0179] Compounds of formula I can be prepared according to Scheme 1:



15 [0180] The synthesis of compounds of Formula I proceeds from commercially available reagents and employs standard techniques. Standard Suzuki coupling reactions conditions can be used to convert dichloropyrimindines of formula A (commercially available from Sigma Aldrich) and boronic acids of formula B (commercially available from Sigma Aldrich,
 Fisher Scientific, or Combi-Blocks Inc.), where R^{25} , Z and $n1$ are as defined in the
 20 Detailed Description of the Invention, to 4-substituted-2-chloropyrimindes of formula C.

- 5 Compounds of Formula **D1** and **I** can be generated by reaction of **C** with the corresponding amines (**F1**, available from Fluka) or anilines (**F2**, available from Sigma Aldrich).
 Compounds of formula **D1** can be further transformed to amides of formula **E** using standard peptide coupling conditions with carboxylic acids or reaction with acid chlorides. For instance, **D1** can be reacted with an intermediate of formula $LG^1C(O)R^4$ where LG^1 is a
 0 leaving group under acylation conditions and R^4 is phenyl optionally substituted with 1, 2, 3, 4, or 5 R^{11} groups, wherein R^{11} is as defined in the Detailed Description of the Invention to yield a compound of formula **E**.


Examples

- [0181] The following examples serve to more fully describe the manner of making the
 5 compounds of the invention, as well as to set forth the best modes contemplated for carrying out the invention. These examples in no way serve to limit the scope of the invention, but rather are presented for illustrative purposes. All references cited herein are incorporated by reference in their entirety. Generally, each example is set out below with a corresponding multi-step synthesis procedure. Following the specific examples is a list of compounds that
 10 were made in a similar way.

Example 1


N-(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 58)

a) *N*-(4-(2-chloropyrimidin-4-yl)phenyl)acetamide (**C₁**)



- 25 [0182] A flask was charged with 2,4-dichloropyrimidine **A₁** (650 mg, 4.4 mmol), 4-acetoamidophenylboronic acid **B₁** (820 mg, 4.6 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (480 mg, 0.56 mmol, 15 mol %), and triethylamine (1.5 mL, 11 mmol). Ethyleneglycoldimethylether (30 mL) was added to the flask and the mixture was purged with N_2 for 5 minutes. The reaction mixture was stirred under an N_2 atmosphere at 80
 30 °C for 12 hours, after which time, ether was added and the reaction mixture was filtered. The product, **C₁**, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: m/z 248 ($M+H$)⁺.

b) *N*-(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)acetamide (**58**)




[0183] A flask containing a solution of **C₁** (500 mg, 2.0 mmol) and 3-boc-amino-aniline **F** (687 mg, 3.3 mmol) in nBuOH (5 mL) was immersed in an oil bath at 180 °C for 30 mins. The mixture was cooled to ambient temperature and to the black residue was added aqueous HCl and MeOH. The aqueous layer was twice washed with ethylacetate. The aqueous layer was then basified with NaOH and extracted twice with ethylacetate. The organic layer was washed with brine and dried with sodium sulfate. The solvent was removed on a rotary evaporator and the product was purified by HPLC with TFA/ACN as eluent. The TFA salt was removed by extraction with sodium hydroxide and ethylacetate to afford the title compound **58**.

10

15

Example 2

N-[3-({4-[4-(acetamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide (Compound 7)



[0184] A flask was charged with **58** (638 mg, 2.0 mmol), 2,6-dichlorobenzoylchloride **G** (350 μ L, 2.4 mmol), diisopropylethylamine (1.1 mL, 6 mmol) and THF (50 mL). The reaction mixture was stirred at 70 °C for 6 hours. The crude mixture was concentrated on a rotary evaporator and the crude product was purified by HPLC with TFA/ACN as eluent. The title compound **7** was isolated by precipitation from ACN and washed with ether.


25

¹H-NMR (400MHz, d₆-DMSO): 10.718 ppm (s, 1H), 10.269 ppm (s, 1H), 9.678 ppm (s, 1H), 8.507 ppm (d, 1H), 8.419 ppm (s, 1H), 8.215 ppm (d, 2H), 7.758 ppm (d, 2H), 7.608 ppm (d, 2H), 7.532 ppm (t, 1H), 7.472 ppm (d, 1H), 7.380 ppm (d, 1H), 7.301 ppm (t, 1H), 7.216 ppm (d, 1H), 2.085 ppm (s, 3H); MS (EI) C₂₅H₁₉Cl₂N₅O₂: 492.2 (MH⁺).

5

Example 3

N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide (18)




[0185] A flask was charged with **C₁** (500 mg, 2.0 mmol), 4-morpholinylaniline **H** (540 mg, 3.0 mmol) and *n*BuOH (10 mL). The flask was immersed in a 180 °C oil bath for 30 minutes. The reaction mixture was cooled to ambient temperature and the black residue dissolved in DMF and MeOH. The product was purified by HPLC with TFA/ACN as eluent. The TFA salt was removed by extracting with sodium hydroxide and ethylacetate to afford the title compound (**18**).

5 ¹H-NMR (400MHz, d₆-DMSO): 10.533 ppm (s, 1H), 9.408 ppm (s, 1H), 8.447 ppm (d, 1H), 8.114 ppm (d, 2H), 7.813 ppm (d, 2H), 7.705 ppm (d, 2H), 7.288 ppm (d, 1H), 6.982 ppm (br s, 2H), 4.65 ppm (br s, 4H), 3.072 ppm (br s, 2H), 2.108 ppm (s, 3H); MS (EI) C₂₂H₂₃N₅O₂: 390.3 (MH⁺).

10

Example 4

N-{1-[(2,6-dichlorophenyl)carbonyl]piperidin-4-yl}-4-(4-methyl-2-thienyl)pyrimidin-2-amine



5 [0186] To a solution of {4-[4-(5-Methyl-thiophen-2-yl)-phenyl]-pyrimidin-2-yl}-piperidin-4-yl-amine hydrochloride **I** (274mg, 1 mmol) and TEA (0.69mL, 5 mmol) in DMF (5 mL) was added 2,6-dichlorobenzoyl chloride **G** (0.21mL, 1.5 mmol) and the solution was stirred for 4 h. To the resulting solution was added ethyl acetate (100mL) and the organic

5 layer was washed with 5% LiCl (3 x 50mL), dried over anhydrous sodium sulfate, filtered, and concentrated to yield a residue. This residue was purified by reverse phase HPLC to yield the product **G2** (195mg, 38.9% yield, acetate salt) as a tan solid.


¹H NMR (400 MHz, *d*₆-DMSO): 8.28 (m, 1H), 7.73 (m, 1H), 7.58-7.54 (m, 2H), 7.48-7.46 (m, 1H), 7.32 (s, 1H), 7.27 (m, 1H), 7.01 (m, 1H), 4.47 (m, 1H), 4.03 (m, 1H), 3.30-3.05 (m, 10 2H), 2.25 (s, 3H), 2.03 (m, 1H), 1.88 (m, 1H), 1.58-1.48 (m, 3H); MS (EI) for C₂₁H₂₀Cl₂N₄OS: 447 (MH⁺).

Example 5

N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide

15 (Compound 574)

a) N-(4-(2-chloro-5-methylpyrimidin-4-yl)phenyl)acetamide (C₂)




[0187] A flask was charged with 5-methyl-2,4-dichloropyrimidine **C₂** (2.45g, 15.0mmol), 4-acetoamidophenylboronic acid (2.95g, 16.5mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium **B₁** (1.22g, 1.5mmol, 10 mol %), and triethylamine (5.23mL, 37.5mmol). Ethyleneglycoldimethylether (20 mL) and H₂O (5ml) were added to the flask and the mixture was purged with N₂ for 5 minutes. The reaction mixture was stirred under an N₂ atmosphere at 90 °C for 2hours, after which time, ether was added and the reaction mixture was filtered. The product, **C₂**, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: m/z 262 (M+H)⁺.

20

25

b) N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide



30 [0188] A flask containing a solution of **C₂** (523 mg, 2.0 mmol) and **H** (392 mg, 2.2mmol) in *n*-BuOH (6 mL) was immersed in an oil bath at 180 °C for 3hr. The mixture was cooled to


5 ambient temperature and concentrated *in vacuo*. The residue was purified by HPLC with Ammonium acetate/ACN as eluent to afford the title compound **574** (531mg, 66%).

¹H-NMR (400MHz, d₆-DMSO): 10.15 ppm (s, 1H), 9.27 ppm (s, 1H), 8.31 ppm (s, 1H), 7.72 ppm (d, J = 8.8Hz, 2H), 7.66-7.62 ppm (m, 4H), 6.88 ppm (d, J = 8.8Hz, 2H), 3.73 (t, J = 4.8Hz, 4H), 3.01 (t, J = 4.8Hz, 4H), 2.21 ppm (s, 3H), 2.09 (s, 3H); MS (EI) C₂₃H₂₅N₅O₂:

10 404 (M+H)⁺.

Example 6

N-(4-(2-(3,5-dimorpholinophenylamino)-5-methylpyrimidin-4-yl)phenyl)acetamide
(Compound **570**)




[0189] A flask containing a solution of **C₂** (288 mg, 1.1 mmol) and **H₁** (263 mg, 1.0mmol) in n-BuOH (3 mL) was immersed in an oil bath at 180 °C for 4hr. The mixture was cooled to ambient temperature and concentrated *in vacuo*. The residue was purified by HPLC with ammonium acetate/acetonitrile (ACN) as eluent to afford the title compound **570** (205mg, 42%).

25 ¹H-NMR (400MHz, d₆-DMSO): 10.15 ppm (s, 1H), 9.22 ppm (s, 1H), 8.34 ppm (s, 1H), 7.72 ppm (d, J = 9.2Hz, 2H), 7.69 ppm (d, J = 8.8Hz, 2H), 7.10 ppm (d, J = 2.0Hz, 2H), 6.09 ppm (s, 1H), 3.71 (t, J = 4.8Hz, 8H), 3.03 (t, J = 4.8Hz, 8H), 2.26 ppm (s, 3H), 2.07 (s, 3H); MS (EI) C₂₇H₃₂N₆O₃: 489 (M+H)⁺.

5

Example 7

N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide



10

[0190] A solution of N-(4-(2-(4-(piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide (300mg, 0.6mmol) and DIPEA (261ul, 1.50mmol) was treated with isobutyryl chloride at room temperature. After stirring for 10 minutes, the reaction mixture was directly concentrated *in vacuo* and the residue was purified by HPLC TFA/ACN as eluent. TFA salt was removed by using basic resin to afford 111mg (40%) of the title compound 572.

15


¹H-NMR (400MHz, d₆-DMSO): 10.21 ppm (s, 1H), 9.40 ppm (s, 1H), 8.44 ppm (d, J = 4.8Hz, 1H), 8.11 ppm (d, J = 9.2Hz, 2H), 8.10 ppm (s, 1H), 7.74 ppm (d, J = 8.8Hz, 2H), 7.68 ppm (d, J = 8.8Hz, 1H), 7.28 ppm (d, J = 5.6Hz, 1H), 6.97 (d, J = 9.6Hz, 2H), 3.65-3.61 (m, 4H), 3.08-3.02 (m, 4H), 2.92 ppm (penth, J = 6.8Hz, 1H), 2.09 ppm (s, 3H), 1.03 ppm (s, 3H), 1.01 ppm (s, 3H); MS (EI) C₂₆H₃₀N₆O₂: 459 (M+H)⁺.

20

Example 8

Methyl (4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)carbamate
(Compound 248)

25



[0191] To a solution of 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine 55 (100 mg, 0.29 mmol) and DIEA (0.435 mmol, 75 μl) in THF (50 mL) was added methyl


5 chloroformate (0.348 mmol, 27 μ l) and the solution was stirred at room temperature for 2 hours. The solution mixture was concentrated, redissolved with MeOH and purified using reverse phase HPLC. The product obtained from the reverse phase HPLC was free base **248**, converted to HCl salt using 3 N HCl and lyophilized to yield the product **248** (60mg, 47% yield) as a yellow solid.

0 $^1\text{H-NMR}$ (400MHz, d_6 -DMSO): 10.063 ppm (s, 1H), 9.976 ppm (s, 1H), 8.521 ppm (d, 1H), 8.153 ppm (d, 2H), 7.878 ppm (d, 2H), 7.661 ppm (d, 2H), 7.554 ppm (bs, 2H), 7.432 ppm (d, 1H), 3.983 ppm (bs, 4H), 3.707 ppm (s, 3H), 4.435 ppm (bs, 4H); MS (EI) $\text{C}_{22}\text{H}_{23}\text{N}_5\text{O}_3\text{HCl}$: 475.4 (MH^+).

15 Example 9


4-[4-(dimethylamino)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (67)

a) 4-(2-chloropyrimidin-4-yl)-N,N-dimethylaniline (C_3)



20 [0192] A flask was charged with A_1 (650 mg, 4.4 mmol), 4-(dimethylamino)phenylboronic acid B_2 (797 mg, 4.8 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (480 mg, 0.56 mmol, 15 mol %), and triethylamine (1.5 mL, 11 mmol). Ethyleneglycoldimethylether (30 mL) was added to the flask and the mixture was purged with N_2 for 5 minutes. The reaction mixture was stirred under an N_2 atmosphere at 80 $^\circ\text{C}$ for 12 hours, after which time, ether was added and the reaction mixture was filtered. The product, C_3 , was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: m/z 234 ($\text{M}+\text{H}^+$).

25 b) 4-[4-(dimethylamino)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (67)




[0193] A flask was charged with **C₃** (500 mg, 2.1 mmol), 4-morpholinoaniline (573 mg, 3.2 mmol) and nBuOH (10 mL). The flask was immersed in a 180 °C oil bath for 30 minutes. The reaction mixture was cooled to ambient temperature and the black residue dissolved in DMF and MeOH. The product **67** was purified by HPLC with TFA/ACN as eluent. The TFA salt was removed by extracting with sodium hydroxide and ethylacetate to afford the free base of **67**.

¹H-NMR (400MHz, d₆-DMSO): 9.24 ppm (s, 1H), 8.33 (d, 1H), 8.03 (d, 2H), 7.68 (d, 2H), 7.18 (d, 1H), 6.92 (d, 2H), 6.81 (d, 2H), 3.72-3.77 (m, 4H), 3.04-3.08 (m, 4H), 3.00 (s, 6H). MS (EI) C₂₂H₂₅N₅O: 376.1 (MH⁺).

Example 10


4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (Compound 319)

a) 4-(2-chloropyrimidin-4-yl)benzonitrile




[0194] A flask was charged with **A₁** (763 mg, 5.16 mmol), 4-cyanophenylboronic acid **B₃** (848 mg, 5.77 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (375 mg, 0.437 mmol, 10 mol %), and triethylamine (1.76 mL, 12.9 mmol). Ethylene glycol dimethyl ether (5.0 mL) was added to the flask and the mixture was purged with N₂. The reaction mixture was stirred under an N₂ atmosphere at 90 °C for 1 hour, after which time, it was cooled to ambient temperature and filtered. The product, **C₄**, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: m/z 216 (M+H)⁺.

5 b) 4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzonitrile (**S**)




[0195] A flask containing a solution of **C₄** (400 mg, 1.85 mmol) and 4-morpholinoaniline (362 mg, 2.04 mmol) in 1-butanol (10 mL) was immersed in an oil bath at 180 °C for 2 h. The mixture was cooled to ambient temperature, concentrated, and the crude product **S** was used without further purification. LCMS: m/z 358 ($M+H$)⁺.

c) 4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzoic acid



[0196] A flask containing a solution of **S** (600 mg, 1.68 mmol) and 10 N HCl (aq., 20 mL) was immersed in an oil bath at 100 °C for 5 hours. The mixture was cooled to ambient temperature, after which time, 5 N LiOH was added until the reaction mixture was pH 6. The white precipitate was filtered and dried to give the product **T**, which was used without further purification. LCMS: m/z 377 ($M+H$)⁺.

d) 4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (**323**)




[0197] To a flask containing a solution of **T** (570 mg, 1.47 mmol) and THF (10 mL) was added 1,1'-carbonyldiimidazole (475 mg, 2.93 mmol). The reaction mixture was immersed in an oil bath at 60 °C for 2 h, after which time, it was cooled to ambient temperature. A mixture of acetamide oxime (120 mg, 1.62 mmol) and NaH (39 mg, 1.6 mmol) in DMF (5 mL) was added to the reaction mixture, after which time, the reaction mixture was immersed in an oil

- 5 bath at 80 °C for 2 hours. The reaction mixture was then cooled to ambient temperature, quenched with saturated NH₄Cl (aq., 10 mL), extracted with ethyl acetate (50 mL), dried over anhydrous sodium sulfate, filtered, and concentrated to yield a residue. The residue was purified by reverse phase HPLC to yield the product **319** (49.6 mg, 8.10% yield) as a light brown solid.
- 10 ¹H-NMR (400MHz, d₆-DMSO): 9.57 ppm (s, 1H), 8.57 ppm (d, 1H), 8.38 ppm (d, 2H), 8.25 ppm (d, 2H), 7.67 ppm (d, 2H), 7.44 ppm (d, 1H), 6.95 ppm (d, 2H), 3.75 ppm (t, 4H), 3.06 ppm (t, 4H), 2.46 ppm (s, 3H); MS (EI) C₂₃H₂₂N₆O₂: 415.0 (MH⁺).

5

Example 11**N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyrrolidin-1-ylacetamide (Compound 292)**

a) 2-chloro-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide




10

[0198] To a flask charged with 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine (100 mg, 0.286 mmol) and THF (1 mL) was added chloroacetyl chloride (0.0230 mL, 0.286 mmol). The solution was stirred at ambient temperature for 1 hour. The crude mixture was then concentrated and used without further purification. LCMS: m/z 424 (M+H)⁺.

15

b) N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyrrolidin-1-ylacetamide (292)




[0199] To a flask charged with U (100 mg, 0.236 mmol), diisopropylethylamine (0.2 mL, 1 mmol), and dimethylacetamide (1 mL) was added pyrrolidine (0.021 mL, 1.3 mmol). The reaction mixture was stirred at 80 °C for 1 hour. The crude mixture was concentrated on a rotary evaporator and the product 292 was purified by reverse phase HPLC.

¹H-NMR (400MHz, d₆-DMSO): 10.90 ppm (s, 1H), 10.20 ppm (br. s, 1H), 9.64 ppm (s, 1H), 8.50 ppm (d, 1H), 8.19 ppm (d, 2H), 7.78 ppm (d, 2H), 7.74 ppm (d, 2H), 7.36 ppm (d, 1H), 7.11 ppm (d, 2H), 4.32 ppm (s, 2H), 3.80 ppm (t, 4H), 3.70-3.65 ppm (m, 2H), 3.19-3.06 ppm (m, 6H), 2.10-1.86 (m, 4H); MS (EI) C₂₆H₃₀N₆O₂: 459.4 (MH⁺).

25

5

Example 12

**3-methoxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide
(Compound 575)**

[0200] To a solution of **249(a)** (0.18 g, 0.05 mmol), HATU (0.4 g, 1.1 mmol), and DIEA (0.5 mL, 4.0 mmol) in DMA (5 mL) was added 3-methoxypropanoic acid (0.1 mL, 1.05 mmol) and the solution was stirred at 60 °C for 2 hours. The solution mixture was diluted with ethyl acetate and the mixture was extracted with 10% LiCl (3X) and brine (1X). The resulting organic layer was dried with sodium sulfate and concentrated *in vacuo*. The product was purified by silica column chromatography (5% MeOH/DCM as eluent) to afford 0.1 g of the title compound **575** (49% yield) as a white solid.


¹H-NMR (400MHz, d₆-DMSO): 10.20 ppm (s, 1H), 9.37 ppm (s, 1H), 8.42 ppm (d, 1H), 8.10 ppm (d, 2H), 7.74 ppm (d, 2H), 7.65 ppm (d, 2H), 7.25 ppm (d, 1H), 6.91 ppm (d, 2H), 3.72 ppm (m, 4H), 3.61 ppm (t, 2H), 3.23 ppm (s, 3H), 3.03 ppm (m, 4H), 2.57 ppm (t, 2H); MS (EI) C₂₂H₂₃N₅O₃HCl: 434.3 (MH⁺).

20

5


Example 13

N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)prolinamide (576)



- 0 [0201] To a solution of **249(a)** (5HCl) (0.2 g, 0.37 mmol) in DMA (5 mL) was added a solution of 1-(*tert*-butoxycarbonyl)pyrrolidine-2-carboxylic acid, (*d,l*-*boc*-proline) (0.1 g, 0.46 mmol), Hunigs base (0.5 mL, 2.5 mmol), HATU (0.2 g, 0.52 mmol) and the solution was stirred at RT for 14 hours. The resulting solution was loaded on the silica gel and was purified by silica gel column chromatography (10-100% gradient of ethyl acetate/hexanes) to yield **Z** (160 mg) in 79% yield as yellow solid. LCMS: m/z 545 ($M+H$)⁺.

d) *N*-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide (**Compound 585**)




- 10 [0202] A flask containing a solution of **Z** (160 mg, 0.29 mmol) in 4M HCl in 1,4-dioxane (5 mL) and MeOH (5 mL) was stirred at 50 C for 1 hour. Concentration of the solvent gave a yellow solid that was purified by reverse phase HPLC using an ammonium acetate buffer to yield 105 mg (68%) of **576** as a yellow solid.

- 5 ¹H NMR (400 MHz, *d*₄-MeOD): 8.36 (m, 1H), 8.14 (m, 2H), 7.78 (m, 2H), 7.62 (m, 2H), 7.22 (m, 1H), 6.98 (m, 2H), 4.15 (m, 1H), 3.83 (m, 4H), 3.21 (m, 2H), 3.13 (m, 4H), 2.41 (m, 1H), 2.06-1.91 (m, 3H); LCMS: for C₂₅H₂₈N₆O₂: 445 (M + H)⁺.

Example 14

- 10 **2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide (Compound 208)**



- [0203] A flask was charged with **249(a)** (140 mg, 0.3 mmol), *N*-(*tert*-butoxycarbonyl)-alanine (57 mg, 0.3 mmol, purchased from Chem-Impex International), HATU (140 mg, 0.37 mmol), diisopropylethylamine (0.6 mL, 3.0 mmol) and DMA (5 mL). The reaction mixture was stirred at RT for 12 hours. The crude mixture was concentrated on a rotary evaporator and the residue was dissolved in 10 mL of MeOH and 5 mL of 4N HCl in dioxane. The reaction mixture was stirred at 70 °C for 1 hour. The crude mixture was concentrated on a rotary evaporator and the product was purified by HPLC with NH₄OAc/ACN as eluent. The resulting solution was concentrated on a rotary evaporator and the final product, **208**, was dried by lyophilization.


- 5 ¹H-NMR (400MHz, *d*₆-DMSO): 9.387 ppm (s, 1H), 8.443 ppm (d, 1H), 8.127 ppm (d, 2H), 7.825 ppm (d, 2H), 7.676 ppm (d, 2H), 7.287 ppm (d, 1H), 6.939 ppm (d, 2H), 3.747 ppm (m, 4H), 3.457 ppm (q, 1H), 3.050 ppm (m, 4H), 1.896 ppm (s, 3H (AcOH)) 1.243 ppm (d, 3H); MS (EI) C₂₃H₂₆N₆O₂: 419.1 (MH⁺).

5

Example 15

**N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide
(Compound 341)**

a) 4-(2-methoxy-4-nitrophenyl)morpholine (AA)




10

[0204] A pressure bottle was charged with 1-chloro-2-methoxy-4-nitrobenzene (10.0 g mg, 53.3 mmol, purchased from TCI America) and morpholine (15 mL, 172.0 mmol). The reaction mixture was stirred at 120 °C for 15 hours and it was allowed to cool to room temperature by itself. The resulting solid was suspended in 20 mL of ethyl acetate, filtered, and washed with 20 mL of *tert*-butyl methyl ether. 8.8 g of yellow solid as the desired product AA was collected (69% yield). ¹H-NMR (400MHz, d₆-DMSO): 7.83 (dd, 1H), 7.67 (d, 1H), 6.98 (d, 1H), 3.88 (s, 3H), 3.71 (m, 4H), 3.16 (m, 4H). MS (EI) C₁₁H₁₄N₂O₄: 239 (M+H)⁺.

15

b) 3-methoxy-4-morpholinoaniline




20

[0205] To a solution of AA (8.8 g, 37.0 mmol) in ethyl acetate (30 mL) and methanol (10 mL) in a Parr bottle was added 1 g 10% palladium on carbon. The reaction mixture was hydrogenated at 40 PSI H₂ for 1 hour, filtered and concentrated. 8.0 g of a pink solid as the product BB was obtained as a crude product and used without further purification. MS (EI) C₁₁H₁₆N₂O₂: 209 (M+H)⁺.

25

5 c) *N*-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide




[0206] A flask was charged with BB (51 mg, 0.24 mmol), *N*-(4-(2-chloropyrimidin-4-yl)phenyl)acetamide (50 mg, 0.2 mmol) and *n*BuOH (2 mL). The flask was immersed in a 180 °C oil bath for 30 minutes, and then cooled to ambient temperature. The residue was suspended in 5 mL of ethyl acetate, stirred for 1 hour, filtered, and washed with 10 mL of ethyl acetate. 50 mg of an off-white powder was obtained as the title compound (341) (60% yield).

15 ¹H-NMR (400MHz, d₆-DMSO): 10.33 (s, 1H), 9.50 (br, 1H), 8.54 (d, 1H), 8.15 (d, 2H), 7.95 (br, 1H), 7.77 (d, 2H), 7.42 (m, 2H), 3.94 (s, 3H), 3.77 (br, 4H), 3.40 (br, 2H), 2.15 (s, 2H). MS (EI) C₂₃H₂₅N₅O₃: 420 (M+H)⁺.

Example 16

20 *N*-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)methanesulfonamide (Compound 326)



[0207] 82 (500 mg, 0.94 mmol) was dissolved in 4 mL of pyridine. Methanesulfonyl chloride (730 μL, 9.4 mmol) was added dropwise to the vigorously stirred pyridine solution. 25 The addition of the sulfonyl chloride was exothermic and caused a significant increase in the temperature of the reaction. The reaction was maintained at 80 °C for several hours. After


5 cooling, the solvent was removed under vacuum and the residue was purified by reverse phase HPLC to afford 200 mg (52% yield) of the title compound (326).

¹H-NMR (400MHz, d₆-DMSO): 10.16 ppm (s, 1H), 9.41 ppm (s, 1H), 8.45 ppm (d, 1H), 8.13 ppm (d, 2H), 7.67 ppm (d, 2H), 7.33 ppm (d, 2H), 7.28 ppm (d, 1H), 6.94 ppm (d, 2H), 3.74 ppm (br s, 4H), 3.09 ppm (s, 3H), 3.05 ppm (br s, 4H); MS (EI) C₂₁H₂₃N₅O₃S: 426 (MH⁺).

10

Example 17


Methyl (4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)carbamate (Compound 248)



15 [0208] To a solution of 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine 249(a) (100 mg, 0.29 mmol) and DIEA (0.435 mmol, 75 μ l) in THF (50 mL) was added methyl chloroformate (0.348 mmol, 27 μ l) and the solution was stirred at room temperature for 2 hours. The solution mixture was concentrated, redissolved with MeOH and purified using reverse phase HPLC. The product obtained from the reverse phase HPLC was free
20 base 248, converted to HCl salt using 3 N HCl and lyophilized to yield the product 248 (60mg, 47% yield) as a yellow solid.

¹H-NMR (400MHz, d₆-DMSO): 10.063 ppm (s, 1H), 9.976 ppm (s, 1H), 8.521 ppm (d, 1H), 8.153 ppm (d, 2H), 7.878 ppm (d, 2H), 7.661 ppm (d, 2H), 7.554 ppm (bs, 2H), 7.432 ppm (d, 1H), 3.983 ppm (bs, 4H), 3.707 ppm (s, 3H), 4.435 ppm (bs, 4H); MS (EI)
25 C₂₂H₂₃N₅O₃HCl: 475.4 (MH⁺).


5

Example 18**(S)-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide (Compound 363)**

- 10 [0209] To a solution of (S)-3-hydroxybutyrate (0.180 g, 1.73 mmol), HATU (0.602 g, 1.58 mmol), DIEA (1.0 mL, 5.4 mmol) in DMF (3.0 mL) was added a solution of 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine (0.500 g, 1.44 mmol) in DMF (1.0 mL). The reaction mixture was stirred at rt for 2 hours, at which time it was
- 15 quenched with saturated NaHCO₃ (10 mL, aq.), extracted into DCM (3X), and washed with brine (1X). The organic layers were dried with sodium sulfate and concentrated. The product was purified by reverse phase HPLC to afford (S)-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide (0.136 g, 22% yield) as a light brown solid. (363) ¹H-NMR (400MHz, DMSO-*d*₆): 10.14 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.94 (d, 2H), 4.79 (d, 1H),
- 20 4.11 (m, 1H), 3.74 (m, 4H), 3.05 (m, 4H), 2.47 (dd, 1H), 2.35 (dd, 1H), 1.15 (d, 3H); MS (EI) m/z for C₂₄H₂₈N₅O₃: 434.3 (MH⁺).

Example 19**2-Hydroxy-2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide (Compound 366)**

25




- [0210] To a solution of 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine (1.0 g, 2.8 mol) 249(a) and DIPEA (0.5 mL, 1 eq.) in anhydrous DMA (5 mL) was added dropwise 2-acetoxy-2-methylpropionyl chloride (3 mol, 1.05 eq., 0.44 mL) at 0 °C. The

5 mixture was stirred for 20 min at room temperature. The solution was diluted with water and EtOAc. The organic layer was concentrated *in vacuo*. The residue **366(a)** was suspended in MeOH (10 mL) and a solution of LiOH-H₂O (8.3 mmol, 3 eq. 0.35 g) in water (3 mL) was added. The reaction was complete within 20 min and then was neutralized. The organic solvent was removed *in vacuo*. The residue was purified to afford 2-hydroxy-2-methyl-*N*-(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)propanamide (**366**) (1.0 g, 85% yield)
 0 as a pale yellow solid. ¹H-NMR (400MHz, DMSO-*d*₆): 9.86 (s, 1H), 9.41 (s, 1H), 8.47 (d, 1H), 8.12 (d, 2H), 7.94 (d, 2H), 7.68 (d, 2H), 7.30 pm (d, 1H), 6.94 (d, 2H), 5.82 (s, 1H), 3.75 (m, 4H), 3.08 (m, 4H), 1.38 (s, 6H); MS (EI) *m/z* for C₂₂H₂₃N₅O₃HCl: 434.2 (MH⁺).

15 Example 20


(*R*)-3-hydroxy-*N*-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide (Compound 364)



20 [0211] To a solution of (*R*)-3-hydroxybutyrate (0.180 g, 1.73 mmol), HATU (0.602 g, 1.58 mmol), DIEA (1.0 mL, 5.4 mmol) in DMF (3.0 mL) was added and a solution of 4-(4-aminophenyl)-*N*-(4-morpholinophenyl)pyrimidin-2-amine (**249(a)**) (0.500 g, 1.44 mmol) in DMF (1.0 mL). The reaction mixture was stirred at room temperature for 2 hours, at which time it was quenched with saturated NaHCO₃ (10 mL, aq.), extracted into DCM
 25 (3X), washed with brine (1X), and the organic layers were dried with sodium sulfate. The solution was concentrated and the product was purified by reverse phase HPLC to afford (*R*)-3-hydroxy-*N*-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide (**364**) (0.129 g, 21% yield) as a light brown solid. ¹H-NMR (400MHz, DMSO)-*d*₆: 10.14 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.94 (d, 2H), 4.79 (d, 1H), 4.11 (m, 1H), 3.74 (m, 4H), 3.05 (m, 4H), 2.47 (dd, 1H), 2.35 (dd, 1H),
 30 1.15 (d, 3H); MS (EI) *m/z* for C₂₄H₂₈N₅O₃: 434.3 (MH⁺).

5

Example 21

(R)-2-amino-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide (Compound 365)

[0212] To a solution of 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine
 10 **249(a)** (521mg, 1.5 mmol), N-CBZ-D-Serine (359mg, 1.5mmol), and DIEA (0.653mL,
 3.75mol) in DMA (4mL) was added HATU (855mg, 2.25mmol) and the solution was stirred
 at room temperature for 0.5 hour. Excess H₂O was added to the reaction mixture. The
 precipitate was collected and were redissolved in CH₂Cl₂, washed with NaHCO₃ (aq) (2X),
 brine, dried over Na₂SO₄, and concentrated *in vacuo*. The residue was purified by silica
 15 column chromatography (1% MeOH/DCM as eluent) to afford (R)-benzyl 3-hydroxy-
 1-(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenylamino)-1-oxopropan-
 2-ylcarbamate **365(a)** (668 mg, 78% yield).


[0213] To a stirred solution of (R)-benzyl 3-hydroxy-1-(4-(2-(4-
 morpholinophenylamino)-pyrimidin-4-yl)phenylamino)-1-oxopropan-2-ylcarbamate from the
 10 step above in MeOH (10 mL) was added Pd(OH)₂ (134 mg) and ammonium formate (369
 mg, 5.85). The mixture was heated at 60 °C for 2 hours, cooled down to room temperature,
 and filtered on Celite by eluting with MeOH. The filtrate was concentrated *in vacuo* and the
 residue was purified by preparatory HPLC (TFA). The TFA salt was removed by using basic
 resin to afford (R)-2-amino-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-
 15 yl)phenyl)propanamide **365** (346 mg, 68%). ¹H-NMR (400MHz, DMSO-*d*₆): 9.39 (s, 1H),
 8.44 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.94pm (d, 2H), 4.94
 (m, 1H), 3.75 (m, 4H), 3.60 (m, 2H), 3.46 (m, 1H), 3.05 (m, 4H); MS (EI) m/z for
 C₂₃H₂₆N₆O₃: 435.4 (MH⁺).

0

5

Example 22

N-{4-[2-({3-[(4-ethylpiperazin-1-yl)methyl]phenyl}amino)pyrimidin-4-yl]phenyl}-acetamide (Compound 122)



10

[0214] Intermediate A (0.5g) was dissolved in THF (5 ml), 20% aqueous H_2SO_4 solution (5ml) was then added to the solution. The mixture was stirred at 50°C for 2 hours and monitored by LC/MS (MH^+ , 333). The solution was then neutralized with 2N NaOH solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 , and concentrated to afford 0.38g of the aldehyde B. (90% yield)

5

[0215] A flask was charged with aldehyde B (0.1g, 0.3 mmol), dichloromethane (10 ml), sodium triacetoxyborohydride (0.32g, 1.5mmol), and 1-ethylpiperazine (0.19 ml, 1.5mmol). The reaction mixture was stirred at room temperature overnight and checked with LC/MS. The product 122 was isolated by removal of the solvent with a rotary evaporator and then purified with a preparative HPLC.


10

^1H NMR (400 MHz, d_6 -DMSO): 10.23 (s, 1H), 9.6 (s, 1H), 8.5 (d, 1H), 8.14 (d, 2H), 7.9 (s, 1H), 7.76 (d, 2H), 7.65 (d, 1H), 7.36 (d, 1H), 7.24 (t, 1H), 6.89 (d, 1H), 3.43 (s, 2H), 2.4 (br, 6H), 2.3 (q, 2H), 2.1 (s, 3H), 0.96 (t, 3H). MS (EI) for $\text{C}_{25}\text{H}_{30}\text{N}_6\text{O}$: 431 (MH^+).

5

Example 23

2-(3-(1H-imidazol-1-yl)propylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide (Compound 143)



10 [0216] A flask was charged with aniline A₁ (100 mg, 0.29 mmol), and THF (1.0 mL). Chloroacetylchloride (23 μ L, 0.29 mmol) was added and the mixture was stirred at ambient temperature for 1 hr, after which time it was concentrated. The product, B₁, was isolated by removal of the solvent with a rotary evaporator and used without further purification.

15 [0217] A flask was charged with alkyl chloride B₁ (20 mg, 0.047 mmol), Na₂CO₃ (30 mg, 0.28 mmol), 1-(3-Aminopropyl)imidazole (5.6 μ L, 0.047 mmol), and DMF (1.0 mL). The mixture was stirred at 150 °C for 1 hr, after which time it was concentrated. The product 143 was purified by reverse phase HPLC to afford 9.7 mg (40% yield from B₁) as a white solid.


¹H-NMR (400MHz, d₆-DMSO): 8.35 (d, 1H), 8.13 (d, 2H), 7.78-7.63 (m, 3H), 7.61 (d, 2H), 7.22 (d, 1H), 7.17 (s, 1H), 7.05-6.95 (m, 2H), 4.62 (s, br, 1H), 4.16 (t, 2H), 3.87-3.77 (m, 4H), 3.49 (s, 1H), 3.34 (s, 1H), 3.15-3.07 (m, 4H), 2.67 (t, 2H), 2.11-2.01 (m, 2H), 1.95 (s, 2H). MS (EI) C₂₈H₃₂N₈O₂: 513.1 (MH⁺).

20

5

Example 24

N-chloro-N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(1H-tetrazol-1-yl)acetamide (Compound 554)



10

[0218] 4-(4-aminophenyl)-N-(3-methoxy-4-morpholinophenyl)pyrimidin-2-amine

hydrochloride: A flask was charged with tert-butyl 4-(2-chloropyrimidin-4-yl)phenylcarbamate A (12.2 g, 40.0 mmol), 3-Methoxy-4-morpholinoaniline (9.7 g, 40.76 mmol) and 50 mL n-butanol. The reaction mixture was stirred under an N₂ atmosphere at 100 °C for 12 hours, after which time, then, cooled to room temperature. 25 mL of 4N HCl in dioxane was added, the reaction mixture was stirred at 50 °C for 5 hours. After cooled to room temperature, it was filtered, washed with ethyl acetate, dried in the air to collect 16 g of yellow-green solid as the desired product. NMR (400 MHz, d₆-DMSO): 10.40 (s, 1H), 8.60 (d, 1H), 8.20 (s, 2H), 7.94 (s, 1H), 7.84 (d, 1H), 7.54 (d, 1H), 7.41 (d, 1H), 7.30 (m, 2H), 4.10 (m, 2H), 3.99 (s, 3H), 3.60 (br, 2H), 3.39 (m, 2H), 1.25 – 1.42 (m, 4H). MS (EI) for C₂₁H₂₃N₅O₂: 378 (MH⁺).

15

20


[0219] A flask was charged with 4-(4-aminophenyl)-N-(3-methoxy-4-morpholinophenyl)pyrimidin-2-amine hydrochloride B (471.0 mg, 0.84 mmol), 2-(1H-tetrazol-1-yl)acetic acid (216.0 mg, 1.69 mmol), HATU (1276.0 mg, 3.38 mmol) and 2 mL of

25

5 DMA. The reaction mixture was stirred at room temperature for 24 hours, and then quenched with 50 mL of water, extracted with ethyl acetate (3X50 mL). The combined organics were washed with water and then brine (50 mL each), dried over anhydrous sodium sulfate, and then concentrated. The crude product was purified with a silica gel column (ethyl acetate to 10% methanol in ethyl acetate), 345.0 mg of the desired product **554** was obtained as
 10 yellowish powder. NMR (400 MHz, d₆-DMSO): 10.85 (s, 1H), 9.50 (s, 1H), 9.43 (s, 1H), 8.47 (s, 1H), 8.19 (d, 2H), 7.75 (d, 2H), 7.63 (s, 1H), 7.20 (m, 3H), 6.84 (d, 1H), 5.56 (s, 2H), 3.80 (s, 3H), 3.74 (m, 4H), 2.94 (m, 4H). MS (EI) for C₂₄H₂₅N₉O₃: 488 (MH⁺).


Example 25

15 **4-[4-(1,1-Dioxidoisothiazolidin-2-yl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (Compound 374)**



[0220] Aniline A (300 mg, 0.78 mmol) was dissolved in 4 mL of dry pyridine. 3-Chloropropanesulfonyl chloride (950 uL, 7.8 mmol) was added dropwise. The reaction
 20 mixture was heated to 80 °C and stirred overnight under a nitrogen atmosphere. The solvent was removed under vacuum and the residue was re-dissolved in 25 mL of ethyl acetate. The reaction mixture was washed one time each with 10 mL portions of water, 0.1 M HCl, and saturated aqueous NaCl. The organic layer was dried with MgSO₄ and concentrated under vacuum. The residue was taken up in DMF (4 mL) and triethylamine (1100 uL, 7.9 mmol).
 25 The reaction mixture was heated to 80 °C and stirred overnight. The product was purified by preparative HPLC to give 85 mg of **374**. ¹H NMR (400 MHz, d₆-DMSO): 9.78 (s, 1H), 8.49 (d, 1H), 8.20 (d, 2H), 7.78 (d, 2H), 7.39 (d, 1H), 7.33 (d, 2H), 7.25 (br s, 2H), 3.84 (br s, 4H), 3.73 (t, 2H), 3.60 (t, 2H), 2.54 (m, 2H), 2.45 (m, 2H) 2.01 (m, 2H); MS (EI) for C₂₃H₂₅N₅O₃S: 452 (MH⁺).

5


Example 26**N-(4-Morpholin-4-ylphenyl)-4-[4-(1H-tetrazol-1-yl)phenyl]pyrimidin-2-amine**

- 0 [0221] Aniline A (200 mg, 0.52 mmol), sodium azide (45 mg, 0.69 mmol), triethylorthoformate (280 μ L, 1.7 mmol) and acetic acid (480 μ L, 8.4 mmol) were combined in a 25 mL round bottom flask. The reaction mixture was stirred for 2 hours at 80 $^{\circ}$ C. The reaction mixture was allowed to cool to room temperature and then it was cooled further in an ice bath. A solution of 670 μ L of 6.0 M HCl in 1.25 mL of water was added to the reaction
- 15 mixture. After stirring in the ice bath for 5 minutes, another solution of sodium nitrite (50 mg, 0.72 mmol) in water (200 μ L) was added slowly. The precipitate was filtered off and purified by reverse phase HPLC to give 24 mg of **375**. ^1H NMR (400 MHz, d_6 -DMSO): 10.19 (s, 1H), 9.51 (s, 1H), 8.53 (d, 1H), 8.40 (dd, 2H), 8.10 (d, 2H), 7.65 (d, 2H), 7.43 (d, 1H), 6.92 (d, 2H), 3.73 (m, 4H), 3.03 (m, 4H); MS (EI) for $\text{C}_{21}\text{H}_{20}\text{N}_8\text{O}$: 401 (MH^+).

5

Example 27

N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-fluoro-6-iodobenzamide (Compound 289)



0

[0222] A flask was charged with C1 (5.0 g, 20.2388 mmol) and (3-aminopropyl)-carbamic acid-t-butyl ester (6 mL, 30.3582 mmol). N-butanol (40 mL) were added to the flask and heated to 175 °C for an hour. Solvent was evaporated and reaction mixture was checked with LC/MS. The reaction mixture was cooled to room temperature and ethyl acetate was added. The precipitate, B, was filtered and used without further purification. LC/MS: m/z 386 ($M+H$)⁺.

5

[0223] A flask was charged with B. 4 N HCl in dioxane was added and stirred at room temperature for 3 hours. The reaction mixture was checked with LC/MS. The product, E, was isolated by removal of the solvent with a rotary evaporation and used without further purification. LC/MS; m/z 286 ($M+H$)⁺.

10

[0224] A flask was charged with E (254 mg, 0.8902 mmol), 2-fluoro-6-iodobenzoyl chloride (90 μ L, 0.6231 mmol), tetrahydrofuran (25 mL), and n-ethyldiisopropylamine (108 μ L, 0.6231 mmol). The reaction mixture was stirred at room temperature for 12 hours. The reaction mixture was monitored with LC/MS. The product, 289, was isolated by removal of the solvent with a rotary evaporator and purified with a TFA preparative HPLC (10:90, 11 min run).

15

¹H-NMR (400MHz, d₆-DMSO): 10.16 ppm (s, 1H), 8.64 ppm (t, 1H), 8.30 ppm (d, 1H), 8.06 ppm (d, 2H), 7.70 ppm (m, 3H), 7.30 ppm (m, 1H), 7.20 ppm (m, 1H), 7.13 ppm (m, 1H), 7.07 ppm (m, 1H), 3.34 ppm (m, 4H), 2.08 ppm (s, 3H), 1.83 ppm (m, 2H); MS (EI)


30

C₂₂H₂₁FIN₅O₂: 533.9 (MH)⁺.

5

Example 28

N-(4-(2-((3-((2,6-dimethylphenyl)methyl)amino)phenyl)amino)pyrimidin-4-yl)phenyl)acetamide (Compound 51)




10 [0225] A flask was charged with C1 (5.0 g, 20.2388 mmol) and tert-butyl 3-aminophenylcarbamate (4.6 g, 22.2627 mmol). N-butanol (40 mL) were added to the flask and heated to 175 °C for 4 hours. Solvent was evaporated and reaction mixture was checked with LC/MS. The reaction mixture was cooled to room temperature and ethyl acetate was added. The precipitate, D, was filtered and used without further purification. LC/MS: m/z 320 (M+H)⁺.

15 [0226] A flask was charged with D (463 mg, 1.4514 mmol), dichloromethane/tetrahydrofuran (2;1, 15 mL), sodium triacetoxyborohydride (615 mg, 2.9028 mmol), and 2,6-dimethylbenzaldehyde (196 µL, 1.4514 mmol). The reaction mixture was stirred at room temperature for 12 hours and monitored with LC/MS. The product, 51, was isolated by removal of the solvent with a rotary evaporator and purified with a TFA preparative HPLC (10:90, 11 min run). ¹H-NMR (400MHz, d₆-DMSO): 10.20 ppm (s, 1H), 9.36 ppm (s, 1H), 8.47 ppm (d, 1H), 8.16 ppm (d, 2H), 7.72 ppm (d, 2H), 7.35 ppm (s, 1H), 7.31 ppm (d, 1H), 7.12 ppm (m, 1H), 7.07 ppm (m, 2H), 6.99 ppm (m, 3H), 6.38 ppm (d, 1H), 5.46 ppm (t, 1H), 4.14 ppm (d, 2H), 2.36 ppm (s, 6H), 2.08 ppm (s, 3H); MS (EI) C₂₇H₂₇N₅O: 438.1 (MH⁺).

5

Example 29

3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(dimethylamino)ethyl]benzamide (Compound 9)



0 [0227] A flask was charged with 2,4-dichloropyrimidine (22.7 g, 152.38 mmol), 4-acetoamidophenylboronic acid (30.0 g, 167.62 mmol), dichloro[1,1'-bis(diphenylphosphino)-ferrocenepalladium (16.726 g, 22.86 mmol, 15 mol %), and triethylamine (53 mL, 380.95 mmol). Ethyleneglycoldimethylether (500 mL) and H₂O (20 mL) was added to the flask. The reaction mixture was stirred at 80 °C for 4 hours. The product, Intermediate A, was isolated

5 by removal of the solvent with a rotary evaporator and purified using glass column chromatography and eluted with ethyl acetate to afford 30.5 g (123.14 mmol, 81% yield) of intermediate A as a yellow solid.

[0228] A seal tube was charged with intermediate A (400 mg, 1.62 mmol) and 3-aminobenzoic acid (222 mg, 1.62 mmol). N-butanol (15 mL) was added to the seal tube and

10 stirred at 180°C. The reaction was done in 1h according to LCMS to afford intermediate B as a yellow solid. Intermediate B was placed on a rotary evaporator to remove excess n-butanol. Intermediate B was carried on to the next step without further purification.

[0229] A flask was charged with intermediate B (282 mg, 0.81 mmol), HATU (464 mg, 1.22 mmol), DMF (15 mL) and DIEA (212 µL, 1.22 mmol). The reaction mixture was


15 stirred at rt and completed in 30 min. to afford the final product (9). The final product was

5 purified using Preparative HPLC and ammonium acetate buffer and lyophilized to afford the product as ACE salt (170 mg, 0.41 mmol). ¹H-NMR (400MHz, d₆-CD₃OD): 8.523 ppm (t, 1H), 8.45 ppm (d, 1H), 8.176 ppm (m, 2H), 7.828 ppm (m, 1H), 7.2 ppm (d, 2H), 7.475-7.404 ppm (m, 2H), 7.326 ppm (d, 1H), 3.738 ppm (t, 2H), 3.244 ppm (t, 2H), 2.877 ppm (s, 6H), 2.162 ppm (s, 3H), 1.955 (s, 3H, ACE). MS (EI) C₂₃H₂₆N₆O₂: 419.1 (MH⁺).

10

Example 30

N-[5-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-2-morpholin-4-ylphenyl]-2,6-dichlorobenzamide (Compound 62)



15 **[0230]** A seal tube was charged with intermediate A (500 mg, 2.02 mmol) and 4-morpholinobenzene-1,3-diamine (400 mg, 2.02 mmol, Zerenex Limited). N-butanol (15 mL) was added to the seal tube and stirred at 180°C. The reaction was done in 1h according to LCMS to afford intermediate C as a yellow solid. Intermediate C was placed on a rotary evaporator to remove excess n-butanol. Intermediate C was carried on to the next step

20 without further purification.


[0231] A flask was charged with intermediate C (816 mg, 2.02 mmol), THF (100 mL), DIEA (705 μL, 4.04mmol), and 2,6-dichlorobenzoyl chloride (290 μL, 2.02 mmol). The reaction mixture was stirred at rt over night to afford the final product 62. The final product

5 was purified using Preparative HPLC and TFA buffer, then was free-based and lyophilized (165 mg, 0.28 mmol, 14% Yield).

¹H NMR (400 MHz, DMSO): 10.194 (s, 1H), 9.8 (s, 1H), 9.607 (s, 1H), 8.585 (s, 1H), 8.484 (d, 1H), 8.235 (d, 2H), 7.711 (d, 2H), 7.592 (d, 2H), 7.496 (m, 2H), 7.356 (d, 1H), 7.183 (d, 1H), 3.74 (t, 4H), 2.89 (t, 4H), 2.07 (s, 3H). MS (EI) for C₂₉H₂₆Cl₂N₆O₃: 579.1 (MH⁺).

Example 31

N-{3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-5-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}-2,6-dichlorobenzamide (Compound 66)



[0232] A seal tube was charged with intermediate A (300 mg, 1.21 mmol) and 3,5-diaminobenzoic acid (204 mg, 1.34 mmol). N-butanol (15 mL) was added to the seal tube and stirred at 180°C. The reaction was done in 1h according to LCMS to afford intermediate D as a yellow solid. Intermediate D was placed on a rotary evaporator to remove excess n-butanol. Intermediate D was carried on to the next step without further purification.

[0233] A flask was charged with intermediate D (439 mg, 1.21 mmol), THF (30 mL), DMF (5 mL), DIEA (632 µL, 3.63 mmol), and 2,6-dichlorobenzoyl chloride (174 µL, 1.21 mmol). The reaction mixture was stirred at rt over night. The reaction mixture was quenched with 2 M NaOH (100 mL) and extracted with ethyl acetate (3x) and the organic layer was


discarded. The aqueous NaOH layer was neutralized with conc. HCl. The solid formed was collected via filtration and washed with excess water to afford intermediate E (274 mg, 0.51 mmol, 62% yield) as a yellow solid. Intermediate E was carried on to the next step without further purification.

[0234] A flask was charged with intermediate E (274 mg, 0.51 mmol), HATU (291 mg, 0.765 mmol), DMF (25 mL), ethylpiperazine (78 μ L, 0.61 mmol) and DIEA (133 μ L, 0.765 mmol). The reaction was stirred at rt and completed in 15 min. The final product **66** was purified using Preparative HPLC and TFA buffer, free-based and lyophilized to afford the product (166 mg, 52% yield).

¹H NMR (400 MHz, DMSO): 10.896 (s, 1H), 10.33 (s, 1H), 9.881 (s, 1H), 8.533 (d, 1H), 8.374 (s, 1H), 8.202 (d, 2H), 7.776 (d, 2H), 7.636-7.6 (m, 3H), 7.529 (m, 1H), 7.419 (d, 1H), 7.296 (s, 1H), 3.628 (br s, 2H), 3.415 (br s, 2H), 2.427-2.314 (m, 6H), 2.091 (s, 3H), 0.996 (t, 3H). MS (EI) for C₃₂H₃₁Cl₂N₇O₃: 634.1(MH⁺).

Example 32

***N*-methyl-*N*-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 118):**




[0235] A flask was charged with intermediate A (250 mg, 1.01 mmol), DMF (10 mL), NaH (30.0 g, 167.62 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (60 mg, 1.5 mmol), and methyl iodide (94 μ L, 1.5 mmol). The reaction mixture was stirred at rt and completed in 30 min. The reaction mixture was quenched with H₂O and extracted with ethyl acetate (3X) and washed with 10% LiCl solution (1X), brine (1X), dried over sodium sulfate, and filtered. The organic layer was removed with a rotary evaporator to afford intermediate F (200 mg, 0.766 mmol) as a yellow gelatin. Intermediate F was carried on to the next step without further purification.

5 **[0236]** A seal tube was charged with intermediate **F** (200 mg, 0.766 mmol), anhydrous DMA (15 mL), cesium carbonate (374 mg, 1.15 mmol), racemic-2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl (70 mg, 0.115 mmol), and tris(dibenzylideneacetone)dipalladium(0). The reaction was flushed with N₂ gas for five minutes and the seal tube was sealed and stirred at 80°C over night. The reaction was filtered
 10 and washed with ethyl acetate and the solid was discarded. The organic solvent was removed using the rotary evaporator. The final product **66** was purified using Preparative HPLC and TFA buffer, free-based and lyophilized to afford the product (95 mg, 0.235 mmol, 28% Yield).

¹H-NMR (400MHz, d₆-DMSO): 9.464 ppm (s, 1H), 8.511 ppm (d, 1H), 8.209 ppm (d, 2H),
 15 7.67 ppm (m, 2H), 7.516 ppm (d, 2H), 7.366 ppm (d, 1H), 6.926 ppm (m, 2H), 3.743 ppm (t, 4H), 3.22 ppm (s, 3H), 3.048 ppm (t, 4H). MS (EI) C₂₃H₂₅N₅O₂: 404.3 (MH⁺).

Example 33

N-(4-(2-(3-(3-morpholinopropoxy)phenylamino)pyrimidin-4-yl)phenyl)acetamide
 20 **(Compound 160)**




[0237] A seal tube was charged with intermediate **A** (500 mg, 2.02 mmol) and 3-benzyloxyaniline (404 mg, 2.02 mmol). N-butanol (15 mL) was added to the seal tube and

5 stirred at 180°C. The reaction was done in 1h according to LCMS to afford intermediate **G** as a yellow solid. Intermediate **G** was placed on a rotary evaporator to remove excess n-butanol. Intermediate **G** was carried on to the next step without further purification. A flask was charged with intermediate **G** and HBr/Acetic acid (33%, 10 mL) and stirred at rt over night. The reaction was done and the solid was collected via filtration and washed with ether
10 to afford intermediate **H** as a yellow and HBr salt solid (800 mg, 1.66 mmol, 82% yield).

[0238] A flask was charged with intermediate **H** (250 mg, 0.52 mmol), DMF (15mL), Cs₂CO₃ (847mg, 2.6 mmol) and 4-(3-chloropropyl)morpholine HCl salt (135 mg, 0.676 mmol, purchased from Apin Chemicals, Ltd.) and stirred at 80°C over night. The reaction mixture had approximately 85% desired product and 15% bis-alkylated by-product. The
15 solid was filtered and washed with ethyl acetate and discarded. The filtrate was concentrated using the rotary evaporator. The final product was purified using Preparative HPLC and TFA buffer, free-based, converted to HCl salt and lyophilized to afford the product (115mg, 0.237 mmol, 46% Yield).

¹ H NMR (400 MHz, DMSO): 11.058 (s, 1H), 10.403 (s, 1H), 9.761 (s, 1H), 8.532 (d, 1H),
20 8.158 (d, 2H), 7.81 (d, 2H), 7.677 (s, 1H), 7.4-7.345 (m, 2H), 7.231 (t, 1H), 6.569 (m, 1H), 4.081 (t, 2H), 3.962 (m, 2H), 3.82 (t, 2H), 3.46 (m, 2H), 3.267 (m, 2H), 3.123 (m, 2H), 2.254 (m, 2H), 2.104 (s, 3H). MS (EI) for C₂₅H₂₉N₅O₃: 448.3 (MH⁺).

5

Example 34***N*-(4-{2-[(2-methyl-4-piperazin-1-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide
(Compound 35):**

0 [0239] A flask was charged with 5-fluoro-2-nitrotoluene (1 mL, 8.2 mmol), DMF (15 mL), Bocpiperazine (1.68 g, 9.02 mmol), and K_2CO_3 (2.27 g, 16.4 mmol). The reaction mixture was stirred at 50 $^\circ\text{C}$ for about 25 h. The reaction was quenched with H_2O and the solid precipitated out of the solution and collected via filtration and washed with excess H_2O to obtain intermediate I (1.765g, 5.4 mmol). Intermediate I was carried on to the next step

5 without further purification. A flask was charged with intermediate I (290 mg, 0.9 mmol), ethanol (18 mL), ammonium formate (340 mg, 5.4 mmol) and Pt/S (10.2 mg, 0.04 mmol). The reaction mixture was stirred at 70 $^\circ\text{C}$ for 3 h and 78 $^\circ\text{C}$ for 4 h. The reaction mixture was filtered through celite and washed with ethanol. The filtrate was removed using the rotary evaporator and then treated with ethyl acetate and washed with H_2O , dried over sodium

20 sulfate, and filtered. The ethyl acetate layer was concentrated using rotary evaporator to afford intermediate J.

[0240] A seal tube was charged with intermediate A (200 mg, 0.81 mmol), and intermediate J (235 mg, 0.81 mmol). N-butanol (15 mL) was added to the seal tube and stirred at 180 $^\circ\text{C}$. The reaction was done in 1h and concentrated to remove excess n-butanol


25 and then treated with 4N HCl/dioxane. The reaction mixture was stirred at rt for 1h to afford the final product 35. The final product was purified using Preparative HPLC and ammonium acetate buffer, then free-based and lyophilized (90 mg, 0.22 mmol, 27% Yield).

$^1\text{H-NMR}$ (400MHz, $\text{d}_6\text{-DMSO}$): 10.31 ppm (s, 1H), 8.569 ppm (s, 1H), 8.323 ppm (d, 1H), 8.022 ppm (d, 2H), 7.715 ppm (d, 2H), 7.271 ppm (d, 1H), 7.186 ppm (d, 1H), 6.8 ppm (m,


5 2H), 3.023 ppm (t, 4H), 2.844 ppm (t, 4H), 2.175 ppm (s, 3H), 2.077 ppm (s, 3H), 1.605 ppm (s, 2H); MS (EI) C₂₃H₂₆N₆O: 403.1 (MH⁺).

Example 35


N-[4-({2-[(4-morpholin-4-ylphenyl)amino]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl}amino)-phenyl]acetamide (Compound 306)




[0241] A flask was charged with methylsulfide (2.1g, 11.6mmol) and THF (50mL). To this, m-CPBA (7.9g, 46mmol) was added and the mixture was stirred at ambient temperature for 20 hours. Volatiles were removed under *vacuo*. The crude mixture was partitioned between EtOAc and DI H₂O. The aqueous layer was extracted with EtOAc (3x15mL). The combined organics were washed with 1N NaHCO₃ (x2), DI H₂O (x2), brine, (x1), dried over sodium sulfate, filtered and concentrated under *vacuo*. The product (1.8g, 75%) was used without further purification. LCMS: m/z 214(M+H)⁺.



[0242] A pressure tube was charged with methylsulfone (1.15g, 5.4mmol) and aniline (2.8g, 16.2mmol). The tube was sealed and the mixture heated at 140°C for 30 minutes. The mixture was cooled. Methanol was added and the resulting solid collected via filtration then washed with methanol. The product (270mg, 8.7%) was used without further purification. LCMS: 312 (M+H)⁺.



[0243] A flask was charged with pyrrolopyrimidinone (250mg, 0.8mmol) and toluene (5mL). Phosphorous oxychloride (218 μ L, 2.41mmol) and DIPEA (165 μ L, 0.96mmol) were added and the mixture stirred at 110°C for 6 hours. The volatiles were removed under *vacuo* and the product used without further purification. LCMS: 330 (M+H)⁺.




[0244] A flask was charged with pyrrolopyrimidine (100mg, 0.3mmol) and isopropanol (1mL). Aniline (55mg, .036mmol) and two drops of conc. HCl were added and the mixture heated to reflux for 6 hours. Volatiles were removed under *vacuo*. The product was purified by preparative HPLC to afford the title compound (306) (12.8mg, 9.6%).

¹H NMR (400MHz, d6-DMSO): 11.13 (s, 1H), 9.95 (s, 1H), 9.04 (s, 1H), 8.56 (s, 1H), 7.86 (d, 2H), 7.66 (d, 2H), 7.54 (d, 2H), 6.88-6.82 (m, 3H), 6.65-6.61 (m, 1H), 3.78-3.71 (m, 4H), 3.05-2.99 (m, 4H), 2.04 (s, 3H). MS (EI) for C₂₄H₂₅N₇O₂: 444 (MH⁺).


5

Example 36

(N-(4-{2-[(3-{[(2,6-dichlorophenyl)sulfonyl]amino}phenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide) (Compound 26)




- 10 [0245] To a mixture of 2,4-dichloro-5-methylpyrimidine (4.17g, 25.6mmol) and 4-acetamidophenylboronic acid (5.0g, 27.9mmol) in DME (40ml) was added Et₃N (8.92ml, 64.0mmol), H₂O (4ml), and dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (2.81g, 3.44mmol, 13%). The mixture was allowed to stir at reflux for 5hrs. After the mixture was cooled down to rt, the crude mixture was directly filtered on silica gel and eluted with EtOAc.
- 15 The filtrate was concentrated *in vacuo*. Further purification was conducted by flash chromatography to afford Intermediate 1 (5.94g, 89%) as a white solid. LCMS: m/z 262 (M+H)⁺.



20

- [0246] To a stirred solution of chloropyrimidine (1.05g, 4.0mmol) in 1-butanol (10ml) was added N-Boc-amino-3-aniline (920mg, 4.4mmol) and the mixture was heated in the sealed tube at 180°C for 1.5hr. The mixture was cooled down to rt and acidified with 1N HCl
- 25 (20ml). The aqueous layer was washed with EtOAc (50ml). The separated aqueous layer was basified with 2N NaOH to pH 8-9 and extracted with EtOAc (50ml*3). The combined organic layer was dried over Na₂SO₄, concentrated *in vacuo*, and purified by flash

- 5 chromatography to afford product **Intermediate K** (943mg, 71% as a light yellow solid. LCMS: m/z 334 ($M+H$)⁺.




- 0 [0247] To a stirred suspension of aniline (250mg, 0.75mmol) in THF (5ml) was added DIPEA (157ml, 0.90mmol) and 2,6-dichlorobenzenesulfonyl chloride (203mg, 0.83mmol) and the mixture containing intermediate K was stirred at reflux for 2hrs. After cooling down to rt, the mixture was diluted with EtOAc, washed with H₂O, brine, and dried over Na₂SO₄. After concentrated *in vacuo*, the residue was purified by flash chromatography to give
5 product **26** (299mg, 73%) as a light pink solid.

¹H-NMR (400MHz, d₆-DMSO): 10.71 (s, 1H), 10.16 (s, 1H), 9.54 (s, 1H), 8.34 (s, 1H), 7.75-7.69 (m, 5H), 7.60 (dd, 2H), 7.51 (dd, 1H), 7.31 (dd, 1H), 7.09 (t, 1H), 6.66 (dd, 1H), 2.25 (s, 3H), 2.08 (s, 3H); MS (EI) C₂₅H₂₁Cl₂N₅O₃S: 542.2 ($M+H$)⁺.

10

Example 37


N-(4-{6-morpholin-4-yl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 47)



- 15 [0248] The mixture of 2,4,6-trichloropyrimidine (1.72ml, 15mmol) 4-acetamidophenylboronic acid (1.79g, 10mmol) in DME (20ml) was added Et₃N (3.5ml, 25.0mmol), H₂O (2ml), and dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (1.22g, 1.5mmol, 15%).


- 5 The mixture was allowed to stir at reflux for 2hrs. After the mixture was cooled down to rt, the crude mixture was directly filtered on silica gel and eluted with EtOAc. The filtrate was concentrated *in vacuo*. Further purification was conducted by flash chromatography to afford intermediate L (1.91g, 68%) as a white solid. LCMS: m/z 282 ($M+H$)⁺.

0



- [0249] To a stirred suspension of pyrimidine (282mg, 1.0mmol) in 1-butanol (5ml) was added morpholine (96ml, 1.10mmol) and DIPEA (209μl, 1.2mmol). The mixture was heated at 120°C for 1hr, cooled down to rt, and concentrated *in vacuo*. The residue was purified by flash chromatography to afford intermediate M (176mg, 53%) as well as isomer (108mg, 32%). LCMS: m/z 333 ($M+H$)⁺.

20




- [0250] The mixture of chloropyrimidine (176mg, 0.53mmol) and 4-morpholinoaniline (104mg, 0.58mmol) in 1-butanol (5ml) was heated in the sealed tube at 160°C for 3hrs. The reaction mixture was cooled down to rt and the crude mixture was directly subjected on silica gel to afford product 47 (122mg, 49%) as a pale pink solid. LCMS: m/z 475 ($M+H$)⁺.

25


- 5 ¹H-NMR (400MHz, d₆-DMSO): 10.13 (s, 1H), 8.87 (s, 1H), 8.07 (d, 2H), 7.70-7.64 (m, 4H), 6.90 (d, 2H), 6.71 (d, 1H), 3.74-3.68 (m, 12H), 3.03 (t, 4H), 2.08 (s, 3H); MS (EI) C₂₆H₃₀N₆O₃: 475 (MH⁺).

Example 38

- 0 **N-[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyridin-2-yl]-2,6-dichloro-benzamide (Compound 299)**



- [0251] To a mixture of 2,6-diaminopyridine A (9.2 mmol, 1.0 g), and diisopropylethylamine (6.9mmol, 1.2ml) in 20ml of THF, was added 2,6-dichlorobenzoylchloride B (4.6mmol, 0.67ml) dropwise. The mixture was stirred at room temperature for 1 hour and LCMS indicated it was done (M+H: 283). THF was removed and replaced with ethyl acetate. The reaction mixture was then extracted with water, brine, and dried over sodium sulfate. The product, C, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: 283 (M+H).



- 25 [0252] A seal tube was charged with intermediate A₁ (0.2g, 0.81 mmol), compound C from the previous step (0.56g, 2.0mmol), tris(dibenzylideneacetone)dipalladium(0) (0.15g,


5 0.16 mmol), racemic-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (0.12g, 0.2mmol), cesium carbonate (0.4g, 1.22mmol). Dimethylacetamide (10ml) was added and the mixture was purged with N₂ for 5 minutes. The tube was sealed and the reaction mixture was stirred at 80°C overnight. LCMS showed the reaction was done (M+H: 493). The reaction mixture was partitioned between ethyl acetate and water, the organic layer extracted with 10% LiCl solution, followed by brine, dried over Na₂SO₄, and then evaporated. The crude product 299 was then purified via prep HPLC.

¹H NMR (400 MHz, d₆-DMSO): 11.12 (s, 1H), 10.25 (s, 1H), 9.43 (s, 1H), 8.58 (d, 1H), 8.2-8.13 (m, 3H), 7.9 (t, 1H), 7.83 (d, 1H), 7.78 (d, 2H), 7.55 (d, 2H), 7.52-7.45 (m, 2H), 2.1 (s, 3H). MS (EI) for C₂₄H₁₈Cl₂N₆O₂ : 493 (MH⁺).

5

Example 39

N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-3-(2-morpholinoethoxy)-benzamide (Compound 123)



0

[0253] A flask was charged with 2,4-dichloropyrimidine (22.7 g, 152.38 mmol), 4-acetoamido-phenylboronic acid (30.0 g, 167.62 mmol), dichloro[1,1'-bis(diphenylphosphino)-ferrocene-palladium (16.726 g, 22.86 mmol, 15 mol %), and triethylamine (53 mL, 380.95 mmol). Ethyleneglycoldimethylether (500 mL) and H₂O (20 mL) were added to the flask. The reaction mixture was stirred at 80 °C for 4 hours. The

5

5 product, Intermediate A, was isolated by removal of the solvent with a rotary evaporator and purified using glass column chromatography and eluted with ethyl acetate to afford 30.5 g (123.14 mmol, 81% yield) of intermediate A as a yellow solid.

[0254] A seal tube was charged with intermediate A (400 mg, 1.62 mmol) and 3-(tert-butoxycarbonylamino)aniline (1.99g, 9.57 mmol). N-butanol (50 mL) was added to the seal
10 tube and stirred at 180°C. The reaction was stopped after 2.5h, monitored by LCMS. The reaction mixture was diluted with methanol and the solid precipitate was filtered to afford intermediate B as a yellow solid. The filter pad was washed with ethyl-acetate, 72% yield. Intermediate B was carried on to the next step without further purification.

[0255] A flask was charged with intermediate B (159mg, 0.5 mmol), 3-(2-morpholinoethoxy)benzoyl chloride (169mg, 0.63 mmol), and Pyridine (8 mL). The reaction
15 mixture was stirred at RT under nitrogen. Reaction was complete after 1h. The final product 123 was purified using Preparative HPLC and trifluoroacetic acid buffer then free based with hydroxide resin in methanol. The filtrate was then concentrated, the yellow oil was then freeze-dried and lyophilized.


20 ¹H-NMR (400MHz, d₆-DMSO): 10.206(s, br, 2H), 9.665(s, br, 1H), 8.512(d, 1H), 8.502(s, 1H), 8.440(d, 2H), 7.755(d, 2H), 7.582(m, 2H), 7.483(m, 2H), 7.375(d, 1H), 7.287(m, 2H), 7.163(d, 1H), 4.188(m, 2H), 3.595 (m, 4H), 3.174(m, 4H), 2.732(m, 2H), 2.083(s, 3H).

MS(EI) for C₃₁H₃₂N₆O₄: 553 (MH⁺).


5

Example 40

4-[4-(methyamino)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (Compound 124)



[0256] A flask was charged with 2,4-dichloropyrimidine (810 mg, 5.5 mmol), 4-(tert-butoxycarbonyl(methyl)amino)phenylboronic acid **B** (4.93 g, 15 mmol), dichloro[1,1'-bis(diphenylphosphino)ferrocenepalladium (590 mg, 0.81 mmol, 15 mol %), triethylamine (1.8 mL, 13 mmol), and water (2 mL). Ethylene glycol dimethyl ether (5.0 mL) was added to the flask and the mixture was purged with N₂. The reaction mixture was stirred under an N₂ atmosphere at 90 °C for 1 hour, after which time, it was cooled to ambient temperature and filtered. The product, **C**, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LCMS: m/z 319 (M+H)⁺.




[0257] A flask containing a solution of **C** (1.9 g, 5.8 mmol) and 4-morpholinoaniline (1.5 g, 8.2 mmol) in 1-butanol (10 mL) was immersed in an oil bath at 180 °C for 4 h. The mixture was cooled to ambient temperature, concentrated, and the residue was dissolved in dichloromethane (10 mL) and 4N HCl in dioxane (10 mL). A portion of this crude product (200 mg) was purified by reverse phase HPLC to yield the product **124** (20 mg) in > 99% purity.


- 5 ¹H-NMR (400MHz, d₆-DMSO): 9.92-9.99 ppm (bs, 1H), 8.20-8.29 (bs, 1H), 8.02 (d, 2H), 7.52-7.68 (bs, 2H), 7.33 (d, 1H), 7.04-7.17 (bs, 1H), 6.67 (d, 2H), 3.71-3.82 (bs, 4H), 3.14-3.24 (bs, 4H), 2.78 (s, 3H). MS (EI) C₂₁H₂₃N₅O: 362.1 (MH⁺).

Example 41


- 10 **2,6-dichloro-N-{3-[(4-{[3-chloro-4-(methoxy)phenyl]oxy}pyrimidin-2-yl)amino]phenyl}-benzamide (Compound 304)**



- [0258] A flask was charged with 2,4-dichloropyrimidine (500 mg, 3.4 mmol), 2-chloro-4-methoxyphenol (580 mg, 3.7 mmol), and diisopropylethylamine (1.2 mL, 6.9 mmol).
 15 Dimethylformamide (20 mL) was added to the flask and the mixture was stirred at 70 °C for 15 hours. The reaction mixture was diluted with water and the mixture was extracted with dichloromethane 2X and 5% LiCl 3X. The crude product, **B**, was isolated by removal of the solvent with a rotary evaporator and the resultant brown oil was used without further purification. LCMS: m/z 272 (M+H)⁺.



- [0259] A flask containing a solution of intermediate **B** (910 mg, 3.4 mmol) and benzene-1,3-diamine (540 mg, 5.0 mmol) in nBuOH (5 mL) was immersed in an oil bath at 180 °C for 30 mins. The intermediate, **C**, was isolated by removal of the solvent with a rotary evaporator
 20 and used without further purification. LCMS: m/z 343 (M+H)⁺.




[0260] A flask was charged with intermediate **C** (1.1 g, 3.4 mmol), 2,6-dichlorobenzoylchloride (1.2 mL, 8.3 mmol), diispropylethylamine (1.8 mL, 10 mmol) and THF (50 mL). The reaction mixture was stirred at 60 °C for 15 hours. The reaction mixture was diluted with ethylacetate, extracted with 5% LiCl 3X, and the organic fraction was concentrated on a rotary evaporator. The crude product was purified by silica column chromatography (1:1 ethylacetate:hexanes as eluent) followed by reverse phase HPLC (TFA/ACN as eluent) to yield the product, **304** (24 mg, 1% yield).

¹H-NMR (400MHz, d₆-DMSO): 10.7 (s, 1H), 9.65 (s, 1H), 8.36 (d, 1H), 7.77 (s, 1H), 7.58-7.47 (m, 3H), 7.36-7.28 (m, 3H), 7.23 (d, 1H), 7.04-6.98 (m, 2H), 6.47 (d, 1H), 3.83 (s, 3H).

MS (EI) C₂₄H₁₇Cl₃N₄O₃: 514.8 (MH⁻).


Example 42

(3S)-1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrrolidine-3-carboxamide (Compound 510)



[0261] To a solution of **249(a)** (300 mg, 0.78 mmol) in DMA (10 mL) was added a solution of (S)-1-(tert-butoxycarbonyl)pyrrolidine-3-carboxylic acid (350 g, 1.6 mmol) diisopropyl-ethylamine (0.5 mL, 2.7 mmol), and HATU (600 mg, 1.6 mmol) in DMA (10 mL) and the solution was stirred at room temperature 15 hours. The solution was diluted with ethyl acetate (100 mL), washed with 10% LiCl (2X) and brine. The resultant solution

5 was dried over Na_2SO_4 , filtered and concentrated to yield a residue that was purified by silica gel column chromatography (3:1 ethyl acetate/hexanes). The Boc intermediate was isolated as a solid (340 mg, 78% yield). LC/MS: m/z 545 ($\text{M}+\text{H}$)⁺. A flask containing the Boc-intermediate was dissolved in 4N HCl in dioxane (10 mL) and dichloromethane (10 mL) and the mixture was stirred at room temperature for 15 hours. Intermediate C was isolated as a
 0 yellow solid after filtration and used without purification.




[0262] A flask was charged with intermediate C (450 mg, 0.78 mmol), 2-hydroxyacetaldehyde (45 mg, 0.75 mmol), sodium triacetoxyborohydride (150 mg, 0.71 mmol), diisopropylethylamine (0.7 mL, 3.8 mmol) and dichloromethane (20 mL) and the mixture
 15 was stirred at room temperature for 4 hours. The reaction mixture was diluted with water and the solution was extracted with saturated NaHCO_3 (2X) and brine. The residue was purified by reverse phase HPLC (ammonium acetate/ACN as eluent) to afford the product **510** (120 mg, 31% yield).

¹H-NMR (400MHz, d_6 -DMSO): 10.2 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.75
 20 (d, 2H), 7.67 (d, 2H), 7.28-7.27 (m, 1H), 6.93 (d, 2H), 4.47 (br 1H), 3.76-3.73 (m, 4H), 3.49 (t, 2H), 3.06-3.03 (m, 4H), 2.91 (t, 1H), 2.70-2.65 (m, 1H), 2.58-2.56 (m, 1H), 2.54-2.49 (m, 4H), 1.99 (t, 2H). MS (EI) $\text{C}_{27}\text{H}_{32}\text{N}_6\text{O}_3$: 489.2 ($\text{M}+\text{H}$)⁺.


5

Example 43

N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}phenyl)acetamide (Compound 329)



- 10 [0263] A mixture of 2-[(4-morpholin-4-ylphenyl)amino]-3,7-dihydro-4H-pyrrolo[2,3-d]pyrimidin-4-one A (312 mg, 1 mmol), phosphorous oxybromide (717 mg, 2.5 mmol), and diisopropylethylamine (130 mg, 1 mmol) in anhydrous toluene (15 ml) was heated at reflux under N₂ overnight. The mixture was cooled down to room temperature, and the solid was filtered, washed with sat. NaHCO₃, water, and dried over MgSO₄. The solvent was removed
- 15 *in vacuo* to give the product 4-bromo-N-(4-morpholin-4-ylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine B (284 mg, 76%) as a black solid. This was clean and used as such without further purification.



- [0264] A mixture of 4-bromo-N-(4-morpholin-4-ylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine B (284 mg, 0.76 mmol), 4-acetoamidophenylboronic acid C (340 mg, 2.5 eq), tetrakis(triphenyl-phosphine)palladium(0) (120 mg, 0.1 mmol), and 1M Na₂CO₃ (1ml, 1 mmol) in 1,4-dioxane (15 ml) was heated at reflux overnight. The mixture was cooled, extracted with 3N HCl. The aqueous layer was washed with ethylacetate, and then basified with 6N NaOH. The solid was filtered, and the crude product was purified by preparative
- 25 HPLC to give the product N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}phenyl)acetamide D (0.8 mg, 0.25%) as a yellow solid.


- 5 ¹H NMR (400 MHz, CD₃OD): 8.10 (d, 2H), 7.75 (d, 2H), 7.68 (d, 2H), 7.12 (d, 1H), 6.98 (d, 2H), 6.70 (d, 1H), 3.85 (t, 4H), 3.09 (t, 4H), 2.17 (s, 3H). MS (EI) for C₂₄H₂₄N₆O₂: 429 (MH⁺).

Example 44

- 10 **2-Methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide (Compound 367)**

Preparation of *tert*-butyl 2-methyl-2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylcarbamoyl)pyrrolidine-1-carboxylate


15



- [0265] An oven dried 50 ml round bottomed flask fitted with a Teflon stirrer and gas inlet was flushed with dry nitrogen and allowed to cool to room temperature. The flask was
- 10 charged with 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine pentahydrochloride (1 equiv., 0.52 g, 0.9631 mmoles) and anhydrous dimethylacetamide (15 ml). The mixture was stirred for 10 minutes to allow for the complete dissolution of the amine. Diisopropylethylamine (10 equiv., 1.24 g, 1.67 ml, 9.631 mmoles) was added in one lot and the reaction mixture was stirred for 5 minutes. 1-(*tert*-Butoxycarbonyl)-2-
- 15 methylpyrrolidine-2-carboxylic acid (4 equiv., 3.852 mmoles, 0.883 g, purchased from Fluka-Sigma Aldrich) was added to the reaction mixture in one lot, followed by 2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU, 4 equiv., 3.852 mmoles, 1.464 g, purchased from Oakland Products). The reaction mixture was stirred at room temperature and the progress of the reaction was monitored by LC/MS. After 72 hours,
- 0 the reaction mixture was quenched with ethyl acetate (20ml), and transferred to separatory funnel. The reaction flask was further rinsed with ethyl acetate (20 ml), transferred to the

5 separatory funnel, shaken and the layer separated off. The aqueous layer was further washed with ethyl acetate (3 x 50 ml). The combined ethyl acetate solutions were washed with cold water (2 x 50 ml) and saturated sodium chloride solution (2 x 50 ml). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to give orange oil. The resulting crude material was purified by silica phase flash chromatography (45 mm x 250 mm) using 3:1 ethyl acetate-hexane to give 0.147 g of *tert*-butyl 2-methyl-2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylcarbamoyl)-pyrrolidine-1-carboxylate as a white solid (27% yield). ¹H NMR (400 MHz, d₆-DMSO) 10.01 (br s, 1H), 9.45 (br s, 1H), 8.19 (d, 1H), 7.70 (d, 2H), 7.46 (d, 2H), 6.74 (d, 1H), 6.66 (d, 2H), 6.28 (d, 2H), 3.67 (m, 4H), 3.40 (m, 1H), 3.30 (m, 1H), 2.29 (m, 4H), 1.76 (m, 1H), 1.64 (m, 1H), 1.58 (s, 3H), 1.54 (m, 1H), 1.40 (s, 9H). MS (EI) for C₃₁H₃₈N₆O₄: 559 (M⁺).

Preparation of 2-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)prolinamide




[0266] *tert*-Butyl 2-methyl-2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylcarbamoyl)pyrrolidine-1-carboxylate (0.140 g, 0.250 mmoles), was dissolved in an ethyl acetate (5 ml) and methanol (1 ml) mixture. 4 M hydrogen chloride in 1,4-dioxane (0.625 ml, 2.5 mmoles, 10 equivalents, purchased from Sigma-Aldrich) was then added in a drop wise fashion over 5-10 minutes. Upon completion of addition, the reaction mixture was stirred at room temperature, and the progress of the reaction monitored by LC/MS. After 16 hours, additional 4M hydrogen chloride in 1,4-dioxane (0.312 ml, 1.25 mmoles, 5 equivalents) was added. After a total of 48 hours the reaction was complete and the resulting slurry was filtered off. The reaction flask was rinsed with ethyl acetate to ensure complete transfer of product. The resulting solid was washed with ethyl acetate (3 x 10 ml) and diethyl ether (2 x

5 25 ml) and dried under reduced pressure to give of 0.061 mg 2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide **367** as its hydrochloride salt (53% yield).

¹H NMR (400 MHz, d₆-DMSO): 10.88 (s, 1H), 9.79 (br s, 1H), 8.47 (d, 1H), 8.12 (d, 1H), 8.10 (d, 1H), 7.80 (br d, 2H), 7.75 (d, 2H), 7.37 (d, 2H), 5.26 (br s, 3H), 3.72 (br s, 4H), 3.27 (br s, 4H), 2.80 (m, 1H), 2.70 (m, 1H), 2.01 (m, 1H), 1.76 (m, 1H), 1.64 (m, 1H), 1.54 (m, 1H), 1.38 (s, 3H). MS (EI) for C₂₆H₃₀N₆O₂: 459 (MH⁺).

Example 45

15 **2-Methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide (Compound 360)**




[0267] An oven dried 50 ml round bottomed flask fitted with a Teflon stirrer and gas inlet was flushed with dry nitrogen and allowed to cool to room temperature. The flask was charged with 4-(4-aminophenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine pentahydrochloride (1 equiv., 0.4 g, 0.756 mmoles) and anhydrous dimethylacetamide (15 ml). The mixture was stirred for 10 minutes to allow for the complete dissolution of the amine. Diisopropylethylamine (10 equiv., 0.977 g, 1.31 ml, 7.561 mmoles) was added in one lot and the reaction mixture was stirred for 5 minutes. N-Boc-D-proline (4 equiv., 3.204 mmoles, 0.65 g, purchased from Fluka-Sigma Aldrich) was added to the reaction mixture in one lot, followed by 2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU, 4 equiv., 3.024 mmoles, 1.149 g, purchased from Oakland Products). The reaction mixture was stirred at room temperature and the progress of the reaction was monitored by LC/MS. After 72 hours, the reaction mixture was quenched with ethyl acetate (20ml), and transferred to separatory funnel. The reaction flask was further rinsed with ethyl acetate (20 ml), transferred to the separatory funnel, shaken and the layer

5 separated off. The aqueous layer was further washed with ethyl acetate (3 x 50 ml). The combined ethyl acetate solutions were washed with chloride solution (2 x 50 ml). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered and evaporated under reduced pressure to give an orange oil. The resulting crude material was purified by silica phase flash chromatography (45 mm x 250 mm) using 3:1 ethyl acetate -hexane to give 0.39 g of 1,1-dimethylethyl (2R)-2-[[[4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]carbonyl]pyrrolidine-1-carboxylate as a white solid (94 % yield).

10 ¹H NMR (400 MHz, d₆-DMSO): 10.26 (br s, 1H), 9.38 (br s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.78 (d, 2H) < 7.68 (d, 2H), 7.28 (d, 1H), 6.94 (d, 2H), 4.22 (m, 1H), 3.74 (m, 4H), 3.43 (m, 1H), 3.34 (m, 1H), 3.04 (m, 4H), 2.20 (m, 1H), 1.90 (m, 1H), 1.81 (m, 1H) < 1.40 (s, 3H), 1.27 (s, 6H). MS (EI) for C₃₀H₃₆N₆O₄: 545 (MH⁺).

15



[0268] 1,1-Dimethylethyl (2R)-2-[[[4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)amino]carbonyl]pyrrolidine-1-carboxylate (0.38 g, 0.698 mmoles), was dissolved in an ethyl acetate (10 ml) and methanol (2 ml) mixture. 4 M hydrogen chloride in 1,4-dioxane (1.75 ml, 6.98 mmoles, 10 equivalents, purchased from Sigma-Aldrich) was then added in a drop wise fashion over 5-10 minutes. Upon completion of addition, the reaction mixture was stirred at room temperature, and the progress of the reaction monitored by LC/MS. After 16 hours, additional 4M hydrogen chloride in 1,4-dioxane (0.87, 1.25 mmoles, 5 equivalents) was added. After a total of 48 hours the reaction was complete and the resulting slurry was filtered off. The reaction flask was rinsed with ethyl acetate to ensure complete transfer of product. The resulting solid was washed with ethyl acetate (3 x 10 ml), followed by diethyl ether (3 x 25 ml) and dried under reduced pressure to give of 0.264 mg 2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-pyrimidin-4-yl}phenyl)prolinamide (68 % yield).

20

25


30

5 ¹H NMR (400 MHz, d₆-DMSO): 11.43 (br s, 1H), 10.07 (br s, 2H), 8.73 (d, 1H), 8.57 (d, 1H), 8.21 (d, 2H), 7.91 (d, 2H), 7.98 (d, 2H), 7.71 (br s, 2H), 7.48 (d, 1H), 4.48 (m, 1H), 4.08 (s, 4H), 3.74 (m, 4H), 3.42 (m, 1H), 3.36 (m, 1H), 3.04 (m, 4H), 2.22 (m, 1H), 1.90 (m, 2H), 1.82 (m, 2H). MS (EI) for C₂₅H₂₈N₆O₂: 445 (MH⁺).

10

Example 46

3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(1-methyl-1H-benzimidazol-2-yl)methyl]benzamide (Compound 83)



15


[0269] 7.9 grams (11.04 mmol, 1.9 eq) of PL-TFP Resin (source: *Polymer Laboratories*) was weighed into a pressure tube. 60 ml of DCM was added. 2 g (5.74 mmol) of 3-(4-(3-acetamidophenyl)pyrimidin-2-ylamino)benzoic acid was dissolved in 15 ml of DMF and
 20 After 10 min, this solution was added to the pressure tube. Dimethylaminopyridine (4.41 mmol, .6 eq, source: *Acros*) was added to the pressure tube as a solid, followed by 1,3-diisopropylcarbodiimide (33.08 mmol, 4.5 eq, source: *Acros*). The pressure tube was sealed and the reaction was placed on a vertical shaker overnight. The resin was filtered, and then washed 3 times with DMF, followed by three times with THF, followed by three times with
 25 DCM. The resin was then dried overnight by vacuum.

[0270] 300 mg of resin prepared above (loading = 0.6 mmol/g, .18 mmol) was added to a 1 dram vial. 2 ml of DMA were added. 1 ml of (1-methyl-1H-benzo[d]imidazol-2-yl)methanamine (0.12 mmol, 0.67 eq) dissolved in DMA was added to the vial. The reaction was stirred overnight at room temperature. The reaction was filtered and rinsed twice with 4
 30 ml of MeOH. The solution was further purified by HPLC to yield (3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)-N-((1-methyl-1H-benzo[d]imidazol-2-yl)methyl)benzamide 83 (10.2 mg, 17%).

- 5 ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.80 (s, 1H), 9.01 (t, 1H), 8.52 (t, 2H), 8.17-8.19 (m, 2H), 7.91-7.93 (m, 1H), 7.75 (d, 2H), 7.50-7.59 (m, 3H), 7.39-7.43 (m, 2H), 7.16-7.26 (m, 2H), 4.80 (d, 2H), 3.86 (s, 3H), 2.10 (s, 3H). MS (EI) for C₂₈H₂₅N₇O₂: 492.4 (MH⁺).

Example 47

- 10 **N-(4-morpholin-4-ylphenyl)-4-{4-[(propylamino)methyl]phenyl} pyrimidin-2-amine (Compound 283)**



- [0271] A flask was charged with 2,4-dichloropyrimidine (1.5 g, 10 mmol), 4-formylphenyl boronic acid (1.65 g, 11 mmol), dichloro[1,1'-bis(diphenylphosphino)-ferrocenepalladium (731 mg, 1 mmol, 10 mol %), and triethylamine (2.6 mL, 15 mmol). Ethyleneglycoldimethylether (50 mL) and H₂O (2mL) was added to the flask. The reaction mixture was stirred at 80 °C for 4 hours. The product, Intermediate A, was isolated by removal of the solvent with a rotary evaporator and purified using glass column chromatography and eluted with ethyl acetate to afford 1.0 g (4.58 mmol, 46% yield) of intermediate A as a yellow solid.
- 15
- 20

[0272] A flask was charged with intermediate A (150 mg, 0.668 mmol), sodium triacetoxy-borohydride (220 mg, 1.032 mmol), propylamine (63 μ L, 0.756 mmol). Dichloromethane (50 mL) was added to the flask and the reaction mixture was stirred at room temperature for 48 h. The reaction was quenched with 2 N NaOH and extracted with ethyl


acetate, washed with brine, dried over sodium sulfate, and filtered. The solvent was removed using the rotary evaporator to afford intermediate **B** as a yellow solid (140 mg, 0.536 mmol, 80% Yield). Intermediate **B** was carried on without further purification.

[0273] A seal tube was charged with intermediate **B** (140 mg, 0.536 mmol) and 4-morpholinoaniline (95 mg, 0.536 mmol). N-butanol (15 mL) was added to the seal tube and stirred at 180°C. The reaction was done in 1h according to LCMS to afford **283** as a yellow solid. **Compound 283** was purified using preparative HPLC and TFA buffer. **Compound 283** was free-based, converted to HCl salt, and lyophilized (20mg, 0.455 mmol).

¹H NMR (400 MHz, DMSO): 9.93 (br s, 1H), 9.396 (br s, 2H), 8.607 (d, 1H), 8.233 (d, 2H), 7.905 (d, 2H), 7.767 (d, 2H), 7.571-7.494 (m, 3H), 4.226 (br s, 2H), 3.998 (br s, 4H), 3.436 (br s, 4H), 2.865 (m, 2H), 1.708 (m, 2H), 0.914 (t, 3H). MS (EI) for C₂₄H₂₉N₅O: 404.4 (MH⁺).

Example 48

N-[(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)methyl]acetamide
(**Compound 282**)




[0274] In a 20 ml round bottomed flask, 36 mg (1 mmol) of compound A was dissolved in 5 ml of dichloromethane and 0.5 ml of triethylamine was added. It was cooled in ice bath and 10 mg (1.2 mmol) of acetyl chloride was added and stirred for 30 min. **Compound 282** precipitated out and purified in a Waters prep column. Yield 40 mg (90%).

¹H NMR (400MHz, CD₃CN): 11.20-11.22(b,1H), 8.40 (d, 2H), 8.05(d,2H), 7.80(d,2H), 7.50 (d, 2H), 7.45 (s, 1H), 7.20 (d, 1H), 7.05-7.10 (b, 1H), 4.40-4.44 (b, 2H), 3.90 (t, 4H), 3.40 (t, 4H), 2.01(s,3H); MS (EI) for C₂₃H₂₅N₅O₂: 404 (MH⁺).

5

Example 49


N-(4-{2-[(4-{4-[(2,4-dichlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 180)



[0275] To a 1 ml vial was added N-(4-{2-[(4-piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)-phenyl)acetamide (38.85 mg, 0.1 mmol), 2,4-dichlorobenzaldehyde (350 mg, 2.0 mmol, 20 eq, source: Aldrich) and 1 ml of DMF. To this mixture was added sodium triacetoxymethylborohydride (106 mg, 0.5 mmol, 5 eq). The mixture was stirred over night at room temperature. Upon completion of the reaction as determined by LC/MS, 0.1 ml of 2M HCl was added. The residue was purified via reverse phase HPLC (ammonium acetate/ACN) to yield N-(4-{2-[(4-{4-[(2,4-dichlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide **180** (20.2 mg, 37 %).

20 ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.80 (s, 1H), 9.10 (t, 1H), 8.53 (d, 2H), 8.18 (d, 2H), 7.94-7.91 (m, 1H), 7.75 (d, 1H), 7.69-7.57 (m, 4H), 7.48-7.39 (m, 2H), 4.58 (d, 6H), 2.50 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₈Cl₂N₆O: 548.5 (MH⁺).

5

Example 50**5-fluoro-N-[2-[(2-methoxy)phenyl]-N-[3-[(methoxy)phenyl]]pyrimidine-2,4-diamine (Compound 306)**


- 10 [0276] A round-bottomed flask was charged with 2,4-dichloro-5-fluoropyrimidine (0.84 g, 5 mmol), 2-methoxyaniline (0.61 g, 5 mmol) and dioxane (5 mL). The reaction mixture was heated at 85 °C overnight. The reaction was cooled down and diluted with acetonitrile/water, stirred for 30 min. and filtered. The collected solid was re-suspended in acetonitrile/water, stirred and filtered to give a 2-chloro-5-fluoro-N-(2-methoxyphenyl)pyrimidin-4-amine (0.9 g, 70% yield).
- 15 To a seal tube was added 2-chloro-5-fluoro-N-(2-methoxyphenyl)pyrimidin-4-amine (254 mg), 3-methoxyaniline (500 mg, 4 eq.) and dioxane (5 mL). The mixture was heated to 130 °C overnight. The reaction mixture was cooled and partitioned between EtOAc and water. The organic layer was concentrated; the residue was triturated with a 1:1 mixture of dichloromethane and acetonitrile, then filtered to give the title product as a white solid
- 20 (150 mg).

¹H NMR (400 MHz, d₆-DMSO): 9.16 (s, 1H), 8.37 (s, 1H), 8.09 (s, 1H), 7.88 (d, 1H), 7.26 (t, 2H), 7.22-7.16 (m, 2H), 7.12-7.08 (m, 1H), 7.08-6.93 (m, 2H), 3.81 (s, 3H), 3.62 (s, 3H). MS (EI) for C₁₈H₁₇FN₄O₂: 341 (MH⁺).


5

Example 51

N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}phenyl)-acetamide (Compound 329)



10



5


[0277] To a flask containing a solution of 1 (0.25g, 1 mmol), 4-(4-ethylpiperazin-1-yl)aniline (0.23g, 1.1 mmol), cesium carbonate (0.5g, 1.5 mmol), *racemic*-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (95mg, 0.15 mmol) in N,N-dimethylacetamide (5mL) purged with N₂ was added tris(dibenzylideneacetone)dipalladium(0) (0.14g, 0.15 mmol). This reaction was heated to 90 °C for 16 h under N₂. At this time the reaction was concentrated and the residue was purified via silica gel column chromatography. The column was eluted with ethyl acetate to remove the impurities and then with 85:10:5 (ethyl acetate/methanol/7M ammonia in methanol) to elute the desired product. The solid obtained was sonicated first in acetone (5mL) and then in ether (10mL) to yield intermediate 2 (0.23g, 48% yield) as a yellow solid. LCMS: m/z 417 (MH⁺). To a flask containing 2 (0.23g, 0.45 mmol) was added 4N HCl in dioxane (5mL) and the solution was heated at 50 °C for 4 h. To the cooled solution was added a 2N aqueous solution of sodium hydroxide (10mL) and the resulting precipitate was filtered and dried to yield 3 (0.2g, 99% yield) as a yellow solid. LCMS: m/z 375 (M+H)⁺. To a flask with 3 (0.3g, 0.8 mmol), phenylacetic acid (0.125mL, 1 mmol), triethylamine (0.97mL, 7 mmol), and DMF (5mL) was added O-(7-azabenzotriazole-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.46g, 1.2 mmol). The reaction mixture was stirred at ambient temperature for 1 h then diluted with 5% aqueous

5 solution of lithium chloride (100mL) and extracted with ethyl acetate (3 x 50mL). The combined organic layers were washed with an aqueous 5% sodium bicarbonate solution, a saturated aqueous solution of sodium chloride, dried over anhydrous sodium sulfate, filtered and concentrated. The residue obtained was purified by silica gel column chromatography (98:2 ethylacetate/methanol) to provide Compound 329 (0.25g, 63% yield) as an off-white
 10 solid.


Example 52

N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide (Compound 662)

15




[0278] A solution of chloropyrimidine (1) (0.28g, 0.64 mmol) and tert-butyl 4-(4-aminophenyl)piperazine-1-carboxylate (0.18g, 0.6 mmol) in n-butanol (5ml) was heated at
 20 180 °C in a sealed tube for 7 h. The reaction mixture was concentrated, the residue dissolved in methanol (5ml) and treated with HCl (3ml, 4M in dioxane) for 1 h at room temperature. After concentration, the residue was dissolved in H₂O (200mL) and the pH adjusted to ca. 8-9
 25 organic layers were dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel column chromatography (85:15 ethyl acetate/methanol) to afford 2 (0.26g, 69%). C₃₃H₃₅N₇O₃: 578 (MH⁺).



[0279] To a solution of **2** (0.41g, 0.7 mmol) and DIPEA (0.31ml, 1.75 mmol) in CH₂Cl₂ (7ml) was added cyclobutylcarbonyl chloride (0.80ml, 0.7 mmol) at room temperature. After 10 min, the reaction mixture was diluted with H₂O (10mL) and CH₂Cl₂ (10mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 5mL). The combined organic layers were dried over anhydrous sodium sulfate, filtered, and concentrated. The residue obtained was purified by silica gel column chromatography (98:2 ethyl acetate/methanol) to afford **3** (0.31g, 68%) as a white powder. C₃₈H₄₁N₇O₄: 660 (MH⁺).


10




[0280] The mixture of **3** (0.31g, 0.47 mmol) and Pd/C (0.94g) AcOH (1mL) and MeOH (5mL) was stirred at room temperature for 24 h under a H₂ balloon. The palladium was filtered through celite and the filtrate was concentrated. The residue was purified by silica gel column chromatography (90:10 ethyl acetate/methanol) to afford product which was then washed with acetonitrile several times to afford Compound **662** (0.14g, 59% yield).

20

5


Example 53**N-ethyl-4-[4-({4-[4-(D-prolylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazine-1-carboxamide (Compound 663)**

- [0281] To a stirred solution of 2 (0.58g, 1 mmol) in DMF (4ml) was added ethyl isocyanate (3mL) at room temperature. After stirring for 30 min, the mixture was diluted with H₂O (5mL) and CH₂Cl₂ (5mL). The separated aqueous layer was extracted with CH₂Cl₂ (3 x 5mL). The combined extracts were dried over anhydrous sodium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography (98:2 ethyl acetate/methanol) to afford 4 (0.46g, 71%). C₃₆H₄₀N₈O₄: 649 (M+H)⁺.



- [0282] A solution of 4 (0.46g, 0.71 mmol) and Pd/C (0.14g) in AcOH (1mL) and MeOH (5mL) was stirred for 24 h under a H₂ balloon. The palladium was filtered through celite and the filtrate was concentrated. The residue obtained was purified by silica gel column chromatography (90:10 ethyl acetate/methanol) to afford product which was then washed with acetonitrile several times to afford N-ethyl-4-[4-({4-[4-(D-prolylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazine-1-carboxamide (663) (0.19g, 51%) as a white powder.

5

2-[[2-(dimethylamino)ethyl]oxy]-3-(4-ethylpiperazin-1-yl)aniline

[0283] To a solution of 2-(diethylamino)ethanol (0.59g, 5mmol) in DMA (5ml), sodium
 10 hydride (0.24g, 10mmol) was added in one portion. Fifteen minute later 2-bromo-4-
 nitroaniline (1.1g, 5mmol) was added and the content was stirred for 4 hr. Water (50ml) was
 added to the reaction mixture followed by chloroform. The organic layer was separated,
 washed with saturated sodium bicarbonate follow by brine. The organic layer was dried over
 sodium sulfate and concentrated to oil. The oil was dissolved in methanol and saturated with
 15 HCl gas. The resulting solution was concentrated and diethyl ester was added. The resulting
 precipitate was washed with ether and dried to yield 0.8g of 2-[(2-bromo-4-
 nitrophenyl)oxy]-N,N-diethylethanamine hydrochloride as a solid. LCMS: m/z 318 (M+H)⁺.

[0284] A mixture of 2-[(2-bromo-4-nitrophenyl)oxy]-N,N-diethylethanamine
 hydrochloride (0.5g, 1.4mmol), Pd(dba)₂ (0.192g, 0.21mmol), BINAP (0.139g, 0.21mmol), 1-
 20 ethylpiperazine (0.182g, 1.68mmol), and cesium carbonate (0.91g, 2.8mmol) in DMA was
 heated at 80 °C with stirring for 72 hr. Saturated aqueous sodium bicarbonate and ethyl
 acetate was added, the phases separated, the solvent was removed under vacuum and the
 residue chromatographed on silica with ethyl acetate/methanol to give 0.32g of N,N-diethyl-
 2-[[2-(4-ethylpiperazin-1-yl)-4-nitrophenyl]oxy]ethanamine. LCMS: m/z 351 (M+H)⁺.

[0285] A solution of N,N-diethyl-2-[[2-(4-ethylpiperazin-1-yl)-4-
 25 nitrophenyl]oxy]ethanamine (0.280mg, 0.8mmol) in methanol (10ml) was added 10% Pd/C
 and stirred in hydrogen atmosphere at ambient temperature for 2hr. The reaction mixture was
 filtered through a pad of Celite and concentrated in vacuum. The residue was taken up in

5 methanol, ether/HCl was added and the hydrochloride (0.270 mg) was precipitated. LCMS: m/z 321 (M+H)⁺.


Example 54

10 **(R)-N-(4-(2-(3-(benzyloxy)-4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide (Compound 376)**

[0286] *R*-N-(4-(2-(3-(benzyloxy)-4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide was synthesized in an analogous fashion to Example 3, wherein 4-morpholinoaniline was substituted with 3-(benzyloxy)-4-morpholinoaniline to afford the title compound.

15

3-(benzyloxy)-4-morpholinoaniline



20 [0287] **4-(2-(Benzyloxy)-4-nitrophenyl)morpholine:** A flask was charged with 2-chloro-5-nitrophenol (3.5 g, 20.2 mmol), potassium carbonate (4.0g, 30.3 mmol), benzyl bromide (2.9 mL, 24.24 mmol,) and acetonitrile (25 mL). The reaction mixture was stirred under an N₂ atmosphere at room temperature for 12 hours, after which time, the reaction mixture was filtered through Celite pat and washed with ethyl acetate (50 mL). The product was isolated by removal of the solvent with a rotary evaporator and used without further purification.


25 NMR (400 MHz, CDCl₃): 7.80 (m, 2H), 7.27 – 7.58 (m, 6H), 5.24 (s, 2H). MS (EI) for C₁₃H₁₀ClNO₃: 264 (MH⁺).

5

Example 55

[0288] (*S*)-2-amino-*N*-(4-(2-(3-methyl-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide (**Compound 384**) was synthesized in an analogous fashion to Example 3, wherein 4-morpholinoaniline was substituted with 3-Methyl-4-morpholinoaniline to afford the title compound.

10

3-Methyl-4-morpholinoaniline

[0289] A flask was charged with 2-fluoro-5-nitrotoluene (3.0 mL, 20.2388 mmol). Excess amount of morpholine (10 mL) were added to the flask and heated to 40 °C for 6 hours. The reaction mixture was checked with LC/MS. Water was added to the reaction mixture and the precipitate, intermediate 1, was filtered and used without further purification. LC/MS: m/z 223 ($M+H$)⁺.

[0290] A hydrogenation flask was charged with intermediate 1 (1.0 g, 4.4209 mmol) and palladium/carbon (200 mg). Ethyl alcohol (50 mL) was added to the flask and hydrogenation technique was used. The reaction mixture was checked with LC/MS. The reaction mixture was filtered through a celite plug and washed with methanol. The product, 3-methyl-4-morpholinoaniline, was isolated by removal of the solvent with a rotary evaporator and used without further purification. LC/MS: m/z 197 ($M+H$)⁺.

25

Example 56

[0291] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-fluoro-3-(methoxy)benzamide (**Compound 49**) was synthesized in an analogous fashion to Example 2, wherein benzoylchloride was substituted with 2-chloro-3-methoxy-6-fluorobenzoylchloride (JRD Fluorochemicals) to afford the title compound.

30

5

Example 57

[0292] *N*-(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 296) was synthesized in an analogous fashion to Example 3, wherein aniline was substituted with 3-chloro-4-morpholinoaniline (Pfaltz and Bauer, Inc.) to afford the title compound.

10

Example 58

[0293] *N*-(4-{2-[(3-bromo-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Compound 315) was synthesized in an analogous fashion to Example 3, wherein aniline was substituted with 3-bromo-4-morpholinoaniline (Ryan Scientific, Inc.) to afford the title compound.

15

Example 59

[0294] (*R*)-*N*-(4-(2-(4-morpholino-3-(trifluoromethyl)-phenylamino)pyrimidin-4-yl)-phenyl)-pyrrolidine-2-carboxamide was synthesized in an analogous fashion to Example 3, wherein aniline was substituted with 3-trifluoromethyl-4-morpholinoaniline (Zerenex Limited) to afford the title compound.

20

Example 60

[0295] (*R*)-*N*-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide was synthesized in an analogous fashion to Example 3, wherein aniline was substituted with 3-fluoro-4-morpholinoaniline (Astatech, Inc.) to afford the title compound.

25

Example 61

[0296] *N*-[4-(2-{[3-(1,3-dioxan-2-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide was synthesized in an analogous fashion to Example 3, wherein aniline was substituted 3-(1,3-dioxan-2-yl)aniline (Oakwood Products, Inc.) to afford the title compound.

30

Example 62

[0297] *N*-(4-{2-[(4-morpholin-4-ylphenyl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}phenyl)acetamide was synthesized in an analogous fashion to Example 5, wherein

35

5 pyrimidine was substituted with 5-trifluoromethyl-2,4-dichloropyrimidine (Astatech, Inc.) to afford the title compound.

[0298] Using the same or analogous techniques as illustrated in the preceding examples the following compounds herein below were made. The skilled artisan would be able to make the necessary modifications and/or substitutions in the above synthetic procedures to arrive at the following compounds:

[0299] *N*-(4-{2-[(3-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 21): ¹H-NMR (400MHz, d₆-DMSO): 10.22 ppm (s, 1H), 9.511 ppm (s, 1H), 8.504 ppm (d, 1H), 8.154 ppm (d, 2H), 7.76 ppm (d, 3H), 7.343 ppm (d, 1H), 7.215-7.153 ppm (m, 2H), 6.584 ppm (d, 1H), 3.775 ppm (t, 4H), 3.14 ppm (t, 4H), 2.094 ppm (s, 3H); MS (EI) C₂₂H₂₃N₅O₂: 390.1 (MH⁺).

[0300] *N*-(4-{2-[(3-piperidin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 22): ¹H-NMR (400MHz, d₆-DMSO): 10.231 ppm (s, 1H), 9.466 ppm (s, 1H), 8.497 ppm (d, 1H), 8.16 ppm (d, 2H), 7.765 ppm (d, 3H), 7.337 ppm (d, 1H), 7.119 ppm (d, 2H), 6.553 ppm (m, 1H), 3.176 ppm (t, 4H), 2.092 ppm (s, 3H), 1.658 ppm (m, 4H), 1.571 ppm (m, 2H); MS (EI) C₂₃H₂₅N₅O: 388.1 (MH⁺).

[0301] *N*-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide (Cmpd 33): ¹H-NMR (400MHz, d₆-DMSO): 10.269 ppm (s, 1H), 9.317 ppm (s, 1H), 8.411 ppm (d, 1H), 8.089 ppm (d, 2H), 7.743 ppm (d, 2H), 7.638 ppm (d, 2H), 7.243 ppm (d, 1H), 6.908 ppm (d, 2H), 3.048 ppm (br s, 4H), 2.350 ppm (q, 2H), 2.07 ppm (s, 3H), 1.027 ppm (t, 3H); MS (EI) C₂₄H₂₈N₆O: 417.4 (MH⁺).

[0302] *N*-(4-{2-[(4-piperidin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 34): ¹H-NMR (400MHz, d₆-DMSO): 10.229 ppm (s, 1H), 9.53 ppm (s, 1H), 8.481 ppm (d, 1H), 8.135 ppm (d, 2H), 7.76 ppm (t, 4H), 7.325 ppm (d, 1H), 7.178 ppm (d, 2H), 3.025 ppm (br d, 2H), 2.59 ppm (m, 2H), 2.094 ppm (s, 3H), 2.08 ppm (br d, 2H), 1.54-1.439 ppm (m, 2H); MS (EI) C₂₃H₂₅N₅O: 388.3 (MH⁺).

[0303] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide (Cmpd 17): ¹H-NMR (400MHz, d₆-DMSO): 10.718 ppm (s, 1H), 10.269 ppm (s, 1H), 9.678 ppm (s, 1H), 8.507 ppm (d, 1H), 8.419 ppm (s, 1H), 8.215 ppm (d, 2H), 7.758 ppm (d, 2H), 7.608 ppm (d, 2H), 7.532 ppm (t, 1H), 7.472 ppm (d, 1H), 7.380 ppm (d, 1H), 7.301 ppm (t, 1H), 7.216 ppm (d, 1H), 2.085 ppm (s, 3H); MS (EI) C₂₅H₁₉Cl₂N₅O₂: 492.2 (MH⁺).

- 5 [0304] *N*-{4-[2-({3-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide (Cmpd 8): ¹H-NMR (400MHz, d₆-MEOD): 8.455 ppm (d, 1H), 8.15 ppm (m, 3H), 7.76-7.7 ppm (m, 3H), 7.435 ppm (t, 1H), 7.311 ppm (d, 1H), 7.1 ppm (d, 1H), 3.832 ppm (br s, 4H), 3.13 ppm (br s, 4H), 3.016 ppm (q, 2H), 2.162 ppm (s, 3H), 1.975 ppm (s, 3H, ACE), 1.299 ppm (t, 3H); MS (EI) C₂₅H₂₈N₆O₂: 445.4 (MH⁺).
- 10 [0305] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluorobenzamide (Cmpd 10): ¹H NMR (DMSO-*D*₆) 10.40 (S, 1H), 10.21 (S, 1H), 9.66 (S, 1H), 8.49 (D, 1H), 8.41 (S, 1H), 8.20 (D, 2H), 7.74 (D, 2H), 7.69 (M, 1H), 7.58 (M, 1H), 7.48 (M, 1H), 7.37 (M, 3H), 7.28 (M, 2H), 2.09 (S, 3H). LCMS: M/Z 442 (M+H)⁺.
- 15 [0306] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluoro-6-iodobenzamide (Cmpd 11): ¹H NMR (DMSO-*D*₆) 10.65 (S, 1H), 10.19 (S, 1H), 9.67 (S, 1H), 8.50 (D, 1H), 8.41 (S, 1H), 8.21 (D, 2H), 7.76 (M, 3H), 7.41 (M, 3H), 7.28 (M, 3H), 2.08 (S, 3H). LCMS: M/Z 567 (M+H)⁺.
- 20 [0307] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-bromobenzamide (Cmpd 23): ¹H NMR (DMSO-*D*₆) 10.48 (S, 1H), 10.20 (S, 1H), 9.66 (S, 1H), 8.49 (D, 1H), 8.42 (S, 1H), 8.20 (D, 2H), 7.73 (D, 3H), 7.55 (M, 2H), 7.45 (M, 2H), 7.36 (M, 1H), 7.37 (D, 1H), 7.25 (M, 2H), 2.08 (S, 3H). LCMS: M/Z 502, 503, 504, 505 (M+H)⁺.
- 25 [0308] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3-fluorobenzamide (Cmpd 24): ¹H NMR (DMSO-*D*₆) 10.33 (S, 1H), 10.23 (S, 1H), 9.69 (S, 1H), 8.50 (D, 1H), 8.44 (S, 1H), 8.21 (D, 2H), 7.85 (M, 1H), 7.79 (M, 1H), 7.73 (D, 2H), 7.61 (M, 1H), 7.50 (M, 1H), 7.36 (M, 1H), 7.29 (M, 2H), 2.09 (S, 3H). LCMS: M/Z 442 (M+H)⁺.
- 30 [0309] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide (Cmpd 12): ¹H NMR (DMSO-*D*₆) 10.37 (S, 1H), 10.20 (S, 1H), 8.55 (S, 1H), 8.49 (D, 1H), 8.22 (D, 2H), 7.72 (D, 2H), 7.36 (D, 2H), 7.22 (M, 3H), 7.12 (D, 2H), 2.33 (S, 6H), 2.08 (S, 3H). LCMS: M/Z 452 (M+H)⁺.
- 35 [0310] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyridine-4-carboxamide (Cmpd 14): ¹H NMR (DMSO-*D*₆) 10.59 (S, 1H), 10.34 (S, 1H), 9.71 (S, 1H), 8.80 (DD, 2H), 8.50 (M, 2H), 8.21 (D, 2H), 7.90 (DD, 2H), 7.75 (D, 2H), 7.51 (M, 1H), 7.37 (D, 1H), 7.31 (M, 2H), 2.09 (S, 3H). LCMS: M/Z 425 (M+H)⁺.

5

[0311] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,3,4,5,6-pentafluorobenzamide (Cmpd 15): ¹H NMR (DMSO-*d*₆) 10.97 (s, 1H), 10.20 (s, 1H), 9.75 (s, 1H), 8.50 (d, 1H), 8.33 (s, 1H), 8.17 (d, 2H), 7.72 (d, 2H), 7.55 (m, 1H), 7.38 (d, 1H), 7.32 (t, 1H), 7.24 (d, 1H), 2.08 (s, 3H). LCMS: *m/z* 514 (M+H)⁺.

10 [0312] *N*-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide (Cmpd 1): ¹H NMR (DMSO-*d*₆) 10.15 (s, 1H), 8.68 (t, 1H), 8.27 (d, 1H), 8.04 (d, 2H), 7.67 (d, 2H), 7.49 (d, 2H), 7.41 (m, 1H), 7.13 (t, 1H), 7.06 (d, 1H), 3.41 (m, 2H), 3.30 (m, 2H), 2.06 (s, 3H), 1.80 (m, 2H). LCMS: *m/z* 458 (M+H)⁺.

[0313] 2,6-dichloro-*N*-(3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}propyl)benzamide (Cmpd 2): ¹H NMR (DMSO-*d*₆) 8.63 (s, 1H), 8.36 (d, 1H), 7.73 (s, 1H), 7.58 (d, 1H), 7.52 (d, 1H), 7.47 (m, 2H), 7.40 (m, 2H), 6.79 (d, 2H), 3.37 (m, 2H), 3.27 (m, 2H), 1.75 (t, 2H). LCMS: *m/z* 471 (M+H)⁺.

[0314] 4-(2,4-dichlorophenyl)-*N*-{3-[(2-piperidin-1-ylethyl)oxy]phenyl}pyrimidin-2-amine (Cmpd 19): ¹H NMR (DMSO-*d*₆) 9.79 (s, 1H), 8.58 (d, 1H), 7.77 (d, 1H), 7.68 (d, 1H), 7.58 (m, 2H), 7.24 (d, 1H), 7.10 (m, 2H), 6.50 (dd, 1H), 3.98 (t, 2H), 2.60 (t, 2H), 2.47 (m, 4H), 1.46 (m, 4H), 1.36 (m, 2H). LCMS: *m/z* 443 (M+H)⁺.

[0315] *N*-(3-{{4-(4-aminophenyl)pyrimidin-2-yl}amino}propyl)-2,6-dichlorobenzamide (Cmpd 6): ¹H NMR (400 MHz, DMSO): δ 8.76-8.79 (m, 1H), 8.45 (d, 6.0 Hz, 1H), 7.94-8.06 (m, 3H), 7.56 (m, 1H), 7.52-7.51 (m, 1H), 7.50 (s, 2H), 7.43-7.52 (m, 2H), 7.36-7.42 (m, 1H), 7.22-7.34 (bs, 1H), 3.44-3.64 (bs, 2H), 3.32-3.40 (m, 2H), 1.02-1.52 (bs, 2H). LC/MS MH=416.

[0316] 2,6-dichloro-*N*-(3-{{4-(2,3-dihydro-1-benzofuran-6-yl)pyrimidin-2-yl}amino}propyl)benzamide (Cmpd 4): ¹H NMR (400 MHz, DMSO): δ 8.724 (t, 5.6Hz, 1H), 8.29 (d, 5.6 Hz, 1H), 8.41-8.18 (bs, 1H), 7.92-8.15 (bs, 1H), 7.49-7.52 (m, 2H), 7.41-7.45 (m, 2H), 7.14-7.23 (bs, 1H), 6.88 (d, 8.4Hz, 1H), 4.61-4.65 (m, 2H), 3.62-3.93 (bs, 1H), 3.39-3.51 (bs, 1H), 3.33-3.36 (m, 2H), 3.23-3.27 (m, 2H), 1.80-1.87 (bs, 2H). LC/MS MH=443.

[0317] 2,6-dichloro-*N*-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-benzamide (Cmpd 3): ¹H NMR (400 MHz, DMSO): δ 8.66 (t, 5.6Hz, 1H), 8.17 (d, 5.2 Hz, 1H), 7.95 (d, 8.8 Hz, 2H), 7.47-7.49 (m, 2H), 7.38-7.42 (m, 1H), 6.95-6.97 (m, 2H), 6.72-6.75 (m, 2H), 3.37-3.43 (m, 2H), 3.26-3.32 (m, 2H), 2.96 (s, 6H), 1.79 (t, 6.8Hz, 2H). LC/MS M-H=442.

5 [0318] *N*-[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide (Cmpd 25): ¹H NMR (400MHz, DMSO-d₆) δ 10.67 (s, 1H), 10.15 (s, 1H), 9.57 (s, 1H), 8.38 (s, 1H), 8.16 (s, 1H), 7.72 (s, 4H), 7.60-7.50 (m, 4H), 7.26-7.23 (m, 2H), 2.26 (s, 3H), 2.09 (s, 3H). LCMS (EI) C₂₆H₂₂Cl₂N₅O₂: 506 (M+H).

[0319] *N*-[3-({4-[4-(acetylamino)phenyl]-5-fluoropyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide (Cmpd 28): ¹H NMR (400MHz, DMSO-d₆) δ 10.72 (s, 1H), 10.25 (s, 1H), 9.78 (s, 1H), 8.59 (d, J = 4.0Hz, 1H), 8.32 (s, 1H), 8.12 (d, J = 8.4Hz, 2H), 7.77 (d, J = 8.8Hz, 2H), 7.61-7.58 (m, 2H), 7.51 (dd, J = 8.8, 6.8Hz, 1H), 7.44 (d, J = 8.4Hz, 1H), 7.28 (t, J = 8.0Hz, 1H), 7.23 (t, J = 8.0Hz, 1H), 2.09 (s, 3H). LCMS (EI) C₂₅H₁₈Cl₂FN₅O₂: 510 (M+H).

15 [0320] *N*-(4-{2-[(4-{[2-(4-ethylpiperazin-1-yl)-2-oxoethyl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 64): ¹H-NMR (400MHz, d₆-DMSO): 10.222 ppm (s, 1H), 9.445 ppm (s, 1H), 8.457 ppm (d, 1H), 8.119 ppm (d, 2H), 7.75 ppm (d, 2H), 7.686 ppm (d, 2H), 7.299 ppm (d, 1H), 6.922 ppm (d, 2H), 4.76 ppm (s, 2H), 3.465 ppm (bs, 4H), 2.4 ppm (m, 4H), 2.31 ppm (m, 2H), 2.091 ppm (s, 3H), 1.02 ppm (t, 3H); MS (EI) C₂₆H₃₀N₆O₃: 475.4 (MH⁺).

[0321] 1-ethyl-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea (cmpd 250·HCl) (Cmpd 250): ¹H-NMR (400MHz, d₆-DMSO): 10.064 ppm (s, 1H), 9.305 ppm (s, 1H), 8.483 ppm (d, 1H), 8.105 ppm (d, 2H), 7.832 ppm (bd, 2H), 7.603 ppm (d, 2H), 7.54 ppm (bs, 2H), 7.431 ppm (d, 1H), 6.55 ppm (bs, 1H), 3.89 ppm (bs, 4H), 3.426 ppm (bs, 25 4H), 3.15 ppm (m, 2H), 1.08 ppm (t, 3H); MS (EI) C₂₃H₂₆N₆O₂HCl: 419.3 (MH⁺).

[0322] *N*-[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyrimidin-4-yl]-2,6-dichlorobenzamide (Cmpd 301): ¹H NMR (400MHz, d₆-DMSO): 11.55 (s, 1H), 10.5 (s, 1H), 10.2 (s, 1H), 9.32 (s, 1H), 8.68 (d, 1H), 8.6 (s, 1H), 8.4 (d, 2H), 7.74 (d, 2H), 7.64 (d, 1H), 7.62-7.5 (m, 3H), 2.03 (s, 3H); MS (EI) for C₂₃H₁₇Cl₂N₇O₂: 494 (MH⁺).

30 [0323] *N*-[4-(2-{[4-(morpholin-4-ylmethyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide (Cmpd 232): ¹H NMR (400MHz, d₆-DMSO): 10.3 (s, 1H), 9.6 (s, 1H), 8.46 (d, 1H), 8.13 (d, 2H), 7.78 (t, 4H), 7.35 (d, 1H), 7.47 (d, 1H), 7.23 (d, 2H), 3.58 (t, 4H), 3.4 (s, 2H), 2.33 (t, 4H), 2.1 (s, 3H); MS (EI) for C₂₃H₂₅N₅O₂: 404 (MH⁺).

[0324] 4-(1*H*-indol-5-yl)-*N*-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (Cmpd 254): 35 ¹H-NMR (400MHz, d₆-DMSO): 11.35 ppm (s, 1H), 9.33 ppm (s, 1H), 8.41 ppm (m, 2H), 7.94 ppm (dd, 1H), 7.71 ppm (d, 2H), 7.50 ppm (d, 1H), 7.44 ppm (t, 1H), 7.33 ppm (d, 1H),

5 6.94 ppm (d, 2H), 6.57 ppm (s, 1H), 3.75 ppm (m, 4H), 3.05 ppm (m, 4H), 2.09 ppm (s, 3H); MS (EI) C₂₄H₂₅N₅O₂: 372.3 (MH⁺).

[0325] *N*-(3-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 79): ¹H-NMR (400MHz, d₆-DMSO): 10.14 ppm (s, 1H), 9.45 ppm (s, 1H), 8.49 ppm (d, 1H), 8.40 ppm (s, 1H), 7.77 ppm (d, 1H), 7.70 ppm (d, 3H), 7.45 ppm (t, 1H), 7.20 ppm (d, 1H), 6.93 ppm (d, 2H), 3.74 ppm (m, 4H), 3.04 ppm (m, 4H), 2.09 ppm (s, 3H); MS (EI) C₂₄H₂₅N₅O₂: 390.1 (MH⁺).

[0326] 4-[4-(methyloxy)phenyl]-*N*-(4-morpholin-4-ylphenyl)pyrimidin-2-amine (Cmpd 252): ¹H-NMR (400MHz, d₆-DMSO): 9.37 ppm (s, 1H), 8.42 ppm (d, 1H), 8.13 ppm (d, 2H), 7.67 ppm (d, 2H), 7.27 ppm (d, 1H), 7.08 ppm (d, 2H), 6.92 ppm (d, 2H), 3.84 ppm (s, 3H), 3.74 ppm (m, 4H), 3.04 ppm (m, 4H); MS (EI) C₂₄H₂₅N₅O₂: 363.1 (MH⁺).

[0327] *N*-(4-{2-[(3-piperazin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 312): ¹H-NMR (400MHz, d₆-DMSO): 10.32 ppm (s, 1H), 9.55 ppm (s, 1H), 8.50 ppm (d, 1H), 8.13 ppm (d, 2H), 7.78 ppm (d, 2H), 7.68 ppm (s, 1H), 7.35 ppm (dd, 2H), 7.20 ppm (t, 1H), 6.63 (dd, 1H), 3.38 (m, 4H), 3.25 (m, 4H), 2.13 ppm (s, 3H); MS (EI) C₂₂H₂₄N₆O: 389.1 (MH⁺).

[0328] 2-amino-*N*-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-phenyl-acetamide (Cmpd 573): ¹H-NMR (400MHz, d₆-DMSO): 11.93 ppm (s, 1H), 10.18 (s, 1H), 9.08 ppm (s, 2H), 8.58 ppm (d, 1H), 8.18 ppm (d, 2H), 7.98-7.88 ppm (m, 2H), 7.82-7.75 ppm (m, 2H), 7.60-7.40 ppm (m, 6H), 7.30 ppm (s, 2H), 5.52-5.46 (m, 1H), 4.10 (t, 4H), 3.57 (t, 4H); MS (EI) C₂₈H₂₈N₆O₂: 481.3 (MH⁺).

[0329] *N*-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide (Cmpd 577): ¹H-NMR (400MHz, d₆-DMSO): 9.387 ppm (s, 1H), 8.443 ppm (d, 1H), 8.127 ppm (d, 2H), 7.825 ppm (d, 2H), 7.676 ppm (d, 2H), 7.287 ppm (d, 1H), 6.939 ppm (d, 2H), 3.747 ppm (m, 4H), 3.457 ppm (q, 1H), 3.050 ppm (m, 4H), 1.896 ppm (s, 3H (AcOH)), 1.243 ppm (d, 3H); MS (EI) C₂₃H₂₆N₆O₂: 419.1 (MH⁺).

[0330] *N*-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)valinamide (Cmpd 575): ¹H-NMR (400MHz, d₆-DMSO): 9.390 ppm (s, 1H), 8.449 ppm (d, 1H), 8.124 ppm (d, 2H), 7.815 ppm (d, 2H), 7.686 ppm (d, 2H), 7.284 ppm (d, 1H), 6.939 ppm (d, 2H), 3.747 ppm (m, 4H), 3.147 ppm (d, 1H), 3.051 ppm (m, 4H), 1.953 ppm (m, 1H), 1.864 ppm (s, 3H (AcOH)), 0.943 ppm (d, 3H), 0.871 ppm (d, 3H); MS (EI) C₂₅H₃₀N₆O₂: 446.2 (MH⁺).

5

[0331] 2-(dimethylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-acetamide (Cmpd 197): ¹H-NMR (400MHz, d₆-DMSO): 10.001 ppm (s, 1H), 9.384 ppm (s, 1H), 8.440 ppm (d, 1H), 8.118 ppm (d, 2H), 7.836 ppm (d, 2H), 7.673 ppm (d, 2H), 7.287 ppm (d, 1H), 6.939 ppm (d, 2H), 3.745 ppm (m, 4H), 3.114 ppm (s, 2H), 3.049 ppm (m, 4H), 2.290 ppm (s, 6H); MS (EI) C₂₄H₂₈N₆O₂: 432.5 (MH⁺).

10

[0332] N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)acetamide (Cmpd 578): ¹H-NMR (400MHz, d₆-DMSO): 10.208 ppm (s, 1H), 9.358 ppm (s, 1H), 8.433 ppm (d, 1H), 8.106 ppm (d, 2H), 7.741 ppm (d, 2H), 7.648 ppm (d, 2H), 7.262 ppm (d, 1H), 6.909 ppm (d, 2H), 3.050 ppm (m, 4H), 2.595 ppm (m, 4H), 2.212 ppm (s, 6H), 2.172 ppm (s, 2H), 2.091 ppm (m, 5H), 0.843 ppm (s, 6H); MS (EI) C₂₉H₃₉N₇O: 502.2 (MH⁺).

15

[0333] N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide (Cmpd 576): ¹H-NMR (400MHz, d₆-DMSO): 10.23 ppm (s, 1H), 9.37 ppm (s, 1H), 8.43 ppm (d, 1H), 8.106 ppm (d, 2H), 7.74 ppm (d, 2H), 7.65 ppm (d, 2H), 7.26 ppm (d, 1H), 7.10 ppm (s, 1H), 6.92 ppm (d, 2H), 6.78 ppm (s, 1H), 3.67 ppm (s, 3H), 3.58 ppm (s, 2H), 3.39-3.34 ppm (m, 4H), 3.02-3.08 ppm (M, 4H), 2.10 ppm (s, 3H); MS (EI) C₂₇H₃₀N₈O 482.6 (MH⁺).

20

[0334] N-(4-(2-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide (Cmpd 349): ¹H-NMR (400MHz, d₆-DMSO): 10.22 ppm (s, 1H), 9.41 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.74 ppm (d, 2H), 7.69 ppm (d, 2H), 7.28 ppm (d, 1H), 6.97 ppm (d, 2H), 3.83 ppm (s, 2H), 3.62 ppm (s, 2H), 3.15 ppm (s, 2H), 3.03 ppm (s, 2H), 2.09 ppm (s, 3H), 2.02-2.05 ppm (m, 1H), 0.74-0.76 ppm (m, 4H); MS (EI) C₂₆H₂₈N₆O₂: 456.5 (MH⁺).

25

[0335] N-{4-[2-({3-[(4-phenylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide (Cmpd 78): ¹H-NMR (400MHz, d₆-DMSO): 10.21 ppm (s, 1H), 9.83 ppm (s, 1H), 8.53 ppm (d, 1H), 8.13 ppm (d, 2H), 8.05 ppm (s, 1H), 7.95 ppm (s, 1H), 7.87 ppm (d, 1H), 7.75 ppm (d, 2H), 7.38-7.43 (m, 2H), 7.20-7.242 (m, 2H), 7.02 ppm (d, 2H), 6.95 ppm (d, 2H), 6.81 ppm (t, 1H), 3.68-3.88 ppm (m, 2H), 3.44-3.65 ppm (m, 2H), 3.02-3.11 ppm (m, 4H), 2.09 ppm (s, 3H); MS (EI) C₂₉H₂₈N₆O₂: 492.6 (MH⁺).

30

[0336] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide (Cmpd 578): ¹H-NMR (400MHz, d₆-DMSO): 11.13 ppm (s, 1H), 9.89 ppm (s, 1H), 8.53 ppm (d, 1H), 8.35 ppm (d, 3H), 8.20 ppm (d, 2H), 7.85 ppm (d, 4H), 7.49 ppm

35

5 (br s, 2H), 7.43 ppm (d, 1H), 4.14 ppm (m, 1H), 3.96 ppm (br s, 4H), 3.40 ppm (br s, 4H), 1.50 ppm (s, 3H); MS (EI) $C_{23}H_{26}N_6O_2$: 419 (MH⁺).

[0337] (N-(4-{5-methyl-2-[(3-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide): ¹H-NMR (400MHz, d₆-DMSO): 10.17 (s, 1H), 9.41 (s, 1H), 8.38 (s, 1H), 7.84 (s, 1H), 7.72 (s, 4H), 7.09 (d, 2H), 6.51 (dd, 1H), 3.74 (t, 4H), 3.07 (t, 4H), 2.26 (s, 3H), 2.09
10 (s, 3H); MS (EI) $C_{23}H_{25}N_5O_2$: 404.3 (M+H)⁺.

[0338] (N-(4-{6-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide): ¹H-NMR (400MHz, d₆-DMSO): 10.18 (s, 1H), 9.31 (s, 1H), 8.09 (d, 2H), 7.74-7.70 (m, 4H), 7.19 (s, 1H), 6.93 (d, 2H), 3.74 (t, 4H), 3.04 (t, 4H), 2.38 (s, 3H), 2.09 (s, 3H); MS (EI) $C_{23}H_{25}N_5O_2$: 404.3 (M+H)⁺.

15 [0339] (N-(4-{2-[(4-morpholin-4-ylphenyl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}phenyl)acetamide): ¹H-NMR (400MHz, CDCl₃): 8.66 (s, 1H), 7.65-7.59 (m, 4H), 7.51 (d, 2H), 7.33 (d, 2H), 6.93 (d, 2H), 3.87 (t, 4H), 3.14 (t, 4H); MS (EI) $C_{26}H_{30}N_6O_3$: 458.1 (M+H)⁺.

[0340] (N-(4-{2-[(3-aminophenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide):
20 ¹H-NMR (400MHz, d₆-DMSO): 10.14 (s, 1H), 9.19 (s, 1H), 8.33 (d, 1H), 7.72 (d, 2H), 7.67 (dd, 2H), 7.03 (t, 1H), 6.94 (dd, 1H), 6.87 (t, 1H), 6.17-6.14 (m, 1H), 4.92 (s, 2H), 2.23 (s, 3H), 2.08 (s, 3H); MS (EI) $C_{19}H_{19}N_5O$: 334.0 (M+H)⁺.

[0341] (N-(4-{2-[(3-aminophenyl)amino]-5-fluoropyrimidin-4-yl}phenyl)acetamide):
25 ¹H-NMR (400MHz, d₆-DMSO): 10.26 (s, 1H), 9.41 (s, 1H), 8.54 (d, 1H), 8.04 (d, 2H), 7.78 (d, 2H), 7.03 (s, 1H), 6.95-6.91 (m, 2H), 6.20 (d, 1H), 5.00 (s, 2H), 2.10 (s, 3H); MS (EI) $C_{18}H_{16}FN_5O$: 338.3 (M+H)⁺.

[0342] (N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-fluoropyrimidin-4-yl)phenyl]acetamide): ¹H-NMR (400MHz, d₆-DMSO): 10.26 (s, 1H), 9.45 (s, 1H), 8.52(d, 1H), 8.02 (d, 2H), 7.78 (d, 2H), 7.59 (d, 2H), 6.91 (d, 2H), 3.35 (bs, 4H), 3.07 (bs, 4H), 2.50
30 (q, 2H), 2.10 (s, 3H), 1.04 (t, 3H); MS (EI) $C_{24}H_{27}FN_6O$: 435.3 (M+H)⁺.

[0343] (N-[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide): ¹H-NMR (400MHz, d₆-DMSO): 10.32 (s, 1H), 10.15 (s, 1H), 9.51 (s, 1H), 8.37(s, 1H), 8.29 (s, 1H), 7.76-7.70 (m, 4H), 7.42-7.41 (m, 1H), 7.25-7.17 (m, 3H), 7.11 (d, 2H), 2.29 (s, 6H), 2.26 (s, 3H), 2.09(s, 3H); MS (EI) $C_{28}H_{27}N_5O_2$: 466.3 (M+H)⁺.

5

[0344] (N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-fluoropyrimidin-4-yl}phenyl)-acetamide): ¹H-NMR (400MHz, d₆-DMSO): 10.27 (s, 1H), 9.45 (s, 1H), 8.58(d, 1H), 8.07 (d, 2H), 7.77 (d, 2H), 7.06 (s, 2H), 6.17 (s, 1H), 3.75 (t, 8H), 3.10 (t, 8H), 2.10 (s, 3H); MS (EI) C₂₆H₂₉FN₆O₃: 493.4 (M+H)⁺.

10 **[0345] (N-{4-[2-(1H-indazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide):** ¹H-NMR (400MHz, d₆-DMSO): 12.80 (s, 1H), 10.18 (s, 1H), 9.72 (s, 1H), 8.44(d, 1H), 8.37 (d, 1H), 7.90 (s, 1H), 7.77-7.70 (m, 4H), 7.59 (d, 1H), 7.28 (dd, 1H), 2.27 (s, 3H), 2.10 (s, 3H); MS (EI) C₂₀H₁₈N₆O: 359.3 (M+H)⁺.

[0346] (N-{4-[2-(1H-indol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide):
15 ¹H-NMR (400MHz, d₆-DMSO): 10.90 (s, 1H), 10.15 (s, 1H), 9.21 (s, 1H), 8.32(d, 1H), 8.00 (d, 1H), 7.72 (dd, 2H), 7.66 (dd, 2H), 7.36 (dd, 1H), 7.28-7.25 (m, 2H), 6.33 (t, 1H), 2.22 (s, 3H), 2.09 (s, 3H); MS (EI) C₂₁H₁₉N₅O: 358.3 (M+H)⁺.

[0347] (N-(4-{5-fluoro-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide): ¹H-NMR (400MHz, d₆-DMSO): 10.26 (s, 1H), 9.48 (s, 1H), 8.52 (d, 1H), 8.02
20 (d, 2H), 7.77 (d, 2H), 7.61 (d, 2H), 6.93 (d, 2H), 3.74 (t, 4H), 3.03 (t, 4H); MS (EI) C₂₂H₂₂FN₅O₂: 408.3 (M+H)⁺.

[0348] N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclopropanecarboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.41 (s, 1H), 9.28 (s, 1H),
25 8.31(d, 1H), 7.73 (d, 2H), 7.66-7.62 (m, 4H), 6.89 (d, 2H), 3.73 (bs, 4H), 3.02 (bs, 4H), 2.22 (s, 3H), 1.85-1.79 (m, 1H), 0.84-0.81 (m, 4H); MS (EI) C₂₅H₂₇N₅O₂: 430 (MH⁺).

[0349] N-{4-[2-(1H-indazol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide: ¹H-NMR (400MHz, d₆-DMSO): 12.88 (s, 1H), 10.16 (s, 1H), 9.49 (s, 1H), 8.37(s, 1H), 8.29 (d, 1H), 7.97 (s, 1H), 7.73 (d, 2H), 7.67 (d, 2H), 7.59 (dd, 1H), 7.44 (d, 1H), 2.24 (s, 3H), 2.10 (s, 3H); MS (EI) C₂₀H₁₈N₆O: 359 (MH⁺).

30 **[0350] N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.48 (d, 1H), 8.15 (d, 2H), 7.74 (d, 2H), 7.32 (d, 1H), 7.12 (d, 2H), 6.17 (s, 1H), 3.75 (t, 4H), 3.11 (t, 4H), 2.09 (s, 3H); MS (EI) C₂₆H₃₀N₆O₃: 475 (MH⁺).

[0351] 4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzonitrile: ¹H-NMR
35 (400MHz, d₆-DMSO): 9.60 (s, 1H), 8.57 (d, 1H), 8.32 (d, 2H), 8.03 (d, 2H), 7.65 (d, 2H), 7.44 (d, 1H), 6.94 (d, 2H), 3.75 (t, 4H), 3.05 (t, 4H), ; MS (EI) C₂₁H₁₉N₅O: 358 (MH⁺).

- 5 **[0352] 4-(4-fluorophenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400MHz, d₆-DMSO): 9.46 (s, 1H), 8.49 (d, 1H), 8.22 (dd, 2H), 7.66 (d, 2H), 7.38 (t, 2H), 7.33 (d, 1H), 6.93 (dd, 2H), 3.74 (t, 4H), 3.05 (t, 4H), ; MS (EI) C₂₀H₁₉FN₄O: 358.3 (M+H)⁺.
- [0353] N-(4-morpholin-4-ylphenyl)-4-(4-pyrimidin-5-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400MHz, d₆-DMSO): 9.50 (s, 1H), 9.26 (s, 2H), 9.24 (s, 1H), 8.53 (d, 1H), 8.32 (d, 10 2H), 8.02 (d, 2H), 7.69 (d, 2H), 7.44 (s, 1H), 6.94 (d, 2H), 3.75 (t, 4H), 3.06 (t, 4H); MS (EI) C₂₄H₂₂N₆O: 411 (MH⁺).
- [0354] N-(4-morpholin-4-ylphenyl)-4-[4-(pyridin-2-ylamino)phenyl]pyrimidin-2-amine:** ¹H-NMR (400MHz, d₆-DMSO): 9.42 (s, 1H), 9.33 (s, 1H), 8.40 (d, 1H), 8.23 (dd, 1H), 8.09 (dd, 2H), 7.86 (d, 2H), 7.70 (d, 2H), 7.65-7.61 (m, 1H), 7.25 (d, 1H), 6.96-6.90 (m, 15 3H), 6.84-6.81 (m, 1H), 3.75 (t, 4H), 3.05 (t, 4H); MS (EI) C₂₅H₂₄N₆O: 425 (MH⁺).
- [0355] N-(4-morpholin-4-ylphenyl)-4-[4-(pyridin-3-ylamino)phenyl]pyrimidin-2-amine:** ¹H-NMR (400MHz, d₆-DMSO): 9.33 (s, 1H), 8.81 (s, 1H), 8.45 (d, 1H), 8.40 (d, 1H), 8.14 (dd, 1H), 8.08 (d, 2H), 7.68 (d, 2H), 7.63-7.60 (m, 1H), 7.32 (dd, 1H), 7.23 (d, 1H), 7.19 (dd, 2H), 6.93 (d, 2H), 3.74 (t, 4H), 3.05 (t, 4H); MS (EI) C₂₅H₂₄N₆O: 425 (MH⁺).
- 20 **[0356] N-[4-(2-([4-(4-D-alanylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.31 (s, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.69 (d, 2H), 7.30 (d, 1H), 6.97 (d, 2H), 4.09-4.04 (m, 1H), 3.77-3.73 (m, 1H), 3.67-3.61 (m, 4H), 3.09-3.03 (m, 4H), 2.92 (t, 2H), 2.10-2.03 (m, 1H), 1.85-1.77 (m, 1H), 1.71-1.64 (m, 2H), 1.19 (d, 3H); MS (EI) C₂₈H₃₄N₈O₂: 515 (MH⁺).
- 25 **[0357] N-[4-(2-([4-(4-L-alanylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.34 (s, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.69 (d, 2H), 7.30 (d, 1H), 6.97 (d, 2H), 4.06-4.01 (m, 1H), 3.77-3.73 (m, 1H), 3.67-3.61 (m, 4H), 3.09-3.02 (m, 4H), 2.92 (t, 2H), 2.10-2.03 (m, 1H), 1.85-1.77 (m, 1H), 1.71-1.64 (m, 2H), 1.18 (d, 3H); MS (EI) C₂₈H₃₄N₈O₂: 515 (MH⁺).
- 30 **[0358] N-{4-[2-([4-(4-(piperazin-1-ylacetyl)piperazin-1-yl)phenyl]amino)pyrimidin-4-yl]phenyl}-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.28 (s, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 3.76-3.73 (m, 1H), 3.69-3.59 (m, 4H), 3.19 (s, 2H), 3.10-3.02 (m, 4H), 2.91 (t, 2H), 2.81 (bs, 4H), 2.44 (bs, 4H), 2.10-2.04 (m, 1H), 1.85-1.77 (m, 1H), 1.71-1.64 (m, 2H); MS (EI)
- 35 C₃₁H₃₉N₉O₂: 570 (MH⁺).

5

[0359] N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide: ¹H-NMR (400MHz, d₆-DMSO): 9.43 (s, 1H), 8.47 (d, 1H), 8.17 (d, 2H), 7.80 (d, 2H), 7.69 (d, 2H), 7.31 (d, 1H), 6.98 (d, 2H), 3.82-3.78 (m, 4H), 3.66-3.62 (m, 4H), 3.09-3.03 (m, 4H), 2.97-2.90 (m, 1H), 1.39 (d, 3H), 1.03 (d, 6H); MS (EI)

10 C₂₇H₃₃N₇O₂: 488 (MH⁺).

[0360] N-[4-(2-{[4-(4-D-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-alaninamide: ¹H-NMR (400MHz, d₆-DMSO): 9.42 (s, 1H), 8.45 (d, 1H), 8.33 (s, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.69 (d, 2H), 7.30 (d, 1H), 6.98 (d, 2H), 4.13-4.07 (m, 1H), 3.70-3.59 (m, 5H), 3.11-3.01 (m, 4H), 1.27 (d, 3H), 1.21 (d, 3H); MS (EI) C₂₆H₃₂N₈O₂: 489

15 (MH⁺).

[0361] N-{4-[2-({4-[4-(tetrahydrofuran-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.19 (s, 1H), 9.42 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 3.89 (t, 1H), 3.74-3.69 (m, 4H), 3.67-3.63 (m, 4H), 3.45-3.37 (m, 1H), 3.09-3.02 (m, 4H), 2.90 (t, 2H), 2.08-1.99 (m, 3H), 1.84-1.76 (m, 1H), 1.70-1.64 (m, 2H); MS (EI) C₃₀H₃₅N₇O₃: 542 (MH⁺).

20

[0362] N-{4-[2-({4-[4-(tetrahydrofuran-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 4.72 (dd, 1H), 3.82-3.72 (m, 3H), 3.69-3.58 (m, 4H), 3.08-3.02 (m, 4H), 2.91 (t, 2H), 2.11-1.99 (m, 3H), 1.88-1.79 (m, 3H), 1.71-1.66 (m, 2H); MS (EI) C₃₀H₃₅N₇O₃: 542 (MH⁺).

25

[0363] N-(4-{5-chloro-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.30 (s, 1H), 9.67 (s, 1H), 8.52 (s, 1H), 7.82 (s, 4H), 7.59 (d, 2H), 6.90 (dd, 2H), 3.84 (dd, 1H), 3.74-3.72 (m, 4H), 3.04-3.02 (m, 4H), 2.97 (t, 2H), 2.15-2.09 (m, 1H), 1.86-1.79 (m, 1H), 1.75-1.69 (m, 2H); MS (EI) C₂₅H₂₇ClN₆O₂: 479 (MH⁺).

30

[0364] (R)-N-(4-(2-(4-(4-(2-(pyrrolidin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 3.75-3.71 (m, 1H), 3.69-3.59 (m, 4H), 3.07-3.01 (m, 4H), 2.91 (t, 2H), 2.50 (t, 4H), 2.48 (s, 2H), 2.11-2.02 (m, 1H), 1.86-1.78 (m, 1H), 1.72-1.63 (m, 6H); MS (EI) C₃₁H₃₈N₈O₂: 555 (MH⁺).

5

- 5 **[0365] (R)-N-(4-(2-(4-(4-(2-morpholinoacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.19 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.84 (dd, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 3.74-3.69 (m, 3H), 3.61-3.57 (m, 6H), 3.19 (s, 2H), 3.11-3.09 (m, 2H), 3.04-3.01 (m, 2H), 2.90 (t, 2H), 2.41 (bs, 4H), 2.11-2.02 (m, 1H), 1.84-1.76 (m, 1H), 1.70-1.63 (m, 2H); MS (EI) C₃₁H₃₈N₈O₃: 571 (MH⁺).
- 10 **[0366] N-{4-[2-({4-[2-(methyloxy)ethyl]-3,4-dihydro-2H-1,4-benzoxazin-7-yl}amino)-pyrimidin-4-yl]phenyl}-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.19 (s, 1H), 9.25 (s, 1H), 8.42 (d, 1H), 8.11 (d, 2H), 7.83 (d, 2H), 7.28-7.25 (m, 2H), 7.17 (dd, 1H), 6.68 (d, 1H), 4.15 (t, 2H), 3.72 (dd, 1H), 3.52 (t, 2H), 3.41-3.36 (m, 4H), 3.27 (s, 3H), 2.90 (t, 2H), 2.10-2.01 (m, 1H), 1.84-1.75 (m, 1H), 1.70-1.63 (m, 2H); MS (EI) C₂₆H₃₀N₆O₃: 475 (MH⁺).
- 15 **[0367] N-(4-{2-[(4-{4-[(2R)-tetrahydrofuran-2-ylcarbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 4.72 (dd, 1H), 3.82-3.72 (m, 3H), 3.69-3.58 (m, 4H), 3.08-3.03 (m, 4H), 2.90 (t, 2H), 2.11-1.98 (m, 3H), 1.88-1.75 (m, 3H), 1.70-1.65 (m, 2H); MS (EI) C₃₀H₃₅N₇O₃: 542 (MH⁺).
- 20 **[0368] N-(4-{2-[(4-{4-[(2S)-tetrahydrofuran-2-ylcarbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 4.72 (dd, 1H), 3.82-3.72 (m, 3H), 3.69-3.58 (m, 4H), 3.08-3.03 (m, 4H), 2.91 (t, 2H), 2.09-1.98 (m, 3H), 1.88-1.75 (m, 3H), 1.70-1.63 (m, 2H); MS (EI) C₃₀H₃₅N₇O₃: 542 (MH⁺).
- 25 **[0369] N-{4-[2-(1,2,3,4-tetrahydroquinolin-6-ylamino)pyrimidin-4-yl]phenyl}-D-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.17 (s, 1H), 9.05 (s, 1H), 8.38 (d, 1H), 8.10 (dd, 2H), 7.84-7.81 (m, 2H), 7.29 (s, 1H), 7.21-7.18 (m, 2H), 6.40 (d, 1H), 5.36 (s, 1H), 3.72 (dd, 1H), 3.15 (t, 2H), 2.90 (t, 2H), 2.67 (t, 2H), 2.10-2.01 (m, 1H), 1.84-1.76 (m, 3H), 1.69-1.63 (m, 2H); MS (EI) C₂₄H₂₆N₆O: 415 (MH⁺).
- 30 **[0370] N-{4-[2-({4-[(phenylmethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.42 (s, 1H), 8.42 (d, 1H), 8.06 (d, 2H), 7.75-7.65 (m, 4H), 7.42-7.24 (m, 6H), 6.95 (d, 2H), 5.04 (s, 2H), 2.04 (s, 3H). MS (EI) for C₂₅H₂₂N₄O₂: 411 (MH⁺).
- 35

- 5 [0371] **4-(4-aminophenyl)-N-[4-(phenyloxy)phenyl]pyrimidin-2-amine:** ¹H-NMR (400 MHz, d6-DMSO): 10.42 (s, 1H), 8.43 (d, 1H), 8.08 (d, 2H), 7.72 (d, 2H), 7.43-7.37 (m, 3H), 7.26-7.15 (m, 3H), 7.07-6.95 (m, 3H). MS (EI) for C₂₂H₁₈N₄O: 355 (MH⁺).
- [0372] **N-[4-(2-{{4-(phenyloxy)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.64 (s, 1H), 8.49 (d, 1H), 8.11 (d, 2H), 7.85 (d, 2H), 7.75 (d, 2H), 7.38-7.32 (m, 3H), 7.10-7.03 (m, 3H), 6.97 (d, 1H). MS (EI) for C₂₄H₂₀N₄O₂: 397 (MH⁺).
- 10 [0373] **N-(4-{2-[[4-{{2-(methyloxy)ethyl}amino}phenyl]amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.18 (s, 1H), 9.08 (s, 1H), 8.38 (d, 1H), 8.05 (d, 2H), 7.73 (d, 2H), 7.43 (d, 2H), 7.19 (d, 1H), 6.58 (d, 2H), 5.20 (t, 1H), 3.43 (t, 2H), 3.25 (s, 3H), 3.16 (t, 2H), 2.04 (s, 3H). MS (EI) for C₂₁H₂₃N₅O₂: 378 (MH⁺).
- 15 [0374] **N-{4-[2-({4-[(pyridin-4-ylmethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.43 (s, 1H), 8.60 (s, 2H), 8.43 (d, 1H), 8.10 (d, 2H), 7.79-7.72 (m, 4H), 7.43 (d, 1H), 7.30 (d, 1H), 7.01 (d, 2H), 5.08 (s, 1H), 2.03 (s, 3H). MS (EI) for C₂₄H₂₁N₅O₂: 412 (MH⁺).
- 20 [0375] **N-(4-{2-[(4-cyclohexylphenyl)amino]pyrimidin-4-yl}phenyl) acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.53 (s, 1H), 8.45 (d, 1H), 8.15 (d, 2H), 7.78-7.70 (m, 4H), 7.35 (d, 1H), 7.18 (d, 1H), 2.43-2.40 (m, 1H), 2.08 (s, 3H), 1.82-1.68 (m, 4H), 1.42-1.20 (m, 6H). MS (EI) for C₂₄H₂₆N₄O: 387 (MH⁺).
- [0376] **N-{4-[2-({4-[(tetrahydrofuran-2-ylmethyl)amino]phenyl}amino) pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.18 (s, 1H), 8.39 (d, 1H), 8.08 (d, 2H), 7.76 (d, 2H), 7.45 (d, 2H), 7.20 (d, 1H), 6.60 (d, 2H), 5.23 (t, 1H), 4.02-3.97 (m, 1H), 3.80-3.77 (m, 1H), 3.65-3.60 (m, 1H), 3.08-3.00 (m, 2H), 2.07 (s, 3H), 2.00-1.80 (m, 3H), 1.62-1.57 (m, 1H). MS (EI) for C₂₃H₂₅N₅O₂: 404 (MH⁺).
- 25 [0377] **N-{4-[2-({4-[(phenylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.17 (s, 1H), 8.05 (d, 2H), 7.72 (d, 2H), 7.44-7.27 (m, 5H), 7.22-7.17 (m, 2H), 6.57 (d, 2H), 6.00 (t, 1H), 4.25 (d, 2H), 2.08 (s, 3H). MS (EI) for C₂₅H₂₃N₅O: 410 (MH⁺).
- 30 [0378] **ethyl [4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl] acetate:** ¹H-NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.53 (s, 1H), 8.45 (d, 1H), 8.15 (d, 2H), 7.78-7.70 (m, 4H), 7.32 (d, 1H), 7.17 (d, 2H), 4.05 (q, 2H), 3.45 (s, 2H), 1.35 (t, 3H). MS (EI) for C₂₂H₂₂N₄O₃: 391 (MH⁺).
- 35

- 5 **[0379] N-(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.25 (s, 1H), 9.73 (s, 1H), 8.55 (d, 1H), 8.12 (d, 2H), 8.05 (s, 1H), 7.80-7.70 (m, 3H), 7.37 (d, 2H), 7.20 (d, 2H), 3.75 (t, 4H), 2.95 (t, 4H), 2.05 (s, 3H). MS (EI) for C₂₂H₂₂ClN₅O₂: 425 (MH⁺).
- [0380] N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]-L-serinamide:** ¹H NMR (400 MHz, d₆-DMSO): 9.50 (s, 1H), 8.45 (d, 1H), 8.18 (d, 2H), 7.83 (d, 2H), 7.70 (s, 1H), 7.37-7.30 (m, 2H), 6.90 (d, 1H), 3.82 (s, 3H), 3.72 (t, 4H), 3.60-3.57 (m, 2H), 3.43 (t, 1H), 2.92 (t, 4H). MS (EI) : 465 (MH⁺).
- [0381] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-(1H-tetrazol-1-yl)acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.90 (s, 1H), 9.47 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.18 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 5.58 (s, 2H), 3.77 (t, 4H), 3.03 (t, 4H). MS (EI): 458 (MH⁺).
- 15 **[0382] (3S)-3-hydroxy-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]butanamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.19 (s, 1H), 9.48 (s, 1H), 8.48 (s, 1H), 8.17 (d, 2H), 7.78 (d, 2H), 7.65 (s, 1H), 7.31 (d, 2H), 6.88 (d, 1H), 4.80 (s, 1H), 4.15-4.07 (m, 1H), 3.81 (s, 3H), 3.75 (t, 4H), 2.96 (t, 4H), 2.44-2.37 (m, 2H), 1.14 (d, 3H). MS (EI): 464 (MH⁺).
- 20 **[0383] (3R)-3-hydroxy-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]butanamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.19 (s, 1H), 9.48 (s, 1H), 8.48 (s, 1H), 8.17 (d, 2H), 7.78 (d, 2H), 7.65 (s, 1H), 7.31 (d, 2H), 6.88 (d, 1H), 4.80 (s, 1H), 4.15-4.07 (m, 1H), 3.83 (s, 3H), 3.75 (t, 4H), 2.96 (t, 4H), 2.44-2.37 (m, 2H), 1.14 (d, 3H). MS (EI): 464 (MH⁺).
- 25 **[0384] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2,5-dihydro-1H-pyrrole-2-carboxamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.19 (s, 1H), 9.40 (s, 1H), 8.43 (s, 1H), 8.17 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.30 (s, 1H), 6.96 (d, 2H), 6.02-5.98 (m, 1H), 5.93-5.89 (m, 1H), 4.60 (s, 1H), 3.82 (s, 2H), 3.75 (t, 4H), 3.05 (t, 4H). MS (EI): 443 (MH⁺).
- 30 **[0385] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-[(2S)-pyrrolidin-2-yl]acetamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.86 (s, 1H), 10.10 (s, 1H), 9.37 (s, br, 1H), 9.28 (s, br, 1H), 8.55 (d, 1H), 8.20 (d, 2H), 7.95-7.85 (m, 4H), 7.70 (s, 2H), 7.45 (d, 1H), 4.10 (t, 4H), 3.83-3.78 (m, 2H), 3.73 (t, 4H), 3.25-3.18 (m, 1H), 3.03-2.95 (m, 2H), 2.20-2.10 (m, 1H), 2.00-1.80 (m, 2H), 1.68-1.56 (m, 1H). MS (EI): 459 (MH⁺).
- 35

- 5 **[0386] 2,3-dihydroxy-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)propanamide:** ¹H-NMR (400 MHz, d₆-DMSO): 9.95 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.25 (s, 1H), 6.96 (d, 2H), 5.95 (s, br, 1H), 4.95 (s, br, 1H), 4.08 (t, 1H), 3.78-3.60 (m, 6H), 3.03 (t, 4H). MS (EI) : 436 (MH⁺).
- [0387] 1-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}-3-ethylurea:** ¹H-NMR (400 MHz, d₆-DMSO): 9.37 (s, 1H), 9.19 (s, 1H), 9.40 (d, 1H), 8.03 (d, 2H), 7.70 (d, 2H), 7.58 (d, 2H), 7.23 (d, 2H), 6.95 (d, 2H), 6.75 (t, 1H), 3.58 (t, 4H), 3.60 (t, 3H), 3.15-3.00 (m, 8H), 2.18 (s, 6H), 1.05 (t, 3H). MS (EI) : 503 (MH⁺).
- [0388] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.78 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.95 (d, 2H), 3.73-3.58 (m, 6H), 3.24 (s, 3H), 3.14-3.00 (m, 6H), 2.60 (t, 3H), 2.20 (s, 6H). MS (EI): 518 (MH⁺).
- [0389] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}cyclopropanecarboxamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.26 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.14 (d, 2H), 7.78 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.95 (d, 2H), 3.70-3.58 (m, 4H), 3.10-3.00 (m, 6H), 2.20 (s, 6H), 1.84-1.80 (m, 1H), 0.83-0.80 (m, 4H). MS (EI): 500 (MH⁺).
- [0390] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}butanamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.12-8.05 (m, 3H), 7.80-7.68 (m, 3H), 7.28 (d, 1H), 6.99 (d, 2H), 2H, 3.70-3.60 (m, 4H), 3.28 (s, 2H), 3.14-3.00 (m, 4H), 2.35-2.20 (m, 8H), 1.64-1.58 (m, 2H), 0.95-0.88 (m, 3H). MS (EI): 502 (MH⁺).
- [0391] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycinamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.00 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.82 (d, 2H), 7.65 (d, 2H), 7.25 (d, 1H), 6.95 (d, 2H), 3.65-3.57 (m, 4H), 3.23 (s, 2H), 3.12-3.00 (m, 6H), 2.28 (s, 6H), 2.20 (s, 6H). MS (EI): 517 (MH⁺).
- [0392] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino) pyrimidin-4-yl]phenyl}-D-alaninamide:** ¹H-NMR (400 MHz, d₆-DMSO): 9.40 (s, 1H), 8.43 (d, 1H), 8.15 (d, 2H), 7.85 (d, 2H), 7.70 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 3.70-3.57 (m, 4H), 3.50 (q, 1H), 3.18-3.00 (m, 6H), 1.83 (s, 6H), 1.21 (d, 3H). MS (EI): 503 (MH⁺).

- 5 **[0393]** N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide: ¹H-NMR (400 MHz, d6-DMSO): 10.30 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.15 (d, 2H), 7.78 (d, 2H), 7.68 (d, 2H), 7.25 (d, 1H), 6.95 (d, 2H), 3.97 (t, 1H), 3.82-3.70 (m, 3H), 3.67-3.60 (m, 4H), 3.22-3.17 (m, 1 H), 3.12-3.00 (m, 4H), 2.35 (s, 6H), 2.12-2.05 (m, 2H). MS (EI): 530 (MH+).
- 10 **[0394]** (2R)-N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide: ¹H-NMR (400 MHz, d6-DMSO): 9.95 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.16 (d, 2H), 7.88 (d, 2H), 7.70 (d, 2H), 7.30 (d, 1H), 6.97 (d, 2H), 4.44-4.41 (m, 1H), 4.02-3.98 (m, 1H), 3.85-3.80 (m, 1H), 3.68-3.57 (m, 4H), 3.18-3.00 (m, 6H), 2.20 (s, 6H), 2.05-1.85 (m, 4H). MS (EI): 530 (MH+).
- 15 **[0395]** (2S)-N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide: ¹H-NMR (400 MHz, d6-DMSO): 9.95 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.16 (d, 2H), 7.88 (d, 2H), 7.70 (d, 2H), 7.30 (d, 1H), 6.98 (d, 2H), 4.45-4.42 (m, 1H), 4.02-3.98 (m, 1H), 3.85-3.80 (m, 1H), 3.70-3.57 (m, 4H), 3.20 (s, 2H), 3.10-3.00 (m, 6H), 2.22 (s, 6H), 2.05-1.85 (m, 4H). MS (EI):
- 20 530 (MH+).
- [0396]** N-(4-{2-[(6-morpholin-4-ylpyridin-3-yl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 11.38 (s, 1H), 10.10 (s, 2H), 9.03 (s, br, 1H), 8.75 (s, br, 1H), 8.60 (d, 2H), 8.30-8.20 (m, 3H), 7.85 (d, 2H), 7.55 (d, 2H), 4.48-4.42 (m, 1H), 3.82-3.70 (m, 8H), 3.37-3.20 (m, 2H), 2.43-2.40 (m, 1H), 2.10-1.95 (m, 3H). MS (EI):
- 25 446 (MH+).
- [0397]** 1-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclopentanecarboxamide: ¹H-NMR (400 MHz, d6-DMSO): 9.95 (s, 1H), 9.45 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.95 (d, 2H), 7.70 (d, 2H), 7.30 (d, 1H), 7.05-6.95 (m, 2H), 5.68 (s, br, 1H), 3.80-3.70 (m, 4H), 3.15-3.05 (m, 4H), 2.10-1.97 (m, 3H), 1.87-1.68 (m, 5H). MS
- 30 (EI) : 460 (MH+).
- [0398]** 2-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide: ¹H NMR (400 MHz, d6-DMSO): 9.95 (s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.70 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 5.75 (t, 1H), 4.03 (d, 2H), 3.78-3.70 (m, 4H), 3.10-3.00 (m, 4H). MS (EI): 406 (MH+).
- 35 **[0399]** 3-chloro-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 11.68 (s, 1H), 11.00

5 (s, 1H), 9.43 (s, 2H), 8.83 (s, 1H), 8.70 (d, 1H), 8.50 (d, 1H), 8.20 (d, 2H), 7.87 (d, 2H), 7.75-7.65 (m, 3H), 7.32 (d, 1H), 6.95 (d, 2H), 3.75 (t, 4H), 3.05 (t, 4H). MS (EI): 487 (MH⁺).

[0400] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 10.24 (s, br, 1H), 9.41 (s, 1H),

8.44 (d, 2H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.95 (d, 2H), 3.78-3.57 (m, 5H), 3.15-3.00 (m, 6H), 2.93 (t, 2H), 2.18 (s, 6H), 2.08-2.00 (m, 1H), 1.93-1.88 (m, 1H), 1.90-1.80 (m, 2H). MS (EI): 529 (MH⁺).

[0401] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide: ¹H-NMR (400 MHz, d₆-DMSO): 10.18 (s, br, 1H), 9.18 (s, 1H),

8.40 (d, 1H), 8.13 (d, 2H), 7.78 (d, 2H), 7.63 (d, 2H), 7.25 (d, 1H), 6.93 (d, 2H), 3.77 (t, 4H), 3.07 (t, 4H), 2.16 (q, 2H), 1.10 (t, 3H). MS (EI): 404 (MH⁺).

[0402] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-3-ylacetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.90 (s, 1H), 9.40 (s, 1H), 8.43 (s, 1H), 8.18 (d, 2H), 7.88 (d, 2H), 7.73-7.45 (m, 5H), 7.32 (d, 1H), 6.95 (d, 2H), 3.77 (t, 4H), 3.03 (t, 4H). MS (EI): 467 (MH⁺).

20 [0403] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrimidine-5-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.82 (s, 1H), 9.42 (s, 1H), 9.38 (s, 1H), 9.31 (s, 2H), 8.45 (d, 1H), 8.20 (d, 2H), 7.97 (d, 2H), 7.70 (d, 2H), 7.33 (d, 1H), 6.95 (d, 2H), 3.77 (t, 4H), 3.03 (t, 4H). MS (EI): 454 (MH⁺).

[0404] N-[4-(2-{[3-(morpholin-4-ylmethyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400MHz, d₆-DMSO): 10.25 (s, 1H), 9.62 (s, 1H), 8.5 (d,

25 1H), 8.16 (d, 2H), 7.94 (s, 1H), 7.76 (d, 2H), 7.62 (d, 1H), 7.36 (d, 1H), 7.26 (t, 1H), 6.9 (d, 1H), 3.6 (t, 4H), 3.45 (s, 2H), 2.39 (t, 4H), 2.1 (s, 3H), 1.86 (s, 3H). MS (EI) for C₂₃H₂₅N₅O₂: 404 (MH⁺).

[0405] N-[4-(2-{[3-(1,3-dioxan-2-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide:

30 ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.68 (s, 1H), 8.5 (d, 1H), 8.32 (s, 1H), 8.21 (d, 2H), 7.77 (d, 2H), 7.58 (d, 1H), 7.39 (d, 1H), 7.28 (t, 1H), 6.98 (d, 1H), 5.52 (s, 1H), 4.2 (dd, 2H), 4.0 (t, 2H), 2.1 (s, 3H), 2.05 (m, 1H), 1.5 (dd, 1H). MS (EI) for C₂₂H₂₂N₄O₃: 391 (MH⁺).

[0406] N-(4-{2-[(6-aminopyridin-2-yl)amino]pyrimidin-4-yl}phenyl)acetamide:

35 ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 8.93 (s, 1H), 8.55 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.57 (d, 1H), 7.45-7.4 (m, 2H), 6.12 (d, 1H), 5.78 (s, 2H), 2.08 (s, 3H). MS (EI) for C₁₇H₁₆N₆O: 321 (MH⁺).

- 5 **[0407] N-(4-{2-[(6-aminopyrimidin-4-yl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.25 (s, 1H), 9.72 (s, 1H), 8.6 (d, 1H), 8.2-8.15 (m, 3H), 7.8 (d, 2H), 7.52 (d, 1H), 7.42 (s, 1H), 6.8 (br, 2H), 2.08 (s, 3H). MS (EI) for C₁₆H₁₅N₇O : 322 (MH⁺).
- [0408] N-(4-morpholin-4-ylphenyl)-4-quinolin-6-ylpyrimidin-2-amine:** ¹H NMR
10 (400MHz, d₆-DMSO): 9.57 (s, 1H), 9.0 (d, 1H), 8.8 (s, 1H), 8.58 (d, 1H), 8.52 (d, 2H), 8.18 (d, 1H), 7.72 (d, 2H), 7.63 (q, 1H), 7.51 (d, 1H), 6.96 (d, 2H), 3.75 (t, 4H), 3.07 (4H). MS(EI) for C₂₃H₂₁N₅O : 384 (MH⁺).
- [0409] N-(4-morpholin-4-ylphenyl)-4-quinoxalin-6-ylpyrimidin-2-amine:** ¹H NMR
15 (400MHz, d₆-DMSO): 9.6 (s, 1H), 9.04 (d, 2H), 8.88 (s, 1H), 8.63 (d, 1H), 8.6 (d, 1H), 8.27 (d, 1H), 7.7 (d, 2H), 7.63 (d, 1H), 6.95 (d, 2H), 3.75 (t, 4H), 3.06 (t, 4H). MS (EI) for C₂₂H₂₀N₆O : 385 (MH⁺).
- [0410] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl]-D-alaninamide:** ¹H NMR (400MHz, d₆-DMSO): 9.25 (s, 1H), 8.3 (s, 1H), 7.7 (d, 2H), 7.66 (d, 2H), 7.6 (d, 2H), 6.85 (d, 2H), 3.5 (q, 1H), 3.03 (t, 4H), 2.5 (t, 4H), 2.35 (q, 2H),
20 2.21 (s, 3H), 1.23 (d, 3H), 1.02 (t, 3H). MS (EI) for C₂₆H₃₃N₇O : 460.5 (MH⁺).
- [0411] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl]-D-prolinamide:** ¹H NMR (400MHz, d₆-DMSO): 10.27 (s, 1H), 9.25 (s, 1H), 8.3 (s, 1H), 7.8 (d, 2H), 7.65 (d, 2H), 7.61 (d, 2H), 6.86 (d, 2H), 3.73 (m, 1H), 3.03 (t, 4H), 2.9 (t, 2H), 2.5 (t, 4H), 2.36 (q, 2H), 2.2 (s, 3H), 2.1-2.0 (m, 1H), 1.85-1.75 (m, 1H), 1.67 (p, 2H),
25 1.02 (t, 3H). MS (EI) for C₂₈H₃₅N₇O : 486 (MH⁺).
- [0412] N-ethyl-4-{4-[(4-{4-[(tetrahydrofuran-2-ylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}piperazine-1-carboxamide:** ¹H NMR (400MHz, d₆-DMSO): 9.93 (s, 1H), 9.4 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.88 (d, 2H), 7.67 (d, 2H), 7.3 (d, 1H), 6.96 (d, 2H), 6.6 (t, 1H), 4.43 (t, 1H), 4.0 (q, 1H), 3.86 (q, 1H), 3.42 (t, 4H), 3.05 (p, 2H), 3.01 (t, 4H), 2.27-2.17 (m, 1H), 2.06-1.97 (m, 1H), 1.88 (p, 2H), 1.02 (t, 3H). MS (EI) for C₂₈H₃₃N₇O₃ : 516 (MH⁺).
- [0413] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1H-imidazole-4-carboxamide:** ¹H NMR (400MHz, d₆-DMSO): 12.76 (br, 1H), 10.12 (s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.13 (d, 2H), 8.01 (d, 2H), 7.87 (s, 2H), 7.7 (d, 2H), 7.3 (d, 1H), 6.93 (d, 2H),
35 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for C₂₄H₂₃N₇O₂ : 442 (MH⁺).
- [0414] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1H-pyrrole-2-carboxamide:** ¹H NMR (400MHz, d₆-DMSO): ¹H NMR (400MHz, d₆-DMSO): 11.75

5 (s, 1H), 10.0 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.15 (d, 2H), 7.92 (d, 2H), 7.7 (d, 2H), 7.3 (d, 1H), 7.12 (s, 1H), 7.0 (s, 1H), 6.93 (d, 2H), 6.2 (d, 1H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for C₂₅H₂₄N₆O₂ : 441 (MH⁺).

[0415] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1H-

imidazole-2-carboxamide: ¹H NMR (400MHz, d₆-DMSO): 13.2 (br, 1H), 10.65 (s, 1H),
10 9.4 (s, 1H), 8.44 (d, 1H), 8.15 (d, 2H), 8.04 (d, 2H), 7.68 (d, 2H), 7.4-7.2 (m, 3H), 6.95 (d, 2H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for C₂₄H₂₃N₇O₂ : 442 (MH⁺).

[0416] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(2-(pyridin-3-

yl)-ethylamino)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 9.00 (s, 1H), 8.92 (d, 1H), 8.64 (d, 1H), 8.35 (d, 1H), 8.17-8.14 (m, 3H), 7.75 (d, 2H), 7.61 (d, 2H), 7.22 (d, 1H), 7.00 (d,
15 2H), 3.89-3.79 (m, 4H), 3.33-3.21 (m, 2H), 3.15-3.07 (m, 4H), 1.92 (s, 2H); MS (EI): 510.4 (MH⁺).

[0417] 2-(3-(4-methylpiperazin-1-yl)propylamino)-N-(4-(2-(4-morpholinophenyl-

amino)pyrimidin-4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 8.36 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.61 (d, 2H), 7.22 (d, 1H), 6.99 (d, 2H), 3.87-3.81 (m, 4H), 3.68
20 (s, 2H), 3.13-3.07 (m, 4H), 2.98-2.88 (m, 2H), 2.82-2.62 (m, 8H), 2.39 (s, 3H), 1.93 (s, 3H), 1.89-1.79 (m, 2H); MS (EI): 512.6 (MH⁺).

[0418] 2-(1-methylpiperidin-4-ylamino)-N-(4-(2-(4-morpholinophenylamino)-

pyrimidin-4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 8.35 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.61 (d, 2H), 7.22 (d, 1H), 6.99 (d, 2H), 3.87-3.81 (m, 4H), 3.47 (s, 2H),
25 3.15-3.09 (m, 4H), 3.06-2.95 (m, 2H), 2.69-2.55 (m, 1H), 2.39 (s, 3H), 2.38-2.22 (m, 2H), 2.03-1.93 (m, 2H), 1.90 (s, 2H), 1.63-1.43 (m, 2H); MS (EI): 545.2 (MH⁺).

[0419] 2-(2-amino-2-oxoethylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-

4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 8.27 (d, 1H), 8.04 (d, 2H), 7.68 (d, 2H), 7.53 (d, 2H), 7.13 (d, 1H), 6.91 (d, 2H), 4.54 (s, 2H), 3.79-3.73 (m, 4H), 3.39 (s, 2H),
30 3.06-3.00 (m, 4H), 1.86 (s, 2H); MS (EI): 462.1 (MH⁺).

[0420] 2-morpholino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-

acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.0 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 3.78-3.70 (m, 4H), 3.69-3.61 (m, 4H), 3.08-3.02 (m, 4H), 2.56-2.46 (m, 4H); MS (EI): 475.3 (MH⁺).

35 **[0421] 2-((2-aminoethyl)(methyl)amino)-N-(4-(2-(4-morpholinophenylamino)-**

pyrimidin-4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 8.36 (d, 1H), 8.13 (d, 2H), 7.76 (d, 2H), 7.62 (d, 2H), 7.22 (d, 1H), 7.00 (d, 2H), 3.89-3.81 (m, 4H), 3.51 (s, 2H),

5 3.14-3.07 (m, 4H), 3.03-3.96 (m, 2H), 2.94-2.88 (m, 2H), 2.67 (s, 3H); MS (EI)

$C_{25}H_{31}N_7O_2$: 462.4 (MH⁺).

[0422] 2-(1H-pyrazol-5-ylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)-phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 8.35 (d, 2H), 8.12 (d, 2H), 7.74 (d, 2H), 7.61 (d, 2H), 7.40 (s, 1H), 7.21 (d, 1H), 6.99 (d, 2H), 5.69 (s, 1H), 3.95 (s, 2H), 3.86-3.80 (m, 4H), 3.14-3.07 (m, 4H), 1.96 (s, 2H); MS (EI): 471.1 (MH⁺).

[0423] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(piperazin-1-yl)-acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.39 (s, 1H), 9.53 (s, 1H), 8.87 (s, 2H), 8.48 (d, 1H), 8.17 (d, 1H), 7.80 (d, 2H), 7.10 (d, 2H), 7.33 (d, 1H), 7.04 (d, 2H), 3.82-3.74 (m, 4H), 3.32-3.24 (m, 2H), 3.19-3.09 (m, 4H), 3.08-3.02 (m, 2H), 2.95 (s, 2H), 2.79 (s, 2H), 1.96 (s, 2H); MS (EI): 474.2 (MH⁺).

[0424] (S)-benzyl 2-(2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylamino)-2-oxoethylamino)propanoate: ¹H-NMR (400MHz, CD₃OD): 8.25 (d, 2H), 8.01 (d, 2H), 7.61 (d, 2H), 7.51 (d, 2H), 7.30-7.08 (m, 5H), 7.12 (d, 1H), 6.90 (d, 2H), 5.15-5.05 (m, 2H), 3.78-3.73 (m, 4H), 3.43 (q, 1H), 3.33 (d, 2H), 3.05-2.97 (m, 4H), 1.28 (d, 3H); MS (EI): 567.2 (MH⁺).

[0425] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyrimidin-4-ylamino)acetamide: ¹H-NMR (400MHz, CD₃OD): 10.90 (s, 1H), 9.63 (s, 1H), 9.18 (d, 2H), 8.78 (s, 1H), 8.50 (s, 1H), 8.26 (d, 1H), 8.18 (d, 2H), 7.82-7.68 (m, 4H), 7.36 (d, 1H), 7.11 (d, 2H), 6.84 (s, 1H), 5.16 (s, 2H), 3.83-3.77 (m, 4H), 2.54-2.47 (m, 4H); MS (EI): 483.2 (MH⁺).

[0426] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(piperidin-1-yl)-acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.98 (s, 1H), 9.84 (s, 1H), 9.65 (s, 1H), 8.50 (d, 1H), 8.20 (d, 2H), 7.90-7.70 (m, 4H), 7.36 (d, 1H), 7.11 (d, 2H), 4.12-4.07 (m, 2H), 3.87-3.77 (m, 4H), 3.62-3.42 (m, 2H), 3.23-3.13 (m, 4H), 2.51 (s, 2H), 1.94-1.64 (m, 6H), 1.45-1.38 (m, 2H); MS (EI): 473.4 (MH⁺).

[0427] 2-(ethylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.78 (s, 1H), 9.45 (s, 1H), 8.87 (s, 2H), 8.48 (d, 1H), 8.17 (d, 2H), 7.77 (d, 2H), 7.71 (d, 2H), 7.33 (d, 1H), 7.04 (d, 2H), 4.04-3.97 (m, 2H), 3.82-3.74 (m, 4H), 3.19-3.02 (m, 6H), 1.12 (t, 3H); MS (EI): 433.3 (MH⁺).

[0428] 2-(1H-imidazol-1-yl)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.87 (s, 1H), 9.56 (s, 1H), 9.11 (s, 1H), 8.45 (d, 1H), 8.15 (d, 2H), 7.76 (d, 2H), 7.70 (d, 2H), 7.32 (d, 1H), 7.05 (d, 2H), 5.26 (s, 2H), 3.82-3.72 (m, 4H), 3.18-3.08 (m, 4H); MS (EI): 456.3 (MH⁺).

- 5 **[0429] 4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzoic acid:** ¹H-NMR (400MHz, d6-DMSO): 9.56 (s, 1H), 8.55 (d, 1H), 8.27 (d, 2H), 8.09 (d, 2H), 7.68 (d, 2H), 7.41 (d, 1H), 6.97 (d, 2H), 3.80-3.72 (m, 4H), 3.11-3.03 (m, 4H); MS (EI): 377.3 (MH⁺).
- [0430] N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(phenylamino)-acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.29 (s, 1H), 9.59 (s, 1H), 8.46 (d, 1H), 8.14 (d, 2H), 7.79 (d, 2H), 7.73 (d, 2H), 7.33 (d, 1H), 7.19-7.00 (m, 4H), 6.70-6.50 (m, 3H), 3.96-3.88 (m, 4H), 3.22-3.12 (m, 4H); MS (EI): 481.1 (MH⁺).
- 10 **[0431] 4-(4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl)-N-(4-morpholinophenyl)-pyrimidin-2-amine:** ¹H-NMR (400MHz, d6-DMSO): 9.53 (s, 1H), 8.54 (d, 1H), 8.35 (d, 2H), 8.12 (d, 2H), 7.65 (d, 2H), 7.40 (d, 1H), 6.92 (d, 1H), 3.76-3.70 (m, 4H), 3.06-3.00 (m, 4H), 2.59 (s, 3H); MS (EI): 415.3 (MH⁺).
- 15 **[0432] (R)-4-(4-(4-(4-(2-aminopropanamido)phenyl)pyrimidin-2-ylamino)phenyl)-N-ethylpiperazine-1-carboxamide:** ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.96 (d, 2H), 6.59 (t, 1H), 3.54-3.46 (m, 1H), 3.44-3.36 (m, 4H), 3.12-2.97 (m, 6H), 1.24 (d, 3H), 1.02 (t, 2H), 0.95 (t, 2H); MS (EI): 487.1 (MH⁻).
- 20 **[0433] (R)-2-amino-N-(4-(2-(4-(4-((R)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenyl)-amino)pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.97-3.92 (m, 1H), 3.72-3.58 (m, 4H), 3.51-3.42 (m, 2H), 3.14-2.99 (m, 4H), 2.68-2.62 (m, 1H), 2.12-2.00 (m, 1H), 1.74-1.70 (m, 1H), 1.70-1.56 (m, 2H), 1.24 (d, 3H); MS (EI): 513.2 (MH⁻).
- 25 **[0434] (R)-2-amino-N-(4-(2-(4-(4-((S)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenyl)-amino)pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 4.65 (t, 1H), 3.89-3.81 (m, 1H), 3.75-3.58 (m, 3H), 3.54-3.28 (m, 2H), 3.15-2.98 (m, 4H), 2.72-2.58 (m, 1H), 2.06-1.97 (m, 2H), 1.74-1.54 (m, 2H), 1.24 (d, 3H); MS (EI): 515.4 (MH⁺).
- 30 **[0435] N-[4-(2-{[4-(4-L-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-alaninamide:** ¹H-NMR (400MHz, d6-DMSO): 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.57 (dd, 1H), 3.68-3.58 (m, 4H), 3.46 (dd, 1H), 3.12-2.98 (m, 4H), 1.23 (d, 3H), 1.12 (d, 3H); MS (EI): 489.4 (MH⁺).
- 35 **[0436] (R)-2-amino-N-(4-(2-(4-(4-((S)-2-aminopropanoyl)piperazin-1-yl)phenylamino)-pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H),

5 6.97 (d, 2H), 3.17 (s, 2H), 3.12-3.07 (m, 1H), 3.05-2.98 (m, 1H), 2.70-2.64 (m, 4H), 2.36-2.28 (m, 4H), 1.23 (d, 3H); MS (EI): 544.4 (MH⁺).

[0437] **3,3,3-trifluoro-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.42 (br s, 1H), 9.41 (s, 1H), 8.46 (d, 1H), 8.15 (d, 2H), 7.89 (d, 2H), 7.68 (d, 2H), 7.57 (br s, 1H), 7.31 (d, 1H), 6.94 (d, 10 2H), 4.83-4.74 (m, 1H), 3.78-3.70 (m, 4H), 3.09-3.01 (m, 4H); MS (EI): 474.3(MH⁺).

[0438] **(R)-2-hydroxy-2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.72 (s, 1H), 9.34 (s, 1H), 8.41 (d, 1H), 8.08 (d, 2H), 7.89 (d, 2H), 7.64 (d, 2H), 7.27 (d, 1H), 6.91 (d, 2H), 5.66 (s, 1H), 4.21-4.13 (m, 1H), 3.72 (q, 1H), 3.15 (d, 3H), 3.12-3.02 (m, 4H), 1.80-1.72 (m, 1H), 1.59-1.51 (m, 15 1H), 1.32 (s, 3H), 0.82 (t, 3H); MS (EI): 448.4 (MH⁺).

[0439] **(S)-2-hydroxy-2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.72 (s, 1H), 9.34 (s, 1H), 8.41 (d, 1H), 8.08 (d, 2H), 7.89 (d, 2H), 7.64 (d, 2H), 7.27 (d, 1H), 6.91 (d, 2H), 5.66 (s, 1H), 4.21-4.13 (m, 1H), 3.72 (q, 1H), 3.19-3.11 (d, 3H), 3.07-3.02 (m, 4H), 1.81-1.71 (m, 1H), 1.60-1.50 (m, 1H), 1.32 (s, 3H), 0.82 (t, 3H); MS (EI): 448.1 (MH⁺). 20

[0440] **(R)-2-methoxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.07 (s, 1H), 9.35 (s, 1H), 8.41 (d, 1H), 8.11 (d, 2H), 7.84 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.90 (d, 2H), 3.89 (q, 1H), 3.75-3.67 (m, 4H), 3.30 (s, 3H), 3.04-2.96 (m, 4H), 1.31 (d, 3H); MS (EI): 434.3 (MH⁺).

25 [0441] **(S)-2-methoxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.07 (s, 1H), 9.35 (s, 1H), 8.41 (d, 1H), 8.11 (d, 2H), 7.84 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.90 (d, 2H), 3.89 (q, 1H), 3.75-3.67 (m, 4H), 3.30 (s, 3H), 3.04-2.96 (m, 4H), 1.31 (d, 3H); MS (EI): 434.3 (MH⁺).

[0442] **1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-cyclopentanecarboxamide:** ¹H-NMR (400MHz, d6-DMSO): 9.39 (s, 1H), 8.44 (d, 1H), 8.14 (d, 2H), 7.87 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.94 (d, 2H), 3.78-3.70 (m, 4H), 3.08-3.00 (m, 4H), 2.10-2.00 (m, 2H), 1.86-1.75 (m, 2H), 1.74-1.62 (m, 2H), 1.60-1.50 (m, 2H); MS (EI): 459.4 (MH⁺). 30

[0443] **(S)-2-hydroxy-3,3-dimethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide:** ¹H-NMR (400MHz, d6-DMSO): 9.83 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.89 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 5.85 (d, 1H), 3.78-3.70 (m, 4H), 3.47 (q, 1H), 3.09-3.01 (m, 4H), 0.97 (d, 9H); MS (EI): 462.4 (MH⁺). 35

- 5 [0444] (R)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d6-DMSO): 9.91 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.89 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (br s, 1H), 3.85 (d, 1H), 3.77-3.69 (m, 4H), 3.10-3.02 (m, 4H), 1.80-1.51 (m, 6H), 1.30-1.02 (m, 5H); MS (EI): 488.1 (MH+).
- 10 [0445] (S)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H-NMR (400MHz, d6-DMSO): 9.91 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.89 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (br s, 1H), 3.85 (d, 1H), 3.77-3.69 (m, 4H), 3.07-2.98 (m, 4H), 1.78-1.50 (m, 6H), 1.25-1.00 (m, 5H); MS (EI): 488.1 (MH+).
- 15 [0446] (S)-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H-NMR (400MHz, d6-DMSO): 9.95 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.87 (br s, 1H), 4.23-4.15 (m, 1H), 3.79-3.71 (m, 4H), 3.08-3.00 (m, 4H), 2.51 (d, 3H); MS (EI): 420.4 (MH+).
- [0447] 1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-cyclobutanecarboxamide: ¹H-NMR (400MHz, d6-DMSO): 9.39 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.87 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 3.79-3.71 (m, 4H), 3.09-3.00 (m, 4H), 2.00-1.85 (m, 4H), 1.84-1.76 (m, 2H); MS (EI): 445.4 (MH+).
- 20 [0448] 4-(4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine: ¹H-NMR (400MHz, d6-DMSO): 9.57 (s, 1H), 8.57 (d, 1H), 8.39 (d, 2H), 8.26 (d, 2H), 7.67 (d, 2H), 7.44 (d, 1H), 6.95 (d, 2H), 3.78-3.72 (m, 4H), 3.09-3.03 (m, 4H), 2.46 (s, 3H); MS (EI): 415.0 (MH+).
- [0449] N-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-2-phenylacetamide: ¹H-NMR (400MHz, d6-DMSO): 10.45 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.76 (d, 2H), 7.64 (d, 2H), 7.38-7.33 (m, 3H), 7.27 (d, 1H), 6.92 (d, 2H), 3.69 (s, 2H), 3.10-3.04 (m, 4H), 2.35 (q, 3H), 1.89 (s, 2H), 1.03 (t, 2H); MS (EI): 493.1 (MH+).
- 30 [0450] 1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidin-2-one: ¹H-NMR (400MHz, d6-DMSO): 8.26 (d, 1H), 8.14 (d, 2H), 7.77 (d, 2H), 7.65 (d, 2H), 7.36 (d, 1H), 7.25 (d, 2H), 3.92-3.84 (m, 5H), 3.82-3.74 (m, 1H), 3.74-3.60 (m, 1H), 3.42-3.30 (m, 4H), 3.06-3.02 (m, 1H), 2.16-2.06 (m, 2H); MS (EI): 416.1 (MH+).
- 35 [0451] (S)-2-hydroxy-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide: ¹H-NMR (400MHz, d6-DMSO): 9.90 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (d, 1H), 3.86

5 (dd, 1H), 3.78-3.73 (m, 4H), 3.08-3.02 (m, 4H), 0.96 (d, 3H), 0.87 (d, 3H); MS (EI): 448.3 (MH⁺).

[0452] (R)-2-hydroxy-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide: ¹H-NMR (400MHz, d6-DMSO): 9.90 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (d, 1H), 3.86
10 (dd, 1H), 3.78-3.73 (m, 4H), 3.08-3.02 (m, 4H), 0.96 (d, 3H), 0.87 (d, 3H); MS (EI): 448.3 (MH⁺).

[0453] (R)-2-amino-N-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H),
15 6.95 (d, 2H), 3.63-3.56 (m, 2H), 3.43-3.37 (m, 3H), 3.18 (d, 1H), 3.07-2.98 (m, 4H), 2.25-2.02 (m, 4H), 1.98-1.83 (m, 1H), 1.82-1.70 (m, 1H), 1.23 (d, 3H); MS (EI): 500.2 (MH⁺).

[0454] (R)-2-amino-N-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.73-3.67 (m, 4H), 3.52-4.42
20 (m, 1H), 3.08-3.02 (m, 4H), 1.25 (s, 3H), 1.23 (d, 3H); MS (EI): 502.4 (MH⁺).

[0455] 4-[4-(ethylamino)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine: ¹H-NMR (400 MHz, DMSO): 9.202 (s, 1H), 8.3 (d, 2H), 7.948 (d, 2H), 7.689 (q, 2H), 7.134 (d, 1H), 6.93 (d, 2H), 6.657(d, 2H), 6.285 (t, 1H), 3.754 (t, 4H), 3.132-3.113 (m, 2H), 3.04 (t, 4H), 1.187 (t, 3H). MS (EI) for C₂₂H₂₅N₅O: 376.3 (MH⁺).

25 [0456] N-{4-[2-(phenylamino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, DMSO): 10.233 (s, 1H), 9.63 (s, 1H), 8.513 (d, 1H), 8.147 (d, 2H), 7.854 (d, 2H), 7.774 (d, 2H), 7.362-7.305 (m, 3H), 6.961 (t, 1H), 2.098 (s, 3H). MS (EI) for C₁₈H₁₆N₄O: 305.3 (MH⁺).

[0457] N-{4-[2-({4-[(4-ethylpiperazin-1 yl)carbonyl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, DMSO): 10.236 (s, 1H), 9.889 (s, 1H), 8.549 (d, 1H), 8.167-8.134 (m, 2H), 7.93-7.903 (m, 2H), 7.782 (d, 2H), 7.418-7.369 (m, 3H), 3.509 (br s, 4H), 2.378-2.324 (m, 6H), 2.097 (s, 3H), 1.025 (t, 3H). MS (EI) for C₂₅H₂₈N₆O₂: 445.4 (MH⁺).

[0458] N-[4-(2-({3-(morpholin-4-ylcarbonyl)phenyl}amino)pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, DMSO): 10.237 (s, 1H), 9.823 (s, 1H), 8.541 (d, 1H), 8.152 (d, 2H), 8.026 (t, 1H), 7.847 (d, 1H), 7.772 (d, 2H), 7.392 (m, 2H), 6.996 (d, 1H), 3.76-3.36 (br s, 8H), 2.094 (s, 3H). MS (EI) for C₂₃H₂₃N₅O₃: 418.3 (MH⁺).

- 5 **[0459] N-(4-(2-(3-(2-(dimethylamino)ethoxy)phenylamino)pyrimidin-4-yl)phenyl)-acetamide:** ¹H NMR (400 MHz, DMSO): 10.471 (br s, 1H), 10.42 (s, 1H), 9.816 (s, 1H), 8.534 (d, 1H), 8.158 (d, 2H), 7.804 (d, 2H), 7.7 (t, 1H), 7.413-7.383 (m, 2H), 7.29 (t, 1H), 6.636 (m, 1H), 4.381 (t, 2H), 3.531 (q, 2H), 2.883 (d, 6H), 2.106 (s, 3H). MS (EI) for C₂₂H₂₅N₅O₂: 392.3 (MH⁺).
- 10 **[0460] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide:** ¹H-NMR (400 MHz, DMSO): 10.532 (s, 1H), 9.407 (s, 1H), 8.47 (d, 1H) 8.189 (d, 2H), 7.982 (m, 4H), 7.7-7.54 (m, 5H), 7.325 (d, 1H), 6.959 (d, 2H), 3.747 (t, 4H), 3.054 (t, 4H). MS (EI) for C₂₇H₂₅N₅O₂: 452.1 (MH⁺).
- [0461] N-[4-(2-{[4-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide:**
- 15 ¹H-NMR (400 MHz, DMSO): 10.222 (s, 1H), 9.433 (s, 1H), 8.455 (s, 1H), 8.121 (d, 2H), 7.725 (q, 4H), 7.293 (d, 1H), 6.91 (d, 2H), 3.739 (s, 3H), 2.093 (s, 3H). MS (EI) for C₁₉H₁₈N₄O₂: 335 (MH⁺).
- [0462] 4-(4-chlorophenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400 MHz, DMSO): 9.488 (s, 1H), 8.514 (d, 1H), 8.185 (d, 2H), 7.665-7.606 (q, 4H), 7.354 (d, 1H), 6.918 (d, 2H), 3.757 (t, 4H), 3.048 (t, 4H). MS (EI) for C₂₀H₁₉ClN₄O: 367 (MH⁺).
- 20 **[0463] N-[4-(2-{[3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide:** ¹H-NMR (400 MHz, DMSO): 10.235 (s, 1H), 9.633 (s, 1H), 8.52 (d, 1H), 8.15 (d, 2H), 7.65 (t, 1H), 7.369 (d, 2H), 7.209 (t, 1H), 6.535 (q, 1H), 3.77 (s, 3H), 2.092 (s, 3H). MS (EI) for C₁₉H₁₈N₄O₂: 335 (MH⁺).
- 25 **[0464] 1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-(phenylmethyl)urea:** ¹H NMR (400 MHz, DMSO): 9.347 (s, 1H), 9 (s, 1H), 8.414 (d, 1H), 8.067 (d, 2H), 7.688 (d, 2H), 7.583 (d, 2H), 7.351-7.31 (m, 4H), 7.237 (d, 2H), 6.943 (d, 2H), 6.831 (t, 1H), 4.33 (d, 2H), 3.742 (t, 4H), 3.044 (t, 4H). MS (EI) for C₂₈H₂₈N₆O₂: 481.4 (MH⁺).
- [0465] N-(4-{2-[(4-{4-[(2S)-pyrrolidin-2-ylmethyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)-D-prolinamide:** ¹H NMR (400 MHz, DMSO): 10.231 (br s, 1H), 9.376 (s, 1H), 8.443 (d, 1H), 8.134 (d, 2H), 7.848 (d, 2H), 7.663 (d, 2H), 7.29 (d, 1H), 6.936 (d, 2H), 3.74 (m, 1H), 3.505 (m, 1H), 3.08-2.89 (m, 6H), 2.64 (m, 2H), 2.374 (m, 1H), 2.069 (m, 1H), 1.938-1.648 (m, 9H), 1.452 (m, 1H). MS (EI) for C₃₀H₃₈N₈O: 527.3 (MH⁺).
- 30 **[0466] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2,3-dihydro-1H-isoindole-1-carboxamide:** ¹H NMR (400 MHz, DMSO): 11.345 (s, 1H), 10.326 (br s, 1H), 9.525 (s, 1H), 9.479 (br s, 1H), 8.474 (d, 1H), 8.19 (d, 2H), 7.799 (d, 2H), 7.7-7.627 (m,

5 3H), 7.469-7.415 (m, 3H), 7.323 (d, 1H), 7.025 (d, 2H), 5.688 (br s, 1H), 4.578 (m, 2H), 3.757 (s, 4H), 3.105 (s, 4H). MS (EI) for $C_{29}H_{28}N_6O_2$: 492.58 (MH^+).

[0467] N-{4-[2-({4-[4-(2-piperazin-1-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide: 1H NMR (400 MHz, DMSO): 9.938 (s, 1H), 9.412 (s, 1H), 8.456 (d, 1H), 7.135 (d, 2H), 7.886 (d, 2H), 7.691 (d, 2H), 7.304 (d, 1H), 10 6.975 (d, 2H), 4.45 (m, 1H), 4.01 (q, 1H), 3.878 (q, 1H), 3.705 (br s, 4H), 3.592 (br s, 4H), 3.148 (s, 2H), 3.1 (br s, 2H), 3.02 (br s, 2H), 2.719 (br s, 4H), 2.354 (br s, 4H), 2.21 (m, 1H), 2.021 (m, 1H), 1.862 (m, 2H). MS (EI) for $C_{31}H_{38}N_8O_3$: 571 (MH^+).

[0468] N-(4-{2-[(4-{4-[(4-chloro-1-methyl-1H-pyrazol-3-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: 1H NMR (400 MHz, DMSO): 15 10.215 (s, 1H), 9.363 (s, 1H), 8.439 (d, 1H), 8.131 (d, 2H), 7.901 (s, 1H), 7.841 (d, 2H), 7.651 (d, 2H), 7.286 (d, 1H), 6.916 (d, 2H), 3.802 (s, 3H), 3.7 (m, 1H), 3.47 (s, 2H), 3.048 (br s, 4H), 2.92 (t, 2H), 2.568 (t, 4H), 2.08 (m, 1H), 1.816 (m, 1H), 1.691 (m, 2H). MS (EI) for $C_{30}H_{34}ClN_9O$: 572.4 (MH^+).

[0469] N-{4-[2-({4-[4-(2-hydroxyethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide: 1H NMR (400 MHz, DMSO): 20 10.176 (s, 1H), 9.927 (s, 1H), 8.374 (d, 1H), 8.065 (d, 2H), 7.773 (d, 2H), 7.587 (d, 2H), 7.219 (d, 1H), 6.857 (d, 2H), 4.396 (br s, 1H), 3.715 (t, 1H), 3.468 (br s, 4H), 2.996-2.954 (m, 6H), 2.873 (t, 2H), 2.74 (s, 1H), 2.368 (t, 2H), 2.369 (t, 2H), 2.368 (t, 2H), 2.038 (m, 1H), 1.749 (m, 1H), 1.61 (m, 2H). MS (EI) for $C_{27}H_{33}N_7O_2$: 488.3 (MH^+).

25 [0470] N-(4-{2-[(4-{4-[(1-methyl-1H-pyrrol-2-yl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)-D-prolinamide: 1H NMR (400 MHz, DMSO): 10.194 (s, 1H), 9.349 (s, 1H), 8.42 (d, 1H), 8.11 (d, 2H), 7.822 (d, 2H), 7.636 (d, 2H), 7.267 (d, 1H), 7.076 (s, 1H), 6.902 (d, 2H), 6.75 (s, 1H), 3.738 (m, 1H), 3.701 (s, 3H), 3.553 (s, 2H), 3.306 (m, 4H), 3.031 (br s, 4H), 2.892 (t, 2H), 2.728 (s, 1H), 2.054 (m, 1H), 1.803 (m, 30 1H), 1.647 (m, 2H). MS (EI) for $C_{31}H_{36}N_8O$: 538.3 (MH^+).

[0471] N-(4-{2-[(4-{4-[(2R)-pyrrolidin-2-ylmethyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)-D-prolinamide: 1H NMR (400 MHz, DMSO): 8.319-8.251 (m, 3H), 7.86 (d, 2H), 7.636 (d, 2H), 7.56 (d, 1H), 7.282 (d, 2H), 4.479 (t, 1H), 4.106 (m, 1H), 3.514 (br s, 4H), 3.496-3.303 (m, 11H), 2.252 (m, 1H), 2.342 (m, 1H), 2.19-2.094 (m, 6H), 35 1.835 (m, 1H). MS (EI) for $C_{30}H_{36}N_8O$: 528.5 (MH^+).

5

[0472] (2S,3aS,7aS)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-octahydro-1H-indole-2-carboxamide: ¹H NMR (400 MHz, DMSO): 10.857 (s, 1H), 9.84 (s, 1H), 9.714 (br s, 1H), 8.531 (d, 1H), 8.207 (m, 3H), 7.816 (d, 4H), 7.42 (d, 1H), 4.448 (m, 1H), 3.874 (t, 4H), 3.681 (br s, 1H), 3.322 (t, 4H), 2.504 (m, 2H), 2.09 (m, 1H),
 10 1.913 (m, 1H), 1.615 (m, 4H), 1.371-1.259 (m, 3H). MS (EI) for C₂₉H₃₄N₆O₂: 499.5 (MH⁺).

[0473] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclopropane-carboxamide: ¹H NMR (400 MHz, DMSO): 10.476 (s, 1H), 9.383 (s, 1H), 8.443 (d, 1H), 8.123 (d, 2H), 7.768 (d, 2H), 7.686 (d, 2H), 7.279 (d, 1H), 6.946 (d, 2H), 3.74 (t, 4H), 3.046 (t, 4H), 1.824 (m, 1H), 0.829 (m, 4H). MS (EI) for C₂₄H₂₅N₅O₂: 416 (MH⁺).

15 **[0474] N-[4-(2-{[4-(dimethylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide:** ¹H-NMR (400 MHz, d₆-DMSO): 10.38 (s, br, 1H), 9.92 (s, br, 1H), 8.85 (d, 2H), 8.11 (d, 2H), 7.92 (d, 2H), 7.79 (d, 2H), 7.67 (s, 2H), 7.42 (d, 1H), 3.10 (s, 6H), 2.10 (s, 3H). MS (EI): 348 (MH⁺).

[0475] N-(4-{2-[(4-chlorophenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.30 (s, 1H), 9.88 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 7.87 (d, 2H), 7.75 (d, 2H), 7.41 (d, 2H), 7.37 (1H), 2.10 (s, 3). MS (EI): 339 (MH⁺).

[0476] N-[4-(2-{[4-(1H-pyrrol-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400 MHz, d₆-DMSO): 10.26 (br s, 1H), 9.78 (br s, 1H), 8.52 (d, 1H), 8.15 (d, 2H), 7.92 (d, 2H), 7.77 (d, 2H), 7.54 (d, 2H), 7.38 (d, 1H), 7.31 (m, 2H), 6.24 (m, 2H). 2.09 (s,
 25 3H). MS (EI): 370 (MH⁺).

[0477] ethyl 1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-piperidine-4-carboxylate: ¹H NMR (400 MHz, d₆-DMSO): 10.25 (s, 1H), 9.49 (br s, 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.75 (d, 4H), 7.30 (d, 1H), 4.10 (q, 2H), 3.56 (d, 2H), 2.86 (br s, 2H), 2.09 (s, 3H), 1.97 (br d, 2H), 1.35 (m, 2H), 1.30 (m, 2H), 0.88 (t, 3H). MS (EI): 460
 30 (MH⁺).

[0478] N-[4-(2-{[4-(4-phenylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.28 (s, 1H), 9.68 (br s 1H), 9.49 (d, 1H), 8.14 (d, 2H), 7.82 (br s 1H), 7.70 (d, 2H), 7.35 (d, 1H), 7.28 (t, 2H), 7.08 (d, 2H), 6.88 (t, 1H), 3.45 (br s, 8H), 2.09 (s, 3H). MS (EI): 465 (MH⁺).

35 **[0479] N-{4-[2-({4-[(2R,6S)-2,6-dimethylmorpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.28 (s, 1H), 9.79 (br s, 1H), 8.47

5 (d, 1H), 8.12 (d, 1H), 8.10 (d, 1H), 7.80 (br d, 2H), 7.75 (d, 2H), 7.37 (d, 2H), 3.69 (br s, 4H), 3.54 (d, 2H), 2.07 (s, 3H), 1.16 (s, 3H), 1.14 (s, 3H) MS (EI): 418 (MH+).

[0480] 4-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
tetrahydro-2H-thiopyran-4-carboxamide 1,1-dioxide: ¹H NMR (400 MHz, d6-
DMSO): 11.06 (br s, 1H), 9.97 (br s, 1H), 9.29 (s, 2H), 8.56 (d, 1H), 8.21 (d, 2H), 7.94 (d,
10 2H), 7.89 (d, 1H), 7.58 (s, 1H), 7.47 (d, 1H), 5.32 (br s, 3H), 3.99 (s, 4H), 3.48 (m, 4H), 3.38
(m, 4H), 2.84 (m, 2H), 2.46 (m, 2H). MS (EI): 523 (MH+).

[0481] (2R)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
piperazine-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.33 (br s, 1H), 10.31 (br s,
1H), 8.58 (d, 1H), 8.22 (d, 2H), 7.94 (d, 2H), 7.89 (d, 2H), 7.77 (d, 2H), 7.53 (d, 1H), 4.55 (d,
15 1H), 4.08 (s, 5H), 3.54 (s, 5H), 3.43 (m, 2H), 3.28 (m, 2H). MS (EI): 460 (MH+).

[0482] 2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
1,2,3,4-tetrahydronaphthalene-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 11.10
(br s, 1H), 10.07 (br s, 1H), 8.89 (d, 2H), 8.56 (d, 1H), 8.20 (d, 2H), 7.98 (d, 2H), 7.92 (d,
2H), 7.70 (d, 2H), 7.48 (d, 1H), 7.10 (m, 4H), 6.10 (br s, 3H), 4.40 (s, 4H), 3.70 (d, 1H), 3.50
20 (s, 4H), 3.39 (d, 4H), 2.89 (m, 1H), 2.71 (m, 1H), 2.1 (m, 1H), 1.26 (m, 1H), 1.17 (m, 1H).
MS (EI): 521 (MH+).

[0483] 4-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
tetrahydro-2H-pyran-4-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.85 (br s, 1H),
9.85 (br s, 1H), 8.97 (s, 2H), 8.53 (d, 1H), 8.20 (d, 2H), 7.94 (d, 2H), 7.83 (d, 2), 7.43 (d,
25 2H), 4.39 (br s, 3H), 3.94 (s, 4H), 3.87 (d, 4H), 3.72 (m, 4H), 2.45 (m, 2H), 1.97 (d, 2H).
MS (EI): 475 (MH+).

[0484] (4S)-4-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-
yl}phenyl)prolinamide: ¹H NMR (400 MHz, d6-DMSO): 11.38 (s, 1H), 10.25 (m, 1H),
10.31 (s, 1H), 8.84 (m, 1H), 8.57 (d, 1H), 8.21 (d, 2H), 7.93 (d, 2H), 7.87 (d, 2H), 7.75 (d,
30 2H), 7.49 (d, 1H), 6.62 (m, 1H), 4.50 (s, 1H), 4.06 (s, 4H), 3.53 (s, 4H), 3.42 (m, 1H) < 3.17
(m, 1H), 2.45 (t, 1H), MS (EI): 461 (MH+).

[0485] 1-acetyl-4-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-
phenyl)piperidine-4-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.72 (s, 1H), 8.51 (d,
1H), 8.14 (dd, 2H), 8.12 (d, 1H), 8.08 (d, 1H), 7.00 (dd, 1H), 7.39 (d, 1H), 7.17 (d, 1H), 4.15
35 (d, 1H), 3.70 (m, 4H), 3.68 (d, 1H), 4.41 (m, 4H), 3.41 (m, 4H), 3.01 (m, 1H), 2.93 (m, 4H),
2.02 (s, 3H), 1.98 (m, 1H), 1.84 (m, 1H). MS (EI): 516 (MH+).

- 5 **[0486] O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.51 (br s, 1H), 9.99 (br s, 1H), 8.55 (d, 1H), 8.50 (br s, 2H), 8.19 (d, 2H), 7.88 (m, 4H), 7.46 (d, 1H), 5.19 (br s, 3H), 4.36 (m, 1H), 4.02 (br s, 4H), 3.88 (m, 1H), 3.46 (br s, 4H), 3.33 (s, 3H). MS (EI): 449 (MH⁺).
- [0487] N-[4-(2-{[4-(2,6-dimethylmorpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.30 (br s, 1H), 9.83 (br s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.81 (br s, 1H), 7.77 (d, 2H), 7.39 (d, 1H), 4.10 (br s, 4H), 3.56 (d, 2H), 2.10 (s, 3H), 1.18 (s, 3H), 1.71 (s, 3H). MS (EI): 418 (MH⁺).
- 10 **[0488] N-(4-{2-[(4-piperidin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400 MHz, d6-DMSO): 10.32 (br s, 1H), 9.89 (br s, 1H), 8.52 (d, 1H), 8.11 (d, 2H), 7.94 (d, 2H), 7.76 (d, 4H), 7.40 (d, 1H), 3.45 (br s, 4H), 3.36 (6H), 2.07 (s, 3H). MS (EI): 388 (MH⁺).
- [0489] O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.51 (br s, 1H), 9.99 (br s, 1H), 8.55 (d, 1H), 8.50 (br s, 2H), 8.19 (d, 2H), 7.88 (m, 4H), 7.46 (d, 1H), 5.19 (br s, 3H), 4.36 (m, 1H), 4.02 (br s, 4H), 3.88 (m, 1H), 3.46 (br s, 4H), 3.33 (s, 3H). MS (EI): 449 (MH⁺).
- 20 **[0490] 1,1-dimethylethyl (2R)-2-{[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]carbonyl}pyrrolidine-1-carboxylate:** ¹H NMR (400 MHz, d6-DMSO): 10.26 (br s, 1H), 9.38 (br s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.78 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.94 (d, 2H), 4.22 (m, 1H), 3.74 (m, 4H), 3.43 (m, 1H), 3.34 (m, 1H), 3.04 (m, 4H), 2.20 (m, 1H), 1.90 (m, 1H), 1.81 (m, 1H), 1.40 (s, 3H), 1.27 (s, 6H). MS (EI): 545 (MH⁺).
- 25 **[0491] 4-[4-(methylsulfonyl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400 MHz, d6-DMSO): 9.62 (s, 1H), 8.59 (d, 1H), 8.39 (d, 2H), 8.09 (d, 2H), 7.68 (d, 2H), 7.45 (d, 1H), 7.0 (s, br, 1H), 3.81-3.71 (m, 4H), 3.29 (s, 3H), 3.04-3.14 (m, 4H). MS (EI): 411 (MH⁺).
- 30 **[0492] 4-[3-(methylsulfonyl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400 MHz, d6-DMSO): 9.60 (s, 1H), 8.72 (s, 1H), 8.57 (d, 1H), 8.47 (d, 1H), 8.10 (d, 1H), 7.85 (d, 1H), 7.66 (d, 1H), 7.46 (d, 1H), 6.92 (d, 1H), 3.81-3.71 (m, 4H), 3.31 (s, 3H), 3.0-3.11 (m, 4H). MS (EI): 411 (MH⁺).
- [0493] 4-[4-(methylthio)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:** ¹H-NMR (400 MHz, d6-DMSO): 9.42 (s, 1H), 8.46 (d, 1H), 8.09 (d, 2H), 7.66 (d, 2H), 7.40 (d, 2H), 7.31 (d, 1H), 6.92 (d, 2H), 3.79-3.69 (m, 4H), 3.1-3.0 (m, 4H), 2.55 (s, 3H). MS (EI): 379 (MH⁺).
- 35

- 5 **[0494] N-(4-{2-[(3-bromo-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.72 (s, 1H), 8.52 (d, 1H), 8.27 (d, 1H), 8.13 (d, 2H), 7.81-7.71 (m, 3H), 7.42-7.32 (m, 1H), 7.18 (d, 1H), 3.78-3.69 (m, 4H), 2.97-2.87 (m, 4H), 2.09 (s, 3H). MS (EI): 469 (MH⁺).
- [0495] N-[4-(2-{4-{[2-(diethylamino)ethyl]oxy}-3-(4-ethylpiperazin-1-yl)phenyl]-amino}pyrimidin-4-yl)phenyl]acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.73 (d, 2H), 7.54 (s, 1H), 7.36-7.26 (m, 2H), 6.88 (d, 1H), 4.02-3.92 (m, 2H), 2.81-2.71 (m, 2H), 2.59-2.49 (m, 4H), 2.45-2.35 (m, 2H), 1.04 (t, 3H), 0.98 (t, 6H). MS (EI): 533 (MH⁺).
- 10 **[0496] N²,N²-dimethyl-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)glycinamide:** ¹H NMR (400 MHz, d6-DMSO): 9.99 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 3.63-3.59 (m, 4H), 3.25 (s, 3H), 3.11 (s, 2H), 3.10-3.05 (m, 2H), 3.04-2.99 (m, 2H), 2.65 (t, 2H), 2.29 (s, 6H). MS (EI): 519 (MH⁺).
- 15 **[0497] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclobutane-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 9.99 (s, 1H), 9.37 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.75 (m, 4H), 3.32 (m, 1H), 3.05 (m, 4H), 2.24 (m, 2H), 2.12 (m, 2H), 1.93 (m, 1H), 1.82 (m, 1H). MS (EI): 430 (MH⁺).
- 20 **[0498] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.10 (s, 1H), 9.36 (s, 1H), 8.42 (d, 1H), 8.09 (d, 2H), 7.74 (d, 2H), 7.64 (d, 2H), 7.24 (d, 1H), 6.91 (d, 2H), 3.71 (m, 4H), 3.61 (m, 1H), 3.52 (m, 4H), 3.02 (m, 4H). MS (EI): 431 (MH⁺).
- 25 **[0499] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.35 (s, 1H), 8.40 (d, 1H), 8.08 (d, 2H), 7.73 (d, 2H), 7.64 (d, 2H), 7.24 (d, 1H), 6.90 (d, 2H), 3.72 (m, 4H), 3.41 (m, 4H), 3.02 (m, 4H), 2.83 (m, 1H), 2.62 (m, 1H), 1.58 (m, 2H), 1.37 (m, 1H). MS (EI): 459 (MH⁺).
- 30 **[0500] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-4-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.12 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.76 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.37 (m, 1H), 3.03 (m, 4H), 3.00 (m, 1H), 2.50 (m, 2H), 2.47 (m, 1H), 1.73 (m, 2H), 1.54 (m, 2H). MS (EI): 459 (MH⁺).
- 35

- 5 [0501] 2-(methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.05 (s, 1H), 9.36 (s, 1H), 8.42 (d, 1H), 8.09 (d, 2H), 7.82 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.91 (d, 2H), 4.02 (s, 2H), 3.72 (m, 4H), 3.38 (s, 3H), 3.02 (m, 4H). MS (EI): 420 (MH⁺).
- [0502] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-2-
10 carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.85 (s, br, 1H), 9.31 (s, 1H), 8.36 (d, 1H), 8.04 (d, 2H), 7.75 (d, 2H), 7.60 (d, 2H), 7.21 (d, 1H), 6.87 (d, 2H), 3.67 (m, 4H), 3.21 (m, 1H), 2.98 (m, 4H), 2.93 (m, 1H), 2.47 (m, 1H), 1.70 (m, 2H), 1.37 (m, 4H). MS (EI) 2: 459 (MH⁺).
- [0503] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide:
15 ¹H NMR (400 MHz, d6-DMSO): 9.37 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.82 (d, 2H), 7.66 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.75 (m, 4H), 3.62 (br s, 2H), 3.32 (m, 2H), 3.05 (m, 4H). MS (EI): 405 (MH⁺).
- [0504] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)furan-2-
20 carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.44 (s, 1H), 9.40 (s, 1H), 8.46 (d, 1H), 8.16 (d, 2H), 7.98 (m, 1H), 7.93 (m, 2H), 7.68 (d, 2H), 7.39 (d, 1H), 7.30 (d, 1H), 6.93 (d, 2H), 6.74 (d, 1H), 3.75 (m, 4H), 3.05 (m, 4H). MS (EI): 3: 442 (MH⁺).
- [0505] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetra-
hydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.91 (s, 1H), 9.36 (s, 1H),
25 8.41 (d, 1H), 8.09 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.22 (d, 1H), 6.91 (d, 2H), 4.41 (dd, 1H), 3.96 (q, 1H), 3.83 (q, 1H), 3.72 (m, 4H), 3.02 (m, 4H), 2.19 (m, 1H), 1.99 (m, 1H), 1.88 (m, 2H). MS (EI): 446 (MH⁺).
- [0506] 5-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
pyrazine-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.90 (s, 1H), 9.38 (s, 1H), 9.17
(s, 1H), 8.71 (s, 1H), 8.43 (d, 1H), 8.16 (d, 2H), 8.08 (d, 2H), 7.66 (d, 2H), 7.31 (d, 1H), 6.92
30 (d, 2H), 3.71 (m, 4H), 3.03 (m, 4H), 2.62 (s, 3H). MS (EI): 468 (MH⁺).
- [0507] 2-(ethyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-
acetamide: ¹H NMR (400 MHz, d6-DMSO): 9.94 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d,
2H), 7.80 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.91 (d, 2H), 4.05 (s, 2H), 3.72 (m, 4H), 3.55
(q, 2H), 3.02 (m, 4H), 1.17 (t, 3H). MS (EI): 434 (MH⁺).
- 35 [0508] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-
(phenyloxy)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.37 (s, 1H), 9.39 (s, 1H), 8.44

5 (d, 1H), 8.14 (d, 2H), 7.82 (d, 2H), 7.67 (d, 2H), 7.31 (m, 3H), 6.98 (m, 4H), 4.75 (s, 2H), 3.73 (m, 4H), 3.04 (m, 4H). MS (EI): 482 (MH+).

[0509] methyl 4-[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-

yl}phenyl)amino]-4-oxobutanoate: ¹H NMR (400 MHz, d6-DMSO): 10.28 (s, 1H), 9.37 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.60 (s, 3H), 3.06 (m, 4H), 2.65 (m, 4H). MS (EI): 462 (MH+).

[0510] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide:

¹H NMR (400 MHz, d6-DMSO): 10.15 (s, 1H), 9.37 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.26 (d, 1H), 6.93 (d, 2H), 3.73 (m, 4H), 3.04 (m, 4H), 2.33 (t, 2H), 1.63 (q, 2H), 0.93 (t, 3H). MS (EI): 418 (MH+).

15 **[0511] 2-(2-methylphenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-**

phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.44 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.76 (d, 2H), 7.67 (d, 2H), 7.27 (m, 2H), 7.16 (m, 3H), 6.93 (d, 2H), 3.75 (m, 6H), 3.03 (m, 4H), 2.31 (m, 3H). MS (EI): 480 (MH+).

[0512] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-

20 **cyclopentane-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.14 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.75 (m, 4H), 3.04 (m, 4H), 2.80 (m, 1H), 1.87 (m, 2H), 1.72 (m, 4H), 1.55 (m, 2H). MS (EI): 444 (MH+).

[0513] (2S)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-

25 **azetidine-2-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.09 (br s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.86 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 4.32 (t, 1H), 3.73 (m, 4H), 3.62 (m, 1H), 3.06 (m, 4H), 2.58 (m, 1H), 2.29 (m, 1H), 0.99 (m, 1H). MS (EI): 431 (MH+).

[0514] N-{4-[2-({4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]phenyl}amino)pyrimidin-

30 **4-yl}phenyl]-D-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.19 (s, 1H), 9.19 (s, 1H), 8.40 (d, 1H), 8.11 (d, 2H), 7.83 (d, 2H), 7.56 (d, 2H), 7.22 (d, 1H), 6.53 (d, 2H), 3.72 (m, 1H), 3.41 (m, 1H), 3.33 (m, 1H), 3.22 (m, 1H), 3.02 (m, 1H), 2.90 (m, 2H), 2.78 (m, 1H), 2.20 (s, 6H), 2.05 (m, 2H), 1.75 (m, 2H), 1.66 (m, 2H). MS (EI): 472 (MH+).

[0515] 4-(4-aminophenyl)-N-{4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]phenyl}-

35 **pyrimidin-2-amine:** ¹H NMR (400 MHz, d6-DMSO): 8.99 (s, 1H), 8.25 (s, 1H), 7.87 (d, 2H), 7.57 (d, 2H), 7.05 (d, 1H), 6.63 (d, 2H), 6.52 (d, 2H), 5.70 (s, 2H), 3.41 (m, 1H), 3.31

5 (m, 1H), 3.21 (m, 1H), 3.01 (m, 1H), 2.77 (m, 1H), 2.20 (s, 6H), 2.15 (m, 1H), 1.77 (m, 1H). MS (EI): 375 (MH+).

[0516] N-{4-[2-({4-[(3R)-3-(dimethylamino)pyrrolidin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.19 (s, 1H), 8.40 (d, 1H), 8.10 (d, 2H), 7.75 (d, 2H), 7.58 (d, 2H), 7.23 (d, 1H), 6.54 (d, 2H), 10 3.63 (t, 2H), 3.41 (m, 1H), 3.25 (m, 3H), 3.21 (m, 1H), 3.01 (m, 1H), 2.77 (m, 1H), 2.59 (t, 2H), 2.52 (m, 1H) 2.20 (s, 6H), 2.13 (m, 1H), 1.79 (m, 1H). MS (EI): 461 (MH+).

[0517] 1-ethyl-3-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)urea: ¹H NMR (400 MHz, d6-DMSO): 9.35 (s, 1H), 8.76 (s, 1H), 8.41 (d, 1H), 8.04 (d, 2H), 7.68 (d, 2H), 7.54 (d, 2H), 7.23 (d, 1H), 6.95 (d, 2H), 6.26 (t, 15 1H), 3.59 (m, 4H), 3.56 (q, 2H), 3.23 (s, 3H), 3.14 (m, 2H), 3.07 (m, 2H), 3.01 (m, 2H), 2.61 (t, 2H), 1.06 (t, 3H). MS (EI): 504 (MH+).

[0518] 3-(methyloxy)-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)propanamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.76 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.95 (d, 20 2H), 3.60 (m, 8H), 3.25 (s, 3H), 3.23 (s, 3H), 3.08 (m, 2H), 3.01 (m, 2H), 2.61 (m, 4H). MS (EI): 519 (MH+).

[0519] N-{4-[2-({4-[4-(3-hydroxypropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 10.19 (s, 1H), 9.40 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 3.74 (dd, 25 1H), 3.66 (t, 2H), 3.62 (m, 4H), 3.09 (m, 2H), 3.03 (m, 2H), 2.91 (t, 2H), 2.52 (m, 2H), 2.05 (m, 1H), 1.79 (m, 1H), 1.66 (m, 2H). MS (EI): 516 (MH+).

[0520] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)-D-alaninamide: ¹H NMR (400 MHz, d6-DMSO): 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 3.59 (m, 30 4H), 3.57 (t, 2H), 3.46 (m, 1H), 3.23 (s, 3H), 3.07 (m, 2H), 3.02 (m, 2H), 2.61 (t, 2H), 1.24 (d, 3H). MS (EI): 504 (MH+).

[0521] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.83 (d, 2H), 7.66 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 35 3.75 (dd, 1H), 3.60 (m, 4H), 3.56 (t, 2H), 3.23 (s, 3H), 3.08 (m, 2H), 3.03 (m, 2H), 2.91 (t, 2H), 2.61 (t, 2H), 2.17 (m, 1H), 1.80 (m, 1H), 1.67 (m, 2H). MS (EI): 530 (MH+).

- 5 **[0522] N-2,N-2-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide:** ¹H NMR (400 MHz, d₆-DMSO): 9.98 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.11 (s, 2H), 3.05 (m, 4H), 2.29 (s, 6H). MS (EI): 433 (MH⁺).
- [0523] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide:**
10 ¹H NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.13 (d, 2H), 7.83 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.71 (m, 1H), 3.04 (m, 4H), 2.90 (t, 2H), 2.05 (m, 1H), 1.80 (m, 1H), 1.66 (m, 2H). MS (EI): 445 (MH⁺).
- [0524] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylpropanamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.12 (s, 1H), 9.31 (s, 1H), 8.37 (d, 1H),
15 8.05 (d, 2H), 7.67 (d, 2H), 7.60 (d, 2H), 7.21 (m, 5H), 7.12 (m, 1H), 6.83 (d, 2H), 3.67 (m, 4H), 2.98 (m, 4H), 2.86 (t, 2H), 2.61 (t, 2H). MS (EI): 480 (MH⁺).
- [0525] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylacetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.44 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.34 (m, 4H), 7.26 (m, 2H), 6.93 (d, 2H), 3.75 (m, 4H),
20 3.69 (m, 2H), 3.04 (m, 4H). MS (EI): 466 (MH⁺).
- [0526] N-[3-({4-[4-(acetyl amino)phenyl]pyrimidin-2-yl}amino)propyl]-2-fluoro-6-iodobenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.16 ppm (s, 1H), 8.64 ppm (t, 1H), 8.30 ppm (d, 1H), 8.06 ppm (d, 2H), 7.70 ppm (m, 3H), 7.30 ppm (m, 1H), 7.20 ppm (m, 1H), 7.13 ppm (m, 1H), 7.07 ppm (m, 1H), 3.34 ppm (m, 4H), 2.08 ppm (s, 3H), 1.83 ppm (m, 2H); MS (EI) C₂₂H₂₁FIN₅O₂: 533.9 (MH⁺).
25
- [0527] N-[3-({4-[4-(acetyl amino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-difluorobenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.78 ppm (s, 1H), 10.12 ppm (s, 1H), 9.70 ppm (s, 1H), 8.50 ppm (d, 1H), 8.39 ppm (s, 1H), 8.12 ppm (d, 2H), 7.73 ppm (d, 2H), 7.61 ppm (m, 1H), 7.47 ppm (m, 1H), 7.40 ppm (m, 1H), 7.27 ppm (m, 4H), 2.09 ppm (s, 3H); MS
30 (EI) C₂₅H₁₉F₂N₅O₂: 460 (MH⁺).
- [0528] N-[3-({4-[4-(acetyl amino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,4,5-trifluorobenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.52 ppm (s, 1H), 10.29 ppm (s, 1H), 9.71 ppm (s, 1H), 8.50 ppm (d, 1H), 8.36 ppm (s, 1H), 8.20 ppm (d, 2H), 7.87 ppm (m, 1H), 7.75 ppm (d, 3H), 7.51 ppm (m, 1H), 7.37 ppm (d, 1H), 7.28 ppm (m, 2H), 2.09 ppm (s, 3H); MS (EI) C₂₅H₁₈F₃N₅O₂: 478 (MH⁺).
35
- [0529] N-[3-({4-[4-(acetyl amino)phenyl]pyrimidin-2-yl}amino)phenyl]benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.27 ppm (s, 1H), 10.21 ppm (s, 1H), 9.67 ppm (s, 1H),

5 8.51 ppm (d, 1H), 8.47 ppm (s, 1H), 8.13 ppm (d, 2H), 8.00 ppm (m, 2H), 7.75 ppm (m, 2H), 7.58 ppm (m, 3H), 7.48 ppm (m, 1H), 7.37 ppm (d, 1H), 7.29 ppm (m, 2H), 2.09 ppm (s, 3H); MS (EI) C₂₅H₂₁N₅O₂: 424(MH⁺).

[0530] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3,5-difluoro-
benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.37 ppm (s, 1H), 10.20 ppm (s, 1H), 9.70
10 ppm (s, 1H), 8.51 ppm (d, 1H), 8.41 ppm (s, 1H), 8.21 ppm (d, 2H), 7.73 ppm (m, 4H), 7.55
ppm (m, 2H), 7.37 ppm (d, 1H), 7.29 ppm (m, 2H), 2.08 ppm (s, 3H); MS (EI)
C₂₅H₁₉F₂N₅O₂: 468.0 (MH⁺).

[0531] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-
fluoro-4-methylbenzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.72 ppm (s, 1H), 10.20
15 ppm (s, 1H), 9.69 ppm (s, 1H), 8.50 ppm (d, 1H), 8.41 ppm (s, 1H), 8.21 ppm (d, 2H), 7.74
ppm (d, 2H), 7.50 ppm (m, 2H), 7.37 ppm (d, 1H), 7.28 ppm (m, 8H), 2.38 ppm (s, 3H), 2.08
ppm (s, 3H); MS (EI) C₂₆H₂₁ClFN₅O₂: 490.0 (MH⁺).

[0532] N-(4-{2-[(3-[(2,6-dimethylphenyl)methyl]amino)phenyl]amino}pyrimidin-4-
yl)-phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.20 ppm (s, 1H), 9.36 ppm (s,
20 1H), 8.47 ppm (d, 1H), 8.16 ppm (d, 2H), 7.72 ppm (d, 2H), 7.35 ppm (s, 1H), 7.31 ppm (d,
1H), 7.12 ppm (m, 1H), 7.07 ppm (m, 2H), 6.99 ppm (m, 3H), 6.38 ppm (d, 1H), 5.46 ppm (t,
1H), 4.14 ppm (d, 2H), 2.36 ppm (s, 6H), 2.08 ppm (s, 3H); MS (EI) C₂₇H₂₇N₅O: 438.1
(MH⁺).

[0533] N-(3-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-
25 carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.66 ppm (s, 1H), 10.28 ppm (s, br, 2H),
10.10 ppm (s, 1H), 8.74 ppm (s, 1H), 8.63 ppm (d, 1H), 8.25 ppm (m, 2H), 7.94 ppm (m,
3H), 7.75 ppm (t, 1H), 7.55 ppm (t, 1H), 7.45 ppm (m, 2H), 7.26 ppm (m, 1H), 6.97 ppm
(m, 1H); MS (EI) C₂₁H₁₇N₅O₂S: 388.0 (MH⁺).

[0534] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-1-
30 methylpiperidine-4-carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 ppm (s, 1H),
9.84 ppm (s, 1H), 9.60 ppm (s, 1H), 8.49 ppm (d, 1H), 8.29 ppm (s, 1H), 8.20 ppm (d, 2H),
7.74 ppm (d, 2H), 7.36 ppm (m, 2H), 7.18 ppm (m, 2H), 2.84 ppm (m, 2H), 2.31 ppm (m,
1H), 2.17 ppm (s, 3H), 2.09 ppm (s, 3H), 1.87 ppm (m, 2H), 1.72 ppm (m, 4H); MS (EI)
C₂₅H₂₈N₅O₂: 445 (MH⁺).

35 [0535] 4-(4-aminophenyl)-N-[3-(morpholin-4-ylsulfonyl)phenyl]pyrimidin-2-amine:
¹H-NMR (400MHz, d₆-DMSO): 9.94 ppm (s, 1H), 8.76 ppm (s, 1H), 8.42 ppm (d, 1H), 7.98

5 ppm (d, dH), 7.90 ppm (m, 1H), 7.57 ppm (t, 1H), 7.28 ppm (m, 2H), 6.65 ppm (d, 2H), 5.81 ppm (s, 2H), 3.64 ppm (m, 4H), 2.90 ppm (m, 4H); MS (EI) $C_{20}H_{21}N_5O_3S$: 412 (MH^+).

[0536] N-(4-{2-[(3-[(2-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 ppm (s, 1H), 9.35 ppm (s,

10 1H), 8.45 ppm (d, 1H), 8.12 ppm (d, 2H), 7.74 ppm (d, 2H), 7.43 ppm (m, 2H), 7.29 ppm (m, 2H), 7.16 ppm (m, 3H), 6.99 ppm (m, 2H), 6.21 ppm (m, 2H), 4.33 ppm (d, 2H), 2.09 ppm (s, 3H); MS (EI) $C_{25}H_{22}FN_5O$: 428 (MH^+).

[0537] N-(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 ppm (s, 1H), 9.31 ppm (s, 1H), 8.46 ppm (d, 1H), 8.13 ppm (d, 2H), 7.75 ppm (d, 2H), 7.30 ppm (d, 1H), 7.13 ppm (s, 1H), 6.94 ppm (m, 2H), 6.20 ppm (d, 15 1H), 5.01 ppm (s, 2H), 2.10 ppm (s, 3H); MS (EI) $C_{18}H_{17}IN_5O$: 320 (MH^+).

[0538] N-(4-{2-[(3-[(4-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}-phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.26 ppm (s, 1H), 9.33 ppm (s, 1H), 8.45 ppm (d, 1H), 8.13 ppm (d, 2H), 7.75 ppm (d, 2H), 7.37 ppm (m, 2H), 7.31 ppm (m, 3H), 7.21 ppm (m, 2H), 6.97 ppm (m, 2H), 6.23 ppm (m, 2H), 4.48 ppm (d, 2H), 2.09 ppm (s, 3H); 20 MS (EI) $C_{25}H_{23}N_5O$: 410 (MH^+).

[0539] N-(4-{2-[(3-[(3-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}-phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.36 ppm (s, 1H), 9.35 ppm (s, 1H), 8.45 ppm (d, 1H), 8.12 ppm (d, 2H), 7.76 ppm (d, 2H), 7.33 ppm (m, 2H), 7.19 ppm (m, 3H), 6.99 ppm (m, 3H), 6.32 ppm (t, 1H), 6.20 ppm (m, 2H), 4.30 ppm (d, 2H), 2.09 ppm (s, 3H); 25 MS (EI) $C_{25}H_{22}FN_5O$: 428 (MH^+).

[0540] N-(4-{2-[(3-[(4-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}-phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.20 ppm (s, 1H), 9.33 ppm (s, 1H), 8.45 ppm (d, 1H), 8.12 ppm (d, 2H), 7.74 ppm (d, 2H), 7.39 ppm (m, 2H), 7.30 ppm (d, 1H), 7.21 ppm (s, 1H), 7.13 ppm (t, 2H), 6.96 ppm (m, 2H), 6.22 ppm (m, 2H), 2.09 ppm (s, 3H); 30 MS (EI) $C_{25}H_{22}FN_5O$: 428 (MH^+).

[0541] 4-[4-({4-[4-(butanoylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-ethyl-piperazine-1-carboxamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 ppm (s, 1H), 9.39 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.77 ppm (d, 2H), 7.68 ppm (d, 2H), 7.28 ppm (d, 1H), 6.96 ppm (d, 2H), 6.59 ppm (t, 1H), 3.43 ppm (t, 4H), 3.07 ppm (m, 2H), 3.02 ppm 35 (t, 4H), 2.34 ppm (t, 2H), 1.63 ppm (m, 2H), 1.02 ppm (t, 3H), 0.98 ppm (t, 3H); MS (EI) $C_{27}H_{33}N_7O_2$: 488 (MH^+).

- 5 **[0542] N-{4-[2-({4-[4-(2-piperazin-1-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.18 ppm (s, 1H), 9.40 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.77 ppm (d, 2H), 7.68 ppm (d, 2H), 7.28 ppm (d, 1H), 6.97 ppm (d, 2H), 3.72 ppm (m, 2H), 3.59 ppm (m, 2H), 3.13 ppm (m, 4H), 3.01 ppm (m, 2H), 2.67 ppm (m, 4H), 2.33 ppm (m, 4H), 1.63 ppm (m, 2H), 0.93 ppm (t, 3H); MS (EI) C₃₀H₃₈N₈O₂: 543 (MH⁺).
- 10 **[0543] N-[4-(2-{[4-(4-L-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-butanamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.15 ppm (s, 1H), 9.40 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.76 ppm (d, 2H), 7.69 ppm (d, 2H), 7.28 ppm (d, 1H), 6.97 ppm (d, 2H), 3.79 ppm (m, 1H), 3.62 ppm (m, 4H), 3.06 ppm (m, 4H), 2.33 ppm (t, 2H), 1.91 ppm (br. s, 2H), 1.63 ppm (m, 2H), 1.09 ppm (d, 3H), 0.93 ppm (t, 3H); MS (EI) C₂₇H₃₃N₇O₂: 488 (MH⁺).
- 15 **[0544] N-[4-(2-{[4-(4-L-prolylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-butanamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 ppm (s, 1H), 9.41 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.77 ppm (d, 2H), 7.69 ppm (d, 2H), 7.28 ppm (d, 1H), 6.97 ppm (d, 2H), 3.85 ppm (m, 1H), 3.62 ppm (m, 4H), 3.04 ppm (m, 5H), 2.62 ppm (m, 1H), 2.34 ppm (t, 2H), 2.00 ppm (m, 1H), 1.62 ppm (m, 6H), 0.93 ppm (t, 3H); MS (EI) C₂₉H₃₅N₇O₂: 514 (MH⁺).
- 20 **[0545] (3R)-1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrrolidine-3-carboxamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 ppm (s, 1H), 9.38 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.76 ppm (d, 2H), 7.67 ppm (d, 2H), 7.28 ppm (d, 1H), 6.94 ppm (d, 2H), 4.50 ppm (m, 1H), 3.75 ppm (m, 4H), 3.49 ppm (m, 2H), 3.05 ppm (m, 6H), 2.91 ppm (t, 1H), 2.67 ppm (m, 1H), 2.56 ppm (m, 3H), 1.98 ppm (m, 2H); MS (EI) C₂₇H₃₂N₆O₃: 489 (MH⁺).
- 25 **[0546] N-(4-{2-[(3-fluoro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 ppm (s, 1H), 9.69 ppm (s, 1H), 8.51 ppm (d, 1H), 8.13 ppm (d, 2H), 7.85 ppm (d, 2H), 7.79 ppm (m, 1H), 7.52 ppm (m, 1H), 7.37 ppm (d, 1H), 7.03 ppm (t, 1H), 3.74 ppm (m, 5H), 3.14 ppm (br.s, 1H), 2.95 ppm (m, 4H), 2.91 ppm (m, 2H), 2.06 ppm (m, 1H), 1.80 ppm (m, 1H), 1.66 ppm (m, 2H); MS (EI) C₂₅H₂₇FN₆O₂: 463 (MH⁺).
- 30 **[0547] N-(4-{2-[(3-fluoro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide:** ¹H-NMR (400MHz, d₆-DMSO): 9.69 ppm (s, 1H), 8.51 ppm (d, 1H), 8.13 ppm (d, 2H), 7.84 ppm (d, 2H), 7.79 ppm (m, 1H), 7.54 ppm (m, 1H), 7.37 ppm (m, 1H),
- 35

5 7.37 ppm (m, 1H), 7.03 ppm (t, 1H), 3.74 ppm (m, 4H), 3.47 ppm (m, 1H), 2.95 ppm (m, 4H), 1.23 ppm (d, 3H); MS (EI) $C_{23}H_{25}FN_6O_2$: 437 (MH⁺).

[0548] N-(4-{2-[(3-methyl-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 ppm (s, 1H), 9.45 ppm (s, 1H), 8.46 ppm (d, 1H), 8.14 ppm (d, 2H), 7.85 ppm (d, 2H), 7.63 ppm (d, 2H), 7.32 ppm (d, 1H), 7.32 ppm (d, 1H), 3.73 ppm (m, 5H), 3.08 ppm (br.s., 1H), 2.90 ppm (t, 2H), 2.80 ppm (m, 4H), 2.27 ppm (s, 3H), 2.06 ppm (m, 1H), 1.80 ppm (m, 1H), 1.66 ppm (m, 2H); MS (EI) $C_{26}H_{30}N_6O_2$: 459 (MH⁺).

[0549] N-(4-{2-[(3-methyl-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide: ¹H-NMR (400MHz, d₆-DMSO): 9.44 ppm (s, 1H), 8.47 ppm (d, 1H), 8.14 ppm (d, 2H), 7.83 ppm (d, 2H), 7.64 ppm (m, 2H), 7.32 ppm (d, 1H), 7.02 ppm (m, 1H), 3.73 ppm (m, 4H), 3.46 ppm (m, 1H), 2.80 ppm (m, 4H), 2.28 ppm (m, 4H), 1.23 ppm (d, 3H); MS (EI) $C_{24}H_{28}N_6O_2$: 433 (MH⁺).

[0550] 1-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclopropanecarboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.14 ppm (s, 1H), 9.39 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.94 ppm (d, 2H), 7.68 ppm (d, 2H), 7.29 ppm (d, 1H), 6.94 ppm (d, 2H), 6.81 ppm (s, 1H), 3.74 ppm (m, 4H), 3.05 ppm (m., 4H), 1.18 ppm (m, 2H), 1.00 ppm (m, 2H); MS (EI) $C_{24}H_{25}N_5O_3$: 432 (MH⁺).

[0551] N-(4-(2-(4-(4-(4-chloro-2,6-dimethylphenylsulfonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.42 (s, 1H), 8.41 (m, 1H), 8.08 (d, 2H), 7.77 (s, 1H), 7.71 (d, 2H), 7.64 (m, 2H), 7.57 (s, 1H), 7.27 (m, 1H), 6.93 (m, 2H), 3.41 (m, 4H), 3.15 (m, 4H), 2.53 (s, 3H), 2.37 (s, 3H), 2.06 (s, 3H). MS (EI): 591 (MH⁺).

[0552] N-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.96 (t, 1H), 3.75 (m, 5H), 3.60 (m, 2H), 3.32 (m, 1H), 3.19 (m, 1H), 3.12 (s, 2H), 3.10 (m, 2H), 3.01 (m, 2H), 2.68 (m, 4H), 2.32 (m, 4H), 2.10 (m, 2H). MS (EI): 571 (MH⁺).

[0553] N-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.30 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.96 (t, 1H), 3.74 (m, 6H), 3.19 (m, 2H), 3.05 (m, 4H), 2.10 (q, 2H), 1.23 (s, 9H). MS (EI): 529 (MH⁺).

- 5 **[0554] 1-ethyl-3-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea:** ¹H NMR (400 MHz, d6-DMSO): 9.24 (s, 1H), 8.87 (s, 1H), 8.29 (s, 1H), 7.64 (d, 2H), 7.59 (m, 2H), 7.53 (m, 2H), 6.88 (d, 2H), 6.41 (m, 1H), 3.73 (m, 4H), 3.12 (m, 2H), 3.02 (m, 4H), 2.23 (s, 3H), 1.06 (t, 3H). MS (EI): 433 (MH⁺).
- [0555] 3-methoxy-N-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide:** ¹H NMR (400 MHz, d6-DMSO): 10.17 (s, 1H), 9.30 (s, 1H), 8.31 (s, 1H), 7.74 (m, 2H), 7.66 (m, 4H), 6.90 (m, 2H), 3.74 (m, 4H), 3.64 (t, 2H), 3.26 (s, 3H), 3.03 (m, 4H), 2.59 (m, 2H), 2.22 (s, 3H). MS (EI): 448 (MH⁺).
- 10 **[0556] N-(4-(2-(4-(4-(ethylsulfonyl)piperazine-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.38 (s, 1H), 8.42 (d, 1H), 8.08 (d, 2H), 7.72 (d, 2H), 7.66 (d, 2H), 7.26 (d, 1H), 6.95 (d, 2H), 3.36 (m, 4H), 3.12 (m, 4H), 2.48 (m, 2H), 2.07 (s, 3H), 1.22 (t, 3H). MS (EI): 481 (MH⁺).
- 15 **[0557] 4-(4-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-N-ethylpiperazine-1-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.41 (d, 1H), 8.08 (d, 2H), 7.72 (d, 2H), 7.64 (d, 2H), 7.25 (d, 1H), 6.94 (d, 2H), 6.57 (d, 1H), 3.50 (m, 2H), 3.39 (m, 4H), 2.99 (m, 4H), 2.07 (s, 3H), 1.00 (t, 3H). MS (EI): 460 (MH⁺).
- 20 **[0558] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)morpholine-2-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 9.87 (s, 1H), 9.37 (s, 1H), 8.43 (s, 1H), 8.04-8.16 (d, 2H), 7.81-7.93 (d, 2H), 7.60-7.72 (d, 2H), 7.27 (s, 1H), 6.85-6.99 (d, 2H), 4.08-4.69 (s, br, 1H), 4.00-4.07 (d, 1H), 3.85-3.94 (d, 1H), 3.72 (s, 3H), 3.51-3.63 (d, 1H), 2.58-2.80 (m, 3H), 1.86 (s, 6H). MS (EI): 461 (MH⁺).
- 25 **[0559] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-beta-alaninamide:** ¹H NMR (400 MHz, d6-DMSO): 9.38 (s, 1H), 10.65 (d, 1H), 8.07-8.16 (d, 2H), 7.72-7.81 (d, 2H), 7.62-7.72 (d, 2H), 7.28 (s, 1H), 6.89-6.98 (d, 2H). MS (EI): 419 (MH⁺).
- 30 **[0560] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)phenylalaninamide:** ¹H NMR (400 MHz, d6-DMSO): 9.39 (s, 1H), 8.42-8.46 (d, 1H), 8.09-8.14 (d, 2H), 7.75-7.81 (d, 2H), 7.64-7.70 (d, 2H), 7.23-7.32 (m, 6H), 7.16-7.22 (m, 2H), 6.90-6.97 (d, 2H), 3.71-3.78 (m, 4H), 3.57-3.63 (m, 1H), 3.02-3.08 (m, 4H), 2.98-3.02 (m, 1H), 2.71-2.79 (m, 1H). MS (EI): 495 (MH⁺).
- 35 **[0561] N²-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-glycinamide:** ¹H NMR (400 MHz, d6-DMSO): 9.39 (s, 1H), 8.45 (s, 1H), 8.09-8.17 (d, 2H),

5 7.78-7.86 (d, 2H), 7.65-7.73 (d, 2H), 7.29 (s, 1H), 6.90-7.00 (d, 2H), 3.74 (s, 4H), 3.05 (s, 4H), 2.33 (s, 3H), 1.92 (s, 1H). MS (EI): 419 (MH⁺).

[0562] 2-cyclopentyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.13 (s, 1H), 9.37 (s, 1H), 8.42-8.45 (d, 1H), 8.07-8.14 (d, 2H), 7.73-7.79 (d, 2H), 7.65-7.71 (d, 2H), 7.25-7.28 (d, 1H), 6.90-10 6.97 (d, 2H), 3.71-3.77 (m, 4H), 3.02-3.07 (m, 4H), 2.33-2.37 (d, 2H), 2.20-2.30 (m, 1H), 1.71-1.82 (m, 2H), 1.48-1.66 (m, 4H), 1.14-1.25 (m, 2H). MS (EI): 458 (MH⁺).

[0563] 6-(methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.50 (s, 1H), 9.44 (s, 1H), 8.80-8.83 (d, 1H), 8.44-8.49 (d, 1H), 8.24-8.28 (m, 1H), 8.15-8.21 (d, 2H), 7.92-7.97 (d, 15 2H), 7.67-7.72 (d, 2H), 7.30-7.34 (d, 1H), 6.93-7.02 (m, 3H), 3.94-3.96 (s, 3H), 3.72-3.79 (m, 4H), 3.04-3.11 (m, 4H). MS (EI): 483 (MH⁺).

[0564] N,N-dimethyl-N'-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanediamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.36 (s, 1H), 8.43 (s, 1H), 8.05-8.17 (d, 2H), 7.71-7.79 (d, 2H), 7.61-7.71 (d, 2H), 7.26-7.31 (d, 1H), 6.89-20 7.00 (d, 2H), 3.68-3.79 (m, 4H), 3.02-3.08 (m, 4H), 3.00 (s, 3H), 2.82 (s, 3H), 2.56-2.66 (m, 4H). MS (EI): 475 (MH⁺).

[0565] N-[4-(2-{[4-morpholin-4-yl-3-(trifluoromethyl)phenyl]amino} pyrimidin-4-yl)-phenyl]-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.94 (s, 1H), 8.53-8.57 (d, 1H), 8.48 (s, 1H), 8.14-8.21 (d, 2H), 7.93-7.98 (m, 1H), 7.82-7.88 (d, 2H), 7.56-7.61 25 (d, 1H), 7.42-7.46 (d, 1H), 3.79-3.86 (m, 1H), 3.67-3.75 (m, 4H), 2.93-3.00 (m, 2H), 2.79-2.86 (m, 4H), 2.05-2.17 (m, 1H), 1.79-1.89 (m, 1H), 1.65-1.75 (m, 2H). MS (EI): 513 (MH⁺).

[0566] 3-(methyloxy)-N-[4-(2-{[4-morpholin-4-yl-3-(trifluoromethyl)phenyl]amino} pyrimidin-4-yl)phenyl]propanamide: ¹H NMR (400 MHz, d₆-DMSO): 10.24 (s, 1H), 9.94 (s, 1H), 8.53-8.56 (d, 1H), 8.47 (s, 1H), 8.13-8.19 (d, 2H), 7.94-8.00 (d, 1H), 7.76-7.81 (d, 30 1H), 7.56-7.62 (d, 1H), 7.41-7.45 (d, 1H), 3.67-3.74 (m, 4H), 3.60-3.67 (m, 2H), 3.35 (s, 3H), 2.80-2.86 (d, 4H), 2.57-2.63 (m, 2H). MS (EI): 502 (MH⁺).

[0567] N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)-5-oxo-L-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 10.34 (s, 1H), 9.37 (s, 1H), 8.43-8.46 (d, 1H), 8.12-8.16 (d, 2H), 7.94 (s, 1H), 7.77-7.81 (d, 35 2H), 7.62-7.67 (d, 2H), 7.26-7.30 (d, 1H), 6.88-6.94 (d, 2H), 4.20-4.26 (m, 1H), 3.02-3.08 (m, 4H), 2.57-2.64 (m, 4H), 2.21 (s, 6H), 2.17 (s, 2H), 2.10 (s, 2H), 1.89 (s, 4H), 0.84 (s, 6H). MS (EI): 571 (MH⁺).

5 **[0568] (2R)-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)tetrahydrofuran-2-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 9.94 (s, 1H), 9.41 (s, 1H), 8.43-8.46 (d, 1H), 8.09-8.15 (d, 2H), 7.85-7.90 (d, 2H), 7.65-7.71 (d, 1H), 6.92-6.98 (d, 2H), 4.39-4.48 (m, 1H), 3.95-4.05 (m, 1H), 3.79-3.89 (m, 1H), 3.52-3.64 (m, 6H), 3.32 (s, 1H), 3.23 (s, 2H), 2.98-3.11 (m, 4H), 2.58-2.65 (m, 2H), 2.15-2.25 (m, 10 1H), 1.96-2.07 (m, 1H), 1.83-1.93 (m, 2H). MS (EI): 531 (MH+).

[0569] (2S)-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.94 (s, 1H), 9.41 (s, 1H), 8.42-8.47 (d, 2H), 8.09-8.16 (d, 2H), 7.84-7.91 (d, 2H), 7.64-7.72 (d, 2H), 7.27-7.37 (d, 1H), 6.93-6.99 (d, 2H), 4.40-4.47 (m, 1H), 3.95-4.05 (m, 1H), 3.80-15 3.89 (m, 1H), 3.53-3.65 (m, 6H), 3.32 (s, 1H), 3.23 (s, 2H), 2.98-3.10 (m, 4H), 2.59-2.65 (m, 2H), 2.15-2.27 (m, 1H), 1.95-2.07 (m, 1H), 1.83-1.93 (m, 2H). MS (EI): 531 (MH+).

[0570] (2R,4S)-4-hydroxy-N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenyl)amino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.42-8.46 (d, 1H), 8.09-8.15 (d, 2H), 7.80-7.86 (d, 2H), 20 7.65-7.71 (d, 2H), 7.27-7.31 (d, 1H), 6.93-6.98 (d, 1H), 4.20-4.26 (m, 1H), 3.88-3.94 (m, 1H), 3.53-3.64 (m, 6H), 3.17 (s, 1H), 2.99-3.10 (m, 4H), 2.89-2.93 (m, 1H), 2.77-2.84 (m, 1H), 2.58-2.64 (m, 3H), 1.99-2.07 (m, 2H), 1.73-1.83 (m, 2H). MS (EI): 546 (MH+).

[0571] N-(4-{2-[(3-fluoro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 11.07 (s, 1H), 9.89 (s, 1H), 8.46-8.56 (d, 25 1H), 8.08-8.21 (d, 2H), 7.74-7.91 (m, 3H), 7.47-7.57 (d, 1H), 7.35-7.41 (d, 1H), 6.98-7.08 (m, 1H), 3.70-3.82 (m, 5H), 2.87-3.03 (m, 5H), 2.01-2.16 (m, 1H), 1.92 (s, 2H), 1.75-1.87 (m, 1H), 1.61-1.74 (m, 2H). MS (EI): 463 (MH+).

[0572] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 30 MHz, d6-DMSO): 10.30 (s, 1H), 9.40 (s, 1H), 8.42-8.47 (d, 1H), 8.09-8.15 (d, 2H), 7.74-7.80 (d, 2H), 7.64-7.71 (d, 2H), 7.27-7.30 (d, 1H), 6.92-7.00 (d, 2H), 3.92-3.99 (m, 1H), 3.68-3.84 (m, 4H), 3.53-3.64 (m, 5H), 3.32 (s, 1H), 3.23 (s, 2H), 3.16-3.22 (m, 1H), 2.98-3.10 (m, 4H), 2.59-2.65 (m, 1H), 2.06-2.15 (m, 2H). MS (EI): 531 (MH+).

[0573] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-pyridin-3-ylpropanamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.37 (s, 1H), 8.39-8.51 (m, 35 3H), 8.08-8.13 (d, 2H), 7.64-7.76 (m, 5H), 7.25-7.35 (m, 2H), 6.91-6.98 (d, 2H), 3.72-3.77 (m, 4H), 3.01-3.08 (m, 4H), 2.91-2.98 (m, 2H), 2.65-2.75 (m, 2H). MS (EI): 481 (MH+).

- 5 **[0574] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2-chlorobenzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.497(s, 1H), 10.201 (s, 1H), 9.668 (s, 1H), 8.505 (d, 1H), 8.505 (d, 1H), 8.427 (s, 1H), 8.223 (d, 2H), 7.748 (d, 2H), 7.59 (m, 2H), 7.477 (m, 3H), 7.374 (d, 1H), 7.253 (m, 2H), 2.083 (s, 3H). MS (EI): 458 (MH+).
- [0575] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2-methylbenzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.284(d, 2H), 9.622 (s, 1H), 8.487(m, 2H), 8.235(d, 2H), 7.749 (d, 2H), 7.352(m, 8H), 2.084(s, 3H). MS (EI): 438 (MH+).
- 10 **[0576] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2,4-dichlorobenzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.533 (s, 1H), 10.198(s, 1H), 9.687(s, 1H), 8.506(d, 1H), 8.406(s, br, 1H), 8.220(d, 2H), 7.787(d, 1H), 7.747(d, 2H), 7.659(d, 1H), 7.591(d, 1H), 7.464 (d, 1H), 7.377(d, 1H), 7.247(m, 2H), 2.085(s, 3H). MS (EI): 492 (MH+).
- 15 **[0577] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2,5-dichlorobenzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.510(s, 1H), 10.198(s, 1H), 9.696(s, 1H), 8.507(s, 1H), 8.389(s, 1H), 8.220(d, 2H), 7.744(m, 3H), 7.617(m, 2H), 7.487(d, 1H), 7.379(d, 1H), 7.282(m, 2H), 2.081(s, 3H). MS (EI): 492 (MH+).
- 20 **[0578] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2-chloro-6-fluoro-3-methoxybenzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.726(s, 1H), 10.204(s, 1H), 9.709(s, 1H), 8.509(d, 1H), 8.402(s, 1H), 8.217(d, 2H), 7.743(d, 2H), 7.488(d, 1H), 7.386(m, 2H), 7.282(m, 2H), 7.221(d, 1H), 3.904(s, 3H), 2.082(s, 3H). MS (EI): 506 (MH+).
- 25 **[0579] N-(3-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)-2,3-dichlorobenzamide:** ¹H-NMR(400MHz, d6-DMSO): 10.6 (s,1H), 10.2(s, 1H), 9.7(s, 1H), 8.5(d, 1H), 8.4(s, 1H), 8.2(d, 2H), 7.8(m, 3H), 7.6(d, 1H), 7.5(m, 2H), 7.4(d, 1H), 7.2(m, 2H), 2.081(s,3H). MS (EI): 492 (MH+).
- [0580] (R)-N-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)-pyrimidin-4-yl) phenyl)cyclopropanecarboxamide:** ¹H-NMR (400MHz, d6-DMSO): 10.476(s, 1H), 9.406(s, 1H), 8.448(d, 1H), 8.124(d, 2H), 7.767(m, 4H), 7.287(d, 1H), 6.977(d, 2H), 3.85(m, 1H), 3.65(m, 4H), 3.0(m, 4H), 2.95(m, 1H), 2.6(m, 1H), 2.0(m, 1H), 1.8(m, 1H), 1.613(m, 3H), 0.84(m, 4H). MS (EI): 512 (MH+).
- 30 **[0581] N-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide:** ¹H-NMR (400MHz, d6-DMSO): 10.480(s, 1H), 9.399(s,1H), 8.447(d, 1H), 8.124(d, 2H), 7.769(d, 2H), 7.692(d, 2H), 7.285(d, 1H), 6.975(d,
- 35

5 2H), 3.170(m, 2H), 3.592(m, 2H), 3.134(s, 2H), 3.099(m, 2H), 3.028(m, 2H), 2.694(m, 4H), 2.331(m, 4H), 0.842(m, 4H). MS (EI): 541 (MH+).

[0582] 4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzamide: ¹H-NMR (400MHz, d6-DMSO): 9.5(s, 1H), 8.5(d, 1H), 8.2(d, 2H), 8.15(s, 1H), 8(d, 2H), 7.7(d, 1h), 7.5(s, 1H), 7.4(d, 1H), 6.9(d, 1H), 4.8(m, 4h), 3.0(m, 4H). MS (EI): 376 (MH+).

10 **[0583] (R)-N-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide:** ¹H-NMR (400MHz, d6-MeOD): 8.355(d, 1H), 8.114(d, 2H), 7.177(d, 2H), 7.639(d, 2H), 7.217(d, 1H), 7.028(d, 2H), 4.394(m, 1H), 3.797 (m, 2H), 3.692(m, 2H), 3.137(m, 4H), 3.1(m, 1H), 2.410(m, 1H), 1.992(m, 2H), 1.827(m, 2H), 0.981(m, 2H), 0.85(m, 2H). MS (EI): 512 (MH+).

15 **[0584] (S)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide:** ¹H-NMR (400MHz, MeOD): 8.357(d, 1H), 8.116(d, 2H), 7.718(d, 2H), 7.642(d, 2H), 7.221(d, 1H), 7.033(d, 2H), 4.1(m, 1H), 3.85(m, 1H), 3.7(m, 4H), 3.2(m, 4H), 1.8(m, 1H), 1.4(d, 3H), 0.9(m, 2H), 0.85(m, 2H). MS (EI): 486 (MH+).

20 **[0585] (R)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide:** ¹H-NMR (400MHz, MeOD): 8.4(d, 1H), 8.15(d, 2H), 7.8(d, 2H), 7.6(d, 2H), 7.2(d, 1H), 7.0(d, 2H), 4.0(m, 1H), 3.7(m, 4H), 3.2(m, 4H), 1.8(m, 1H), 1.3(d, 3H), 0.9(m, 2H), 0.85(m, 2H). MS (EI): 486 (MH+).

[0586] (R)-N-(4-(2-(4-(4-(2-acetylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide: ¹H-NMR (400MHz, d6-DMSO): 10.193(s, 1H), 9.411(s, 1H), 8.452-8.439(d, 1H), 8.136-8.144(d, 2H), 7.849-7.828 ((d, 2H), 7.690-7.688(d, 2H), 7.302-7.289(d, 1H), 6.971-6.948(d, 2H), 3.743(m, 1H), 3.588(m, 4H), 3.085-3.016(m, 4H), 2.905(t, 2H), 2.046(s, 3H), 1.808(m, 1H), 1.663(m, 2H). MS (EI): 486 (MH+).

[0587] (R)-N-(4-(2-(4-(4-(2-methoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide: ¹H-NMR (400MHz, d6-DMSO): 10.198(s, 1H), 9.414(s, 1H), 8.452(d, 1H), 8.136(d, 2H), 7.850(d, 2H), 7.691(d, 2H), 7.302(d, 1H), 6.970(d, 2H), 4.136(s, 2H), 3.744(m, 1H), 3.599-3.535(m, 4H), 3.302(s, 3H), 3.078(m, 4H), 2.905(t, 2H), 2.079(m, 1H), 1.791(m, 1H), 1.646((m, 2H). MS (EI): 516 (MH+).

[0588] N-{1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyrrolidin-3-yl}acetamide: NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.20 (s, 1H), 8.60 (s, 1H), 8.15-8.20 (m, 3H), 7.79-7.86 (m, 4H), 7.20 (s, 1H), 6.58 (d, 2H), 4.39 (m, 1H), 3.43 (m, 1H),

5 3.23 (m, 1H), 3.10 (m, 1H), 2.18 (m, 1H), 2.07 (s, 3H), 1.85 (m, 1H), 1.80 (s, 3H). MS (EI): 431 (MH⁺).

[0589] N-[4-(2-([4-(3-oxopiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide: NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.41 (s, 1H), 8.06 (d, 2H), 8.02 (s, 1H), 7.65 – 7.80 (m, 4H), 7.25 (s, 1H), 6.97 (d, 2H), 3.64 (s, 2H), 3.35 –
10 3.40 (m, 4H), 2.05 (s, 3H). MS (EI): 403 (MH⁺).

[0590] ethyl N-[4-([4-(4-(acetylamino)phenyl)pyrimidin-2-yl]amino)phenyl]-N-methylglycinate: MS (EI): 420 (MH⁺).

[0591] ethyl 1-[4-([4-(4-(acetylamino)phenyl)pyrimidin-2-yl]amino)phenyl]piperidine-3-carboxylate: NMR (400 MHz, d₆-DMSO): 10.40 (s, 1H),
15 10.00 (s, 2H), 8.65 (d, 1H), 8.14 (d, 2H), 7.78 (d, 2H), 7.50 – 7.62 (m, 4H), 7.40 (d, 1H), 2.09 (s, 3H), 2.00 (s, 3H). MS (EI): 362 (MH⁺).

[0592] ethyl 1-[4-([4-(4-(acetylamino)phenyl)pyrimidin-2-yl]amino)phenyl]piperidine-3-carboxylate: MS (EI): 460 (MH⁺).

[0593] N-(4-{2-([4-{bis[2-(methyloxy)ethyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for C₂₄H₂₉N₅O₃: 436 (MH⁺).
20

[0594] N-[4-(2-([4-(morpholin-4-ylsulfonyl)phenyl]amino)pyrimidin-4-yl)phenyl]-acetamide: MS (EI) for C₂₂H₂₃N₅O₄S: 454 (MH⁺).

[0595] 3-hydroxy-3-methyl-N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]butanamide: NMR (400 MHz, d₆-DMSO): 10.04 (s, 1H), 9.46 (s,
25 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.65 (s, 1H), 7.29 (d, 1H), 6.91 (d, 1H), 3.79 (s, 3H), 3.68 (m, 4H), 2.89 (m, 4H), 2.44 (s, 2H), 1.23 (s, 6H). MS (EI) for C₂₆H₃₁N₅O₄: 478 (MH⁺).

[0596] 1-methyl-N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-D-prolinamide: NMR (400 MHz, d₆-DMSO): 10.0 (s, 1H), 9.44 (s, 1H), 8.42
30 (d, 1H), 8.18 (d, 2H), 7.82 (d, 2H), 7.62 (s, 1H), 7.30 (m, 2H), 6.81 (d, 1H), 3.80 (s, 3H), 3.68 (m, 4H), 2.85 – 3.10 (m, 6H), 2.37 – 2.48 (m, 5H), 2.20 (m, 1H), 1.80 (m, 2H). MS (EI) for C₂₇H₃₂N₆O₃: 489 (MH⁺).

[0597] N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-D-alaninamide: NMR (400 MHz, d₆-DMSO): 11.40 (s, 1H), 10.10 (s, 1H), 8.57
35 (d, 1H), 8.45 (d, 2H), 8.02 (d, 2H), 7.87 (m, 3H), 7.47 (m, 2H), 4.15 (m, 1H), 3.95 – 4.10 (m, 7H), 3.58 (m, 4H), 1.48 (d, 3H). MS (EI) for C₂₄H₂₈N₆O₃: 449 (MH⁺).

- 5 [0598] N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-cyclopropanecarboxamide: NMR (400 MHz, d₆-DMSO): 10.45 (s, 1H), 9.43 (s, 1H), 8.42 (d, 1H), 8.17 (d, 2H), 7.75 (d, 2H), 7.64 (s, 1H), 7.15 (m, 2H), 6.84 (d, 1H), 3.80 (s, 3H), 3.75 (m, 4H), 2.96 (m, 4H), 2.52 (m, 2H), 0.80 (m, 2H). MS (EI) for C₂₅H₂₇N₅O₃: 446 (MH⁺).
- 10 [0599] N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-butanamide: NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 9.43 (s, 1H), 8.44 (d, 1H), 8.17 (m, 2H), 7.75 (d, 2H), 7.64 (s, 1H), 7.25 (m, 2H), 6.84 (d, 1H), 3.80 (s, 3H), 3.75 (m, 4H), 2.96 (m, 4H), 2.35 (q, 2H), 1.62 (m, 2H), 0.92 (q, 3H). MS (EI) for C₂₅H₂₉N₅O₃: 448 (MH⁺).
- 15 [0600] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)butanamide: NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 9.40 (s, 1H), 8.41 (d, 1H), 8.17 (d, 2H), 7.78 (d, 2H), 7.68 (d, 2H), 7.24 (s, 1H), 6.94 (d, 2H), 3.60 (m, 6H), 3.21 (s, 3H), 3.0 – 3.09 (m, 4H), 2.60 (q, 2H), 2.35 (m, 2H), 1.60 (m, 2H), 0.95 (q, 3H). MS (EI) for C₂₈H₃₄N₆O₃: 503 (MH⁺).
- 20 [0601] O-methyl-N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-L-serinamide: NMR (400 MHz, d₆-DMSO): 11.60 (s, 1H), 10.1 (s, 1H), 8.60 (s, 1H), 8.55 (m, 2H), 8.20 (m, 2H), 7.98 (s, 1H), 7.90 (d, 2H), 7.80 (s, 1H), 7.48 (m, 2H), 4.35 (m, 1H), 4.04 (m, 5H), 3.98 (s, 3H), 3.85 (m, 4H), 3.60 (m, 4H). MS (EI) for C₂₅H₃₀N₆O₄: 479 (MH⁺).
- 25 [0602] N-[4-(2-([3-(methyloxy)-4-morpholin-4-ylphenyl]amino)pyrimidin-4-yl)phenyl]-D-prolinamide: NMR (400 MHz, d₆-DMSO): 11.57 (s, 1H), 10.25 (br, 1H), 10.06 (s, 1H), 8.76 (br, 1H), 8.60 (d, 1H), 8.22 (d, 2H), 8.05 (s, 1H), 7.87 (m, 3H), 7.50 (m, 2H), 4.18 – 4.52 (m, 5H), 4.08 (m, 2H), 3.99 (s, 3H), 3.62 (m, 4H), 3.30 (m, 2H), 1.95 (m, 2H). MS (EI) for C₂₆H₃₀N₆O₃: 475 (MH⁺).
- 30 [0603] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)cyclopropanecarboxamide: NMR (400 MHz, d₆-DMSO): 10.45 (s, 1H), 9.40 (s, 1H), 8.41 (s, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.98 (d, 2H), 3.60 (m, 6H), 3.22 (s, 3H), 3.0 – 3.11 (m, 4H), 6.62 (q, 2H), 0.82 (m, 4H). MS (EI) for C₂₈H₃₂N₆O₃: 501 (MH⁺).
- 35 [0604] N-{4-[2-({4-[4-(Piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 11.40 (s, 1H), 10.0 (m, 1H), 9.96 (s, 1H), 9.11 (br d, 1H), 8.7-8.8 (m, 2H), 8.55 (d, 1H), 8.20 (d, 2H), 7.87

5 (m, 4H), 7.59 (br s, 2H), 7.45 (d, 1H), 4.48 (m, 1H), 3.4-3.5 (m, 4H), 3.25-3.30 (m, 4H) 3.0-3.1 (m, 1H), 2.9-3.0 (m, 2H), 2.4-2.5 (m, 1H), 1.9-2.0 (m, 3H), 1.7-1.9 (m, 4H); MS (EI) for $C_{31}H_{38}N_8O_2$: 555 (MH^+).

[0605] 3-(Methyloxy)-N-{4-[2-({4-[4-(piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}propanamide: 1H NMR (400 MHz, d_6 -DMSO):
10 10.49 (s, 1H), 9.93 (s, 1H), 9.07 (m, 1H), 8.72 (m, 1H), 8.50 (d, 1H), 8.13 (d, 2H), 7.85 (d, 2H), 7.79 (d, 2H), 7.56 (br s, 2H), 7.42 (d, 1H), 3.61 (t, 2H), 3.3-3.5 (m, 4H), 3.2-3.30 (m, 5H) 3.0-3.1 (m, 1H), 2.85-3.0 (m, 2H), 2.59 (t, 2H), 1.7-1.9 (m, 4H); MS (EI) for $C_{30}H_{37}N_7O_3$: 544 (MH^+).

[0606] 1-Ethyl-3-{4-[2-({4-[4-(piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}urea: 1H NMR (400 MHz, d_6 -DMSO): 9.96 (s, 1H), 9.20 (s, 1H),
15 8.93 (m, 1H), 8.65 (m, 1H), 8.46 (d, 1H), 8.09 (d, 2H), 7.79 (m, 2H), 7.58 (d, 2H), 7.41 (m, 3H), 6.51 (br s, 1H), 3.2-3.4 (m, 6H), 3.13 (q, 2H) 3.0-3.1 (m, 1H), 2.9-3.0 (m, 2H), 1.7-1.9 (m, 4H), 1.06 (t, 3H); MS (EI) for $C_{29}H_{36}N_8O_2$: 529 (MH^+).

[0607] N-(4-{2-[(4-{4-[3-(Dimethylamino)-2,2-dimethylpropanoyl]piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: 1H NMR (400 MHz, d_6 -DMSO):
20 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 6.57 (br s, 1H), 3.70 (m, 4H), 3.07 (m, 4H) 2.60 (br s, 2H), 2.28 (br s, 6H), 2.09 (s, 3H), 1.23 (s, 6H); MS (EI) for $C_{29}H_{37}N_7O_2$: 516 (MH^+).

[0608] 2-(Methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-ethanesulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 9.39 (s, 1H), 9.40 (s, 1H),
25 8.44 (d, 1H), 8.11 (d, 2H), 7.67 (d, 2H), 7.31 (d, 2H), 7.26 (d, 1H), 6.94 (d, 2H), 3.74 (m, 4H), 3.67 (t, 2H) 3.43 (t, 2H), 3.18 (s, 3H), 3.05 (m, 4H); MS (EI) for $C_{23}H_{27}N_5O_4S$: 470 (MH^+).

[0609] 3-(Methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propane-1-sulfonamide: MS (EI) for $C_{24}H_{29}N_5O_4S$: 484 (MH^+).

[0610] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(tetrahydrofuran-2-ylmethyl)benzamide: 1H NMR (400 MHz, d_6 -DMSO): 10.23 (s, 1H), 9.77 (s, 1H), 8.52 (d, 1H), 8.45 (d, 1H), 8.43 (d, 1H), 8.19 (d, 1H), 8.17 (d, 1H), 7.88-7.86 (m, 1H), 7.76 (s, 1H),
35 7.74 (s, 1H), 7.43-7.38 (m, 3H), 4.01-3.98 (m, 1H), 3.81-3.76 (m, 2H), 3.65-3.62 (m, 2H), 2.09 (s, 3H), 1.85-1.80 (m, 3H), 1.64-1.61 (m, 1H). MS (EI): 432.5 (MH^+).

[0611] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[3-(2-oxopyrrolidin-1-yl)propyl]benzamide: 1H NMR (400 MHz, d_6 -DMSO): 10.23 (s, 1H), 9.78 (s, 1H), 8.52 (d,

- 5 1H), 8.49 (s, 1H), 8.41 (t, 1H), 8.19 (dd, 2H), 7.87-7.84 (m, 1H), 7.76 (s, 1H), 7.74 (s, 1H), 7.40-7.38 (m, 3H), 3.27-3.23 (m, 6H), 2.22 (t, 2H), 2.09 (s, 3H), 1.96-1.88 (m, 2H), 1.73-1.70 (m, 2H). MS (EI): 473.5 (MH⁺).
- [0612] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(3s,5s,7s)-tricyclo-[3.3.1.1~3,7~]dec-1-yl]benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.73 (s, 1H), 8.52 (d, 1H), 8.35 (s, 1H), 8.16 (d, 2H), 7.84-7.81 (m, 1H), 7.76 (d, 2H), 7.55 (s, 1H), 7.39-7.31 (m, 3H), 2.09 (s, 3H), 1.67 (m, 15H). MS (EI): 482.6 (MH⁺).
- [0613] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(methyloxy)ethyl]-benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.23 (s, 1H), 9.77 (s, 1H), 8.52 (d, 1H), 8.46 (d, 2H), 8.19 (d, 1H), 8.18 (d, 1H), 7.88-7.85 (m, 1H), 7.77 (s, 1H), 7.75 (s, 1H), 7.43-7.39 (m, 3H), 3.48-3.41 (m, 4H), 3.29-3.27 (m, 3H), 2.09 (s, 3H). MS (EI): 406.3 (MH⁺).
- [0614] N-[4-(2-{{3-(1,3-thiazolidin-3-ylcarbonyl)phenyl}amino}pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.23 (s, 1H), 9.84 (s, 1H), 8.52 (d, 1H), 8.19-8.12 (m, 3H), 7.90-7.87 (m, 1H), 7.77-7.75 (m, 2H), 7.43-7.38 (m, 2H), 7.11 (d, 1H), 4.64 (m, 2H), 3.77 (m, 2H), 3.06 (m, 2H), 2.09 (s, 3H). MS (EI): 420.6 (MH⁺).
- 20 [0615] N-{4-[2-({3-[(4-pyridin-2-yl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.83 (s, 1H), 8.53 (d, 1H), 8.14-8.11 (m, 3H), 8.04 (t, 1H), 7.89-7.86 (m, 1H), 7.73 (d, 2H), 7.57-7.53 (m, 1H), 7.43-7.39 (m, 2H), 7.03-7.01 (m, 1H), 6.83 (d, 1H), 6.69-6.66 (m, 1H), 3.74 (m, 4H), 3.49 (m, 4H), 2.08 (s, 3H). MS (EI): 494.5 (MH⁺).
- 25 [0616] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{2-(methyloxy)phenyl}-methyl}benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 8.80 (d, 1H), 8.53-8.49 (m, 2H), 8.19 (dd, 2H), 7.94-7.91 (m, 1H), 7.74 (d, 2H), 7.52-7.49 (m, 1H), 7.44-7.39 (m, 1H), 7.26-7.19 (m, 2H), 6.99 (dd, 1H), 6.93-6.89 (m, 1H), 4.46 (d, 2H), 3.84 (s, 3H), 2.09 (s, 3H). MS (EI): 468.5 (MH⁺).
- 30 [0617] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{3-(methyloxy)phenyl}-methyl}benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 8.97 (t, 1H), 8.53-8.49 (m, 2H), 8.20-8.18 (m, 2H), 7.93-7.90 (m, 1H), 7.75 (d, 2H), 7.48-7.46 (m, 1H), 7.43-7.39 (m, 1H), 7.27-7.23 (m, 1H), 6.92-6.90 (m, 2H), 6.83-6.80 (m, 1H), 4.47 (d, 2H), 3.71 (s, 3H), 2.09 (s, 3H). MS (EI): 468.4 (MH⁺).
- 35 [0618] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-fluorophenyl)methyl]-benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 8.97 (t, 1H), 8.53-8.49 (m, 2H), 8.19-8.17 (m, 2H), 7.93-7.90 (m, 1H), 7.75 (d, 2H),

- 5 7.50-7.47 (m, 1H), 7.43-7.37 (m, 2H), 7.32-7.29 (m, 1H), 7.22-7.16 (m, 2H), 4.40 (d, 2H), 2.09 (s, 3H). MS (EI): 456.4 (MH⁺).
- [0619] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(4-fluorophenyl)methyl]-benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 9.00 (t, 1H), 8.53-8.50 (m, 2H), 8.19-8.17 (m, 2H), 7.90-7.88 (m, 1H), 7.75 (d, 2H), 10 7.47-7.36 (m, 4H), 7.19-7.13 (m, 2H), 4.47 (d, 2H), 2.09 (s, 3H). MS (EI): 456.5 (MH⁺).
- [0620] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(3,3-dimethylbutyl)-benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.76 (s, 1H), 8.53-8.51 (d, 1H), 8.41 (s, 1H), 8.33 (t, 1H), 8.18-8.17 (m, 2H), 7.88-7.85 (m, 1H), 7.75 (d, 2H), 7.39-7.37 (m, 2H), 3.30-3.26 (m, 2H), 2.08 (s, 3H), 1.48-1.44 (m, 2H), 0.94 (s, 9H). MS (EI): 432.4 (MH⁺).
- 15 [0621] N-[4-(2-{{3-(thiomorpholin-4-ylcarbonyl)phenyl}amino}pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.82 (s, 1H), 8.53 (d, 1H), 8.15-8.20 (m, 2H), 7.99 (t, 1H), 7.85-7.82 (m, 1H), 7.76 (d, 2H), 7.41-7.37 (m, 2H), 6.98-6.96 (m, 1H), 3.88 (m, 4H), 3.60 (m, 4H), 2.09 (s, 3H). MS (EI): 434.5 (MH⁺).
- [0622] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-thienylmethyl)benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 20 9.09 (t, 1H), 8.53-8.50 (m, 2H), 8.18 (d, 2H), 7.89-7.86 (m, 1H), 7.75 (d, 2H), 7.44-7.38 (m, 3H), 7.03 (m, 1H), 6.98-6.96 (m, 1H), 4.64 (d, 2H), 2.09 (s, 3H). MS (EI): 444.4 (MH⁺).
- [0623] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[3-(dimethylamino)propyl]benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 8.74 (m, 25 1H), 8.55-8.50 (m, 2H), 8.21-8.18 (m, 2H), 7.95-7.87 (m, 1H), 7.75 (d, 2H), 7.45-7.40 (m, 2H), 3.05 (s, 2H), 2.75 (s, 6H), 2.55-2.54 (m, 2H), 2.09 (s, 3H), 1.25-1.21 (m, 2H). MS (EI): 433.4 (MH⁺).
- [0624] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(2-chlorophenyl)ethyl]benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.77 (s, 1H), 8.54-8.51 30 (m, 2H), 8.44 (m, 1H), 8.18 (d, 2H), 7.88-7.86 (m, 1H), 7.75 (d, 2H), 7.45-7.36 (m, 4H), 7.28-7.25 (m, 2H), 3.55-3.50 (m, 2H), 3.01-2.98 (m, 2H), 2.08 (s, 3H). MS (EI): 486.8 (MH⁺).
- [0625] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{2-(trifluoromethyl)phenyl}methyl}benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.81 (s, 1H), 35 9.07 (t, 1H), 8.53-8.49 (m, 2H), 8.19-8.17 (m, 2H), 7.96-7.94 (m, 1H), 7.75-7.72 (m, 2H), 7.55-7.39 (m, 6H), 4.68 (d, 2H), 2.08 (s, 3H). MS (EI): 506.5 (MH⁺).

- 5 **[0626]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{3-(trifluoromethyl)-phenyl}methyl}benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.80 (s, 1H), 9.10 (t, 1H), 8.53-8.49 (m, 2H), 8.19-8.17 (m, 2H), 7.94-7.91 (m, 1H), 7.74 (d, 2H), 7.68-7.58 (m, 3H), 7.48-7.39 (m, 3H), 4.58 (d, 2H), 2.09 (s, 3H). MS (EI): 506.4 (MH⁺).
- [0627]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-{{4-(trifluoromethyl)-phenyl}methyl}benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.80 (s, 1H), 9.10 (t, 1H), 8.52 (m, 2H), 8.18 (d, 2H), 7.91-7.89 (m, 1H), 7.75-7.69 (m, 4H), 7.55 (d, 2H), 7.49-7.38 (m, 3H), 4.58 (d, 2H), 2.08 (s, 3H). MS (EI): 506.5 (MH⁺).
- 10 **[0628]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2,4-difluorophenyl)-methyl]benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.79 (s, 1H), 8.98 (t, 1H), 8.52 (m, 2H), 8.18 (d, 2H), 7.91-7.89 (m, 1H), 7.54 (d, 2H), 7.47-7.38 (m, 4H), 7.27-7.22 (m, 1H), 7.09-7.04 (m, 1H), 4.48 (d, 2H), 2.09 (s, 3H). MS (EI): 474.4 (MH⁺).
- 15 **[0629]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-ethyl-N-methylbenzamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.80 (s, 1H), 8.53 (d, 1H), 8.13 (d, 2H), 7.76-7.74 (m, 2H), 7.40-7.32 (m, 3H), 6.94 (m, 2H), 3.28-3.24 (m, 2H), 2.94 (m, 3H), 2.09 (s, 3H), 1.15-1.08 (m, 3H). MS (EI): 390.4 (MH⁺).
- 20 **[0630]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-({4-[(trifluoromethyl)-oxy]phenyl}methyl)benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.79 (s, 1H), 9.04 (t, 1H), 8.52 (m, 2H), 8.18 (d, 2H), 7.91-7.88 (m, 1H), 7.75 (d, 2H), 7.48-7.39 (m, 4H), 7.34-7.32 (m, 2H), 4.51 (d, 2H), 2.09 (s, 3H). MS (EI): 522.5 (MH⁺).
- 25 **[0631]** N-{4-[2-({3-[(4-acetylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.82 (s, 1H), 8.53 (t, 1H), 8.19 (d, 2H), 8.14-8.12 (m, 2H), 7.84-7.84 (m, 1H), 7.75 (d, 2H), 7.40-7.38 (m, 2H), 3.53-3.44 (m, 8H), 2.09 (s, 6H). MS (EI): 459.5 (MH⁺).
- [0632]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(cyclopropylmethyl)-benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.77 (s, 1H), 8.53-8.50 (m, 2H), 8.48 (d, 1H), 8.19(dd, 2H), 7.86-7.84 (m, 1H), 7.75 (d, 2H), 7.41-7.38 (m, 2H), 3.17-3.14 (m, 2H), 2.09 (s, 3H), 1.06 (m, 1H), 0.45-0.42 (m, 2H), 0.25-0.23 (m, 2H). MS (EI): 402.5 (MH⁺).
- 30 **[0633]** 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(2-fluorophenyl)-ethyl]benzamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.77 (s, 1H), 8.54-8.51 (m, 2H), 8.44 (m, 1H), 8.20-8.17 (m, 2H), 7.88-7.86 (m, 1H), 7.75 (d, 2H), 7.40-7.26 (m,
- 35

5 4H), 7.18-7.13 (m, 2H), 3.52-3.49 (m, 2H), 2.92-3.88 (m, 2H), 2.09 (s, 3H). MS (EI): 470.4 (MH⁺).

[0634] N-[4-(2-([3-(pyrrolidin-1-ylcarbonyl)phenyl]amino)pyrimidin-4-yl)phenyl]-
acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 8.53 (d, 1H), 8.20-
8.12 (m, 4H), 7.83 (d, 1H), 7.76-7.73 (m, 2H), 7.39-7.35 (m, 2H), 3.50-3.40 (m, 4H), 2.09 (s,
10 3H), 1.95-1.80 (m, 4H). MS (EI): 402.5 (MH⁺).

[0635] N-{4-[2-([3-[(4-pyrimidin-2-yl)piperazin-1-
yl]carbonyl]phenyl]amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d₆-
DMSO): 10.20 (s, 1H), 9.83 (s, 1H), 8.53 (d, 2H), 8.39 (d, 1H), 8.14-8.12 (m, 2H), 8.04 (t,
1H), 7.89-7.87 (m, 1H), 7.74 (d, 1H), 7.43-7.38 (m, 2H), 7.03-7.01 (m, 1H), 6.67 (t, 2H), 3.81
15 (m, 4H), 3.73 (m, 4H), 2.08 (s, 3H). MS (EI): 495.6 (MH⁺).

[0636] N-{4-[2-([4-[4-(9H-fluoren-2-ylmethyl)piperazin-1-
yl]phenyl]amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO):
10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.87 (t, 2H), 7.73 (d, 2H), 7.65 (d, 2H),
7.59-7.56 (m, 2H), 7.37-7.25 (m, 4H), 6.92 (d, 2H), 3.92 (s, 2H), 3.60 (s, 2H), 3.10 (m, 4H),
20 2.56 (m, 4H), 2.08 (s, 3H). MS (EI): 567.7 (MH⁺).

[0637] N-(4-{2-[(4-{4-[(3-methyl-2-thienyl)methyl]piperazin-1-
yl}phenyl]amino)pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO):
10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.65 (d, 2H), 7.33 (d,
1H), 7.26 (d, 1H), 6.92 (d, 2H), 6.85 (d, 1H), 3.63 (s, 2H), 3.07 (m, 4H), 2.56 (m, 4H), 2.17
25 (s, 3H), 2.08 (s, 3H). MS (EI): 499.5 (MH⁺).

[0638] N-(4-{2-[(4-{4-[(5-ethylfuran-2-yl)methyl]piperazin-1-yl}phenyl]amino]-
pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.35 (s,
1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 6.18 (d,
1H), 6.01 (d, 1H), 3.48 (s, 2H), 3.07 (m, 4H), 2.62-2.56 (m, 6H), 2.09 (s, 3H), 1.16 (t, 3H).
30 MS (EI): 497.6 (MH⁺).

[0639] N-(4-{2-[(4-{4-[(3-([4-(1,1-dimethylethyl)phenyl]oxy)phenyl)methyl]piperazin-
1-yl}phenyl]amino)pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO):
10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.64 (d, 2H), 7.42-7.38
(m, 2H), 7.33 (t, 1H), 7.26 (d, 1H), 7.08 (d, 1H), 6.97-6.86 (m, 6H), 3.52 (s, 2H), 3.06 (m,
35 4H), 2.52 (m, 4H), 2.09 (s, 3H), 1.27 (s, 9H). MS (EI): 627.7 (MH⁺).

[0640] N-{4-[2-([4-[4-(3-thienylmethyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]-
phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d,

5 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.64 (d, 2H), 7.51-7.45 (m, 1H), 7.35 (d, 1H), 7.29-7.25 (m, 1H), 7.08-7.04 (m, 1H), 6.91 (d, 2H), 3.53 (s, 2H), 3.07 (m, 4H), 2.52 (m, 4H), 2.09 (s, 3H). MS (EI): 485.6 (MH+).

[0641] methyl 4-({4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-phenyl]piperazin-1-yl}methyl)benzoate: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H),
10 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.94 (d, 2H), 7.73 (d, 2H), 7.65 (d, 2H), 7.50 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.85 (s, 3H), 3.61 (s, 2H), 3.09 (m, 4H), 2.54 (m, 4H), 2.09 (s, 3H). MS (EI): 537.7 (MH+).

[0642] N-(4-{2-[(4-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.43
15 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.34 (m, 4H), 3.07 (m, 4H), 2.39 (m, 4H), 2.09 (s, 3H), 2.03 (m, 3H), 1.75-1.68 (m, 2H). MS (EI): 477.5 (MH+).

[0643] N-(4-{2-[(4-{4-[(4-{3-(dimethylamino)propyl}oxy)phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO):
20 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.64 (d, 2H), 7.26-7.21 (m, 3H), 6.92-6.87 (m, 4H), 3.97 (t, 2H), 3.44 (m, 6H), 3.06 (m, 4H), 2.37 (t, 2H), 2.16 (s, 6H), 2.09 (s, 3H), 1.87-1.82 (m, 2H). MS (EI): 580.7 (MH+).

[0644] N-[4-(2-{[4-(4-{2-[(phenylmethyl)oxy]ethyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO):
25 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.38-7.26 (m, 6H), 6.92 (d, 2H), 4.50 (s, 2H), 3.59 (m, 3H), 3.07 (m, 3H), 2.58 (m, 6H), 2.09 (s, 3H). MS (EI): 523.5 (MH+).

[0645] N-(4-{2-[(4-{4-[(2-chloroquinolin-3-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.37 (s,
30 1H), 8.49 (s, 1H), 8.44 (d, 1H), 8.13-8.09 (m, 3H), 7.96 (d, 1H), 7.83-7.79 (m, 1H), 7.74 (d, 1H), 7.69-7.65 (m, 3H), 7.26 (d, 2H), 6.95 (d, 2H), 3.78 (s, 2H), 3.15 (m, 4H), 2.69 (m, 4H), 2.09 (s, 3H). MS (EI): 565.1 (MH+).

[0646] N-{4-[2-({4-[4-(2,2'-bithien-5-ylmethyl)piperazin-1-yl]phenyl)amino]-pyrimidin-4-yl}phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.36 (s,
35 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.49 (dd, 1H), 7.27-7.25 (m, 2H), 7.15 (d, 1H), 7.09-7.07 (m, 1H), 6.96-6.92 (m, 3H), 3.73 (s, 2H), 3.10 (m, 4H), 2.59 (m, 4H), 2.09 (s, 3H). MS (EI): 567.6 (MH+).

- 5 [0647] N-[4-(2-([4-(4-([4-(2-thienyl)phenyl)methyl]piperazin-1-yl)phenyl)amino)-pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.66-7.62 (m, 4H), 7.54-7.50 (m, 2H), 7.38 (d, 2H), 7.26 (d, 1H), 7.15-7.13 (m, 1H), 6.92 (d, 2H), 3.54 (s, 2H), 3.09 (m, 4H), 2.55-2.52 (m, 4H), 2.09 (s, 3H). MS (EI): 561.6 (MH⁺).
- 10 [0648] N-(4-{2-[(4-{4-[(4-cyanophenyl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.82 (d, 2H), 7.73 (d, 2H), 7.65 (d, 2H), 7.56 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.63 (s, 2H), 3.10-3.08 (m, 4H), 2.54-2.52 (m, 4H), 2.09 (s, 3H). MS (EI): 504.5 (MH⁺).
- 15 [0649] N-[4-(2-([4-(4-{[2,5-bis(methyloxy)phenyl]methyl}piperazin-1-yl)phenyl)-amino)pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.95-6.90 (m, 4H), 6.81-6.78 (m, 1H), 3.74 (s, 3H), 3.70 (s, 3H), 3.50 (s, 2H), 3.09 (m, 4H), 2.55 (m, 4H), 2.09 (s, 3H). MS (EI): 539.7 (MH⁺).
- 20 [0650] N-{4-[2-([4-(4-(2,2-diphenylethyl)piperazin-1-yl)phenyl)amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.82 (d, 2H), 7.65-7.59 (m, 6H), 7.50-7.46 (m, 4H), 7.26 (d, 1H), 7.21 (d, 2H), 6.92 (d, 2H), 4.45 (t, 1H), 3.61 (t, 2H), 3.09 (m, 4H), 2.54 (m, 4H), 2.09 (s, 3H). MS (EI): 569.6 (MH⁺).
- 25 [0651] N-{4-[2-([4-(4-(1H-pyrrol-2-ylmethyl)piperazin-1-yl)phenyl)amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.71 (s, 1H), 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.91 (d, 2H), 6.65-6.63 (m, 1H), 5.94-5.90 (m, 2H), 3.44 (s, 2H), 3.06 (m, 4H), 2.48 (m, 4H), 2.09 (s, 3H). MS (EI): 468.6 (MH⁺).
- 30 [0652] N-[4-(2-([4-(4-propylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.06 (m, 4H), 2.27 (m, 4H), 2.08 (s, 3H), 1.80 (s, 2H), 1.48 (m, 2H), 0.88 (t, 3H). MS (EI): 431.6 (MH⁺).
- [0653] N-[4-(2-([4-(4-butylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.74 (d, 2H), 7.64 (d, 2H), 7.25 (d, 1H), 6.92 (d, 2H), 3.07 (m, 4H), 2.31

5 (m, 4H), 2.09 (s, 3H), 1.87 (m, 2H), 1.44 (m, 2H), 1.30 (m, 2H), 0.90 (t, 3H). MS (EI): 445.6 (MH⁺).

[0654] N-{4-[2-({4-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.73 (d, 2H), 7.65 (d, 2H), 7.25 (d, 1H), 6.92 (d, 2H), 3.08 (m, 4H), 2.58
10 (m, 4H), 2.22 (d, 2H), 2.09 (s, 3H), 1.86 (s, 1H), 0.49 (m, 2H), 0.09 (m, 2H). MS (EI): 443.6 (MH⁺).

[0655] N-[4-(2-({4-[4-(pentanoylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)]-acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.60 (m, 4H), 3.07 (m, 2H),
15 3.02 (m, 2H), 2.35 (t, 2H), 2.09 (s, 3H), 1.49 (m, 2H), 1.31 (m, 2H), 0.89 (t, 3H). MS (EI): 473.6 (MH⁺).

[0656] N-{4-[2-({4-[4-(pyridin-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.62 (d, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.95 (t, 1H), 7.74 (d, 2H), 7.69 (d, 2H), 7.61 (d, 1H), 7.27
20 (d, 1H), 6.97 (d, 2H), 3.82 (t, 2H), 3.57 (t, 2H), 3.18 (t, 2H), 3.06 (t, 2H), 2.09 (s, 3H). MS (EI): 494.6 (MH⁺).

[0657] N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.68 (m, 2H), 8.44 (d, 1H), 8.10 (d, 2H), 7.89 (d, 1H), 7.74 (d, 2H), 7.69 (d, 2H), 7.51 (m, 1H),
25 7.28 (d, 2H), 3.80 (m, 2H), 3.49 (m, 2H), 3.18 (m, 2H), 3.09 (m, 2H), 2.09 (s, 3H). MS (EI): 494.6 (MH⁺).

[0658] N-{4-[2-({4-[4-(pyridin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.69 (d, 2H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.44 (d, 2H), 7.28 (d, 1H), 6.96
30 (d, 2H), 3.79 (m, 2H), 3.41 (m, 2H), 3.18 (m, 2H), 3.07 (m, 2H), 2.09 (s, 3H). MS (EI): 494.6 (MH⁺).

[0659] N-{4-[2-({4-[4-(1H-pyrazol-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO):
10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.75-7.68 (m, 4H), 7.28 (d, 2H), 6.97
35 (d, 3H), 3.75 (m, 4H), 3.11 (m, 4H), 2.09 (s, 3H). MS (EI): 483.5 (MH⁺).

[0660] N-(4-{2-[(4-{4-[(1-acetylpiperidin-4-yl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO):

5 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.26 (d, 1H), 6.97 (d, 2H), 3.65 (m, 4H), 3.02 (m, 8H), 2.62 (m, 1H), 2.09 (s, 3H), 1.99 (s, 3H), 1.66 (m, 2H), 1.56 (m, 2H). MS (EI): 542.7 (MH⁺).

[0661] N-(4-(2-(4-(4-(2-cyclopropylacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.54 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.59 (m, 4H), 3.04 (m, 4H), 2.30 (d, 2H), 2.09 (s, 3H), 0.97 (m, 1H), 0.45 (m, 2H), 0.14 (m, 2H). MS (EI): 471.6 (MH⁺).

[0662] N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.95 (d, 2H), 3.58 (m, 6H), 3.23 (s, 3H), 3.08 (m, 2H), 3.02 (m, 2H), 2.62 (t, 2H), 2.09 (s, 3H). MS (EI): 475.6 (MH⁺).

[0663] N-{4-[2-({4-[4-(2-[(2-(methyloxy)ethyl]oxy)acetyl]piperazin-1-yl}phenyl)amino)-pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.23 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.68 (d, 1H), 7.27 (d, 1H), 6.96 (d, 2H), 4.20 (s, 2H), 3.58 (m, 6H), 3.47 (m, 2H), 3.25 (s, 3H), 3.06 (m, 4H), 2.09 (s, 3H). MS (EI): 505.6 (MH⁺).

[0664] N-(4-(2-(4-(4-(2-(pyridin-3-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.44 (m, 3H), 8.10 (d, 2H), 7.75 (d, 2H), 7.68 (d, 2H), 7.64 (m, 1H), 7.34 (m, 1H), 7.28 (d, 1H), 6.96 (d, 1H), 3.83 (s, 2H), 3.70 (m, 2H), 3.63 (m, 2H), 3.05 (m, 4H), 2.09 (s, 3H). MS (EI): 508.6 (MH⁺).

[0665] (2R,4S)-4-hydroxy-N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.48 (d, 1H), 8.44 (d, 1H), 8.39 (dd, 1H), 8.11 (d, 1H), 7.74 (d, 2H), 7.69 (m, 3H), 7.28 (m, 3H), 6.94 (d, 2H), 3.59 (m, 4H), 3.01 (m, 4H), 2.86 (t, 2H), 2.73 (t, 2H), 2.09 (s, 3H). MS (EI): 522.6 (MH⁺).

[0666] N-{4-[2-({3-[4-(1,3-thiazol-2-ylmethyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.19 (s, 1H), 9.45 (s, 1H), 8.47 (d, 1H), 8.11 (d, 2H), 7.73 (d, 2H), 7.66 (d, 1H), 7.60 (s, 1H), 7.31 (d, 1H), 7.21 (d, 1H), 7.12 (t, 1H), 6.55 (dd, 1H), 3.90 (s, 2H), 3.17 (m, 4H), 2.66 (m, 4H), 2.07 (s, 3H). MS (EI): 486.6 (MH⁺).

- 5 [0667] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-5-oxo-L-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.33 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.14 (d, 2H), 7.93 (s, 1H), 7.79 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 4.23 (dd, 1H), 3.75 (m, 4H), 3.06 (m, 4H), 2.35 (m, 1H), 2.21 (m, 2H), 2.02 (m, 1H). MS (EI): 459.5 (MH⁺).
- 10 [0668] **(3S)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrrolidine-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.75 (s, 1H), 9.84 (s, 1H), 9.34 (m, 1H), 9.14 (m, 1H), 8.51 (d, 1H), 8.16 (d, 2H), 7.82 (d, 3H), 7.41 (d, 2H), 3.93 (m, 4H), 3.82 (m, 6H), 3.36 (m, 2H), 3.24 (m, 1H), 2.33-2.24 (m, 1H), 2.11-2.04 (m, 2H). MS (EI): 445.5 (MH⁺).
- 15 [0669] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-threoninamide:** ¹H NMR (400 MHz, d6-DMSO): 11.53 (s, 1H), 10.11 (s, 1H), 8.51 (d, 1H), 8.31 (d, 2H), 8.15 (d, 2H), 7.86 (t, 3H), 7.73 (d, 2H), 7.44 (d, 2H), 4.01 (m, 6H), 3.48 (m, 4H), 1.18 (s, 3H). MS (EI): 449.5 (MH⁺).
- 20 [0670] **N-{4-[2-({4-[4-(N,N-diethyl-beta-alanyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.38 (s, 1H), 8.42 (d, 1H), 8.09 (d, 2H), 7.73 (d, 2H), 7.67 (d, 2H), 7.25 (d, 1H), 6.95 (d, 2H), 3.58 (m, 4H), 3.05 (m, 4H), 2.81 (t, 2H), 2.66-2.55 (m, 6H), 1.93 (s, 3H), 0.99 (t, 6H). MS (EI): 516.7 (MH⁺).
- 25 [0671] **N¹-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-glutamamide:** ¹H NMR (400 MHz, d6-DMSO): 9.36 (s, 1H), 8.42 (d, 2H), 8.11 (d, 3H), 7.80 (d, 2H), 7.66 (d, 2H), 7.26 (d, 2H), 6.92 (d, 3H), 3.72 (m, 4H), 3.32 (m, 1H), 3.02 (m, 4H), 1.93 (s, 2H), 1.82 (s, 2H). MS (EI): 476.6 (MH⁺).
- 30 [0672] **(S)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.19 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.94 (d, 2H), 3.04 (m, 4H), 2.90 (m, 2H), 2.65 (m, 2H), 2.43 (m, 3H), 1.98 (m, 6H), 1.03 (t, 3H). MS (EI): 473.6 (MH⁺).
- 35 [0673] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-norvalinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.50 (s, 1H), 10.15 (s, 1H), 8.56 (m, 3H), 8.24 (d, 2H), 7.94 (d, 2H), 7.76 (m, 3H), 7.51 (d, 2H), 4.08 (m, 4H), 3.67 (d, 1H), 1.97 (m, 4H), 1.43 (m, 2H), 1.19 (m, 2H), 0.94 (m, 3H). MS (EI): 447.6 (MH⁺).

- 5 **[0674] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-norleucinamide:** ¹H NMR (400 MHz, d₆-DMSO): 11.55 (s, 1H), 10.19 (s, 1H), 8.57 (d, 2H), 8.26 (d, 1H), 8.01 (m, 4H), 7.80 (m, 3H), 7.53 (d, 2H), 4.05 (m, 4H), 3.68 (d, 1H), 1.92 (m, 4H), 1.36 (m, 4H), 1.19 (m, 2H), 0.99 (d, 3H). MS (EI): 461.6 (MH⁺).
- [0675] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alloisoleucinamide:** ¹H NMR (400 MHz, d₆-DMSO): 11.32 (s, 1H), 10.01 (s, 1H), 8.56 (d, 2H), 8.44 (d, 3H), 8.21 (d, 2H), 7.89 (m, 3H), 7.46 (d, 2H), 4.02 (m, 4H), 3.56 (d, 1H), 1.99 (m, 4H), 1.63 (m, 1H), 1.17 (m, 2H), 1.00 (d, 3H), 0.91 (d, 3H). MS (EI): 461.6 (MH⁺).
- [0676] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-leucinamide:** ¹H NMR (400 MHz, d₆-DMSO): 11.32 (s, 1H), 9.97 (s, 1H), 8.56 (d, 2H), 8.45 (d, 3H), 8.21 (d, 2H), 7.89 (m, 3H), 7.46 (d, 2H), 4.02 (m, 4H), 3.57 (d, 1H), 1.99 (m, 4H), 1.91 (m, 1H), 1.71 (t, 2H), 1.17 (t, 3H), 0.95 (t, 3H). MS (EI): 461.6 (MH⁺).
- [0677] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-ethylphenyl)benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.85 (d, 2H), 8.59 (s, 1H), 8.53 (d, 1H), 8.18 (d, 2H), 7.93 (m, 1H), 7.73 (d, 2H), 7.57 (d, 1H), 7.46 (t, 1H), 7.40 (d, 1H), 7.30-7.34 (m, 2H), 7.24-7.27 (m, 2H), 2.62-2.67 (m, 2H), 2.08 (s, 3H), 1.13-1.7 (t, 3H). MS (EI) for C₂₇H₂₅N₅O₂: 452.58 (MH⁺).
- [0678] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(phenylmethyl)-benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.80 (s, 1H), 8.98-9.01 (t, 1H), 8.52 (d, 1H), 8.50 (s, 1H), 8.18 (d, 2H), 7.9 (dd, 1H), 7.76 (s, 1H), 7.74 (s, 1H), 7.47 (d, 1H), 7.39-7.43 (m, 2H), 7.43 (s, 2H), 7.34 (d, 2H), 7.23-7.25 (m, 1H), 4.50 (d, 2H), 2.09 (s, 3H). MS (EI) for C₂₆H₂₃N₅O₂: 438.48 (MH⁺).
- [0679] N-{4-[2-({3-[(4-cyclopentylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.82 (s, 1H), 8.53 (d, 1H), 8.12 (s, 2H), 8.05 (s, 1H), 7.81 (dd, 1H), 7.76 (d, 2H), 7.38 (t, 2H), 6.95 (d, 1H), 2.89 (s, 2H), 2.73 (s, 2H), 2.34-2.45 (m, 5H), 2.09 (s, 3H), 2.73-2.75 (m, 2H), 1.52-1.59 (m, 2H), 1.46-1.50 (m, 2H), 1.27-1.33 (m, 2H): MS (EI) for C₂₈H₃₂N₆O₂: 485.8 (MH⁺).
- [0680] N-{4-[2-({3-[(4-pyrazin-2-ylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.17 (s, 1H), 9.84 (s, 1H), 8.54 (d, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 8.06-8.10 (m, 2H), 7.86-7.88 (m, 2H), 7.73 (d, 2H), 7.42 (d, 1H), 7.39 (d, 1H), 7.02-7.04 (m, 1H), 3.69 (m, 4H), 3.57 (m, 4H), 2.08 (s, 3H). MS (EI) for C₂₇H₂₆N₈O₂: 495.7 (MH⁺).

- 5 **[0681] N-(4-{2-[(3-{[4-(3-chlorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.84 (s, 1H), 8.53 (d, 1H), 8.13 (dd, 2H), 8.064 (t, 1H), 7.85-7.87 (m, 1H), 7.74 (d, 2H), 7.39-7.43 (m, 2H), 7.22 (t, 1H), 7.01-7.03 (m, 1H), 6.96 (t, 1H), 6.90 (dd, 1H), 6.81 (dd, 1H), 3.76 (m, 4H), 3.52 (m, 4H), 2.08 (s, 3H). MS (EI) for C₂₉H₂₇ClN₆O₂: 528.1 (MH⁺).
- 10 **[0682] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(1-methyl-1H-benzimidazol-2-yl)methyl]benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.80 (s, 1H), 9.01 (t, 1H), 8.52 (t, 2H), 8.17-8.19 (m, 2H), 7.91-7.93 (m, 1H), 7.75 (d, 2H), 7.50-7.59 (m, 3H), 7.39-7.43 (m, 2H), 7.16-7.26 (m, 2H), 4.80 (d, 2H), 3.86 (s, 3H), 2.10 (s, 3H). MS (EI) for C₂₈H₂₅N₇O₂: 492.4 (MH⁺).
- 15 **[0683] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-propylbenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.77 (s, 1H), 8.52 (d, 1H), 8.46 (s, 1H), 8.40 (t, 1H), 8.18 (d, 2H), 7.83-7.86 (m, 1H), 7.75 (d, 2H), 7.38-7.40 (m, 3H), 3.21-3.26 (m, 2H), 2.09 (s, 3H), 1.52-1.58 (m, 2H), 0.89-0.93 (t, 3H). MS (EI) for C₂₂H₂₃N₅O₂: 390.7 (MH⁺).
- [0684] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-**
 20 **cyclopropylbenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.24 (s, 1H), 9.77 (s, 1H), 8.52 (d, 1H), 8.47 (s, 1H), 8.40 (d, 1H), 8.18 (d, 2H), 7.82-7.85 (m, 1H), 7.76 (d, 2H), 7.39 (d, 1H), 7.36-7.37 (m, 2H), 2.84-2.89 (m, 1H), 2.09 (s, 3H), 0.69-0.73 (m, 2H), 0.56-0.60 (m, 2H). MS (EI) for C₂₂H₂₁N₅O₂: 388.7 (MH⁺).
- [0685] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(3-fluorophenyl)-**
 25 **methyl]benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.80 (s, 1H), 9.04 (t, 1H), 8.53 (d, 1H), 8.50 (s, 1H), 8.17-8.20 (m, 2H), 7.90-7.93 (m, 1H), 7.74 (d, 2H), 7.46-7.50 (m, 1H), 7.36-7.43 (m, 3H), 7.08-7.19 (m, 3H), 4.51 (d, 2H), 2.09 (s, 3H). MS (EI) for C₂₆H₂₂FN₅O₂: 456.5 (MH⁺).
- [0686] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(naphthalen-1-**
 30 **ylmethyl)-benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.79 (s, 1H), 9.03 (t, 1H), 8.52 (d, 2H), 8.15-8.23 (m, 3H), 7.95-7.97 (m, 1H), 7.90-7.93 (m, 1H), 7.85-7.87 (dd, 1H), 7.75 (d, 2H), 7.55-7.60 (m, 2H), 7.48-7.50 (m, 3H), 7.38-7.42 (m, 2H), 4.97 (d, 2H), 2.08 (s, 3H). MS (EI) for C₃₀H₂₅N₆O₂: 488.6 (MH⁺).
- [0687] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-**
 35 **(dimethylamino)ethyl]-N-methylbenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.80 (s, 1H), 8.53 (d, 1H), 8.13 (d, 2H), 8.00 (d, 1H), 7.82 (s, 1H), 7.75 (d, 2H), 7.35-7.39 (m,

5 2H)6.93 (d, 1H), 3.39-3.419 (m, 2H), 2.94 (s, 3H), 2.21 (m, 2H), 2.09 (s, 6H), 1.95 (s, 3H). MS (EI) for $C_{24}H_{28}N_6O_2$: 433.7 (MH^+).

[0688] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-methylphenyl)-methyl]benzamide: 1H -NMR (400MHz, d_6 -DMSO):10.22 (s, 1H), 9.79 (s, 1H), 8.56 (t, 1H), 8.52 (d,1H), 8.50 (s, 1H), 8.17-8.19 (m, 2H), 7.90-7.90 (m, 1H), 7.75 (d, 2H), 7.45-7.50 (s, 10 1H), 7.39-7.43 (m, 2H), 7.25-7.27 (1H), 7.14-7.18 (m, 3H), 4.47 (d, 2H), 2.34 (s, 3H), 2.09(s, 3H). MS (EI) for $C_{27}H_{25}N_5O_2$: 452.6 (MH^+).

[0689] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(3-chlorophenyl)methyl]-benzamide: 1H -NMR (400MHz, d_6 -DMSO):10.25 (s, 1H), 9.80 (s, 1H), 9.05 (t, 1H), 8.53 (d, 1H), 8.50 (s, 1H), 8.17-8.20 (m, 2H), 7.91-7.94 (m, 1H), 7.60 (s, 15 1H), 7.40 (s, 1H), 7.46-7.48 (m, 1H), 7.43 (d, 1H), 7.36-7.40 (m, 3H), 7.30-7.32 (m, 2H), 4.95 (d, 2H), 2.09 (s, 3H). MS (EI) for $C_{26}H_{22}ClN_5O_2$: 472.8 (MH^+).

[0690] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-phenylethyl)-benzamide: 1H -NMR (400MHz, d_6 -DMSO):10.22 (s, 1H), 9.78 (s, 1H), 8.51-8.53 (m, 2H), 8.46 (s, 1H), 8.19 (d, 2H), 7.85-7.88 (m, 1H), 7.76 (d, 2H), 7.37-7.38 (m, 3H), 7.27-7.32 (m, 20 4H), 7.19-7.23 (m, 1H), 3.47-3.52 (m, 2H), 2.84-2.88 (m, 2H), 2.08 (s, 3H). MS (EI) for $C_{27}H_{25}N_5O_2$: 452.6 (MH^+).

[0691] N-{4-[2-({3-[(4-methylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: 1H -NMR (400MHz, d_6 -DMSO):10.24 (s, 1H), 9.82 (s, 1H), 8.53 (d, 1H), 8.12-8.15(m, 2H), 8.04(s, 1H), 7.80-7.82 (m, 1H), 7.76 (d, 2H), 7.36-7.40 (m, 2H), 6.94-6.96 25 (m, 1H), 2.94 (s, 2H), 2.78 (s, 2H), 2.37 (s, 2H), 2.27 (s, 2H), 2.18 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{24}H_{26}N_6O_2$: 431.6 (MH^+).

[0692] N-(4-{2-[(3-{[4-(2-fluorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):10.22 (s, 1H), 9.84 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 8.08 (m, 1H), 7.85 (d, 1H), 7.77 (d, 2H), 7.39-7.43 (m, 2H), 30 7.05-7.17 (m, 2H), 6.99-7.05 (m, 4H), 3.81(s, 2H), 3.55 (s, 2H), 3.09 (s, 2H), 2.98 (s, 2H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{27}FN_6O_2$: 511.7 (MH^+).

[0693] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(phenyloxy)ethyl]-benzamide: 1H -NMR (400MHz, d_6 -DMSO):10.22 (s, 1H), 9.79 (s, 1H), 8.65 (t, 1H), 8.2 (d, 2H), 8.50 (s br, 1H), 8.9 (d, 2H), 7.88-7.90 (m, 1H), 7.76 (d, 2H), 7.38-7.45 (m, 3H), 7.27- 35 7.30 (m, 2H), 6.91-6.98 (m, 3H), 4.11-4.14 (m, 2H), 3.63-3.78 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{27}H_{25}N_5O_3$: 468.4 (MH^+).

- 5 **[0694] methyl 1-([3-([4-(acetylamino)phenyl]pyrimidin-2-yl)amino]phenyl)carbonyl]-piperidine-4-carboxylate:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.82 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 7.97 (t, 1H), 7.86-7.88 (m, 1H), 7.75 (d, 2H), 7.36-7.70 (m, 2H), 6.94-6.96 (m, 1H), 3.61 (s, 3H), 3.30-3.36 (m, 2H), 2.77-3.29 (m, 2H), 2.65-2.70 (m, 1H), 2.09 (s, 3H), 1.99-2.19 (m, 2H), 1.79-1.88 (m, 2H).
- 10 MS (EI) for C₂₆H₂₇N₅O₄: 474.6 (MH⁺).
- [0695] N-[4-(2-([3-([4-(methyloxy)phenyl]piperazin-1-yl)carbonyl]phenyl)amino)-pyrimidin-4-yl]phenyl]acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.83 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 8.04 (m, 1H), 7.86-7.89 (m, 1H), 7.75 (d, 2H), 7.39-7.43 (m, 2H), 7.12 (t, 1H), 7.00-7.03 (m, 1H), 6.53 (dd, 1H), 6.47 (t, 1H), 6.40 (dd, 1H), 3.78-3.80 (m, 2H), 3.71 (s, 3H), 3.20-3.24 (m, 2H), 3.12-3.19 (m, 2H), 2.09 (s, 3H). MS (EI) for
- 15 C₃₀H₃₀N₆O₃: 523.5 (MH⁺).
- [0696] 3-([4-(acetylamino)phenyl]pyrimidin-2-yl)amino)-N-(2-[2-(methyloxy)phenyl]-ethyl)benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 8.52-8.53 (m, 1H), 8.45 (s, 2H), 8.19 (d, 2H), 7.88 (s, 1H), 7.75 (d, 2H), 7.38-7.39 (m, 3H), 7.17-7.23 (m, 2H), 6.97 (d, 1H), 6.87 (t, 1H), 3.78 (s, 3H), 3.45-3.47 (m, 2H), 2.84-2.87 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₈H₂₇N₅O₃: 482.7 (MH⁺).
- 20 **[0697] N-[4-(2-([3-(1,3-dihydro-2H-isoindol-2-ylcarbonyl)phenyl]amino)pyrimidin-4-yl)-phenyl]acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.82 (s, 1H), 8.53 (d, 1H), 8.16 (s, 1H), 8.12-8.14 (m, 2H), 7.94-7.97 (m, 1H), 7.74 (s, 1H), 7.72 (s, 1H), 7.39-7.45 (m, 3H), 7.27-7.32 (m, 1H), 7.26 (d, 2H), 7.19-7.21 (m, 1H), 7.89 (s, 2H), 4.82 (s, 2H), 2.09 (s, 3H). MS (EI) for C₂₇H₂₃N₅O₂: 450.7 (MH⁺).
- 25 **[0698] 3-([4-(acetylamino)phenyl]pyrimidin-2-yl)amino)-N-(biphenyl-4-ylmethyl)-benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.80 (s, 1H), 9.04 (s br, 1H), 8.53 (d, 2H), 8.19 (d, 2H), 7.89 (d, 1H), 7.76 (d, 2H), 7.64 (m, 4H), 7.35-7.50 (m, 8H), 4.55 (s, 2H), 2.08 (s, 3H). MS (EI) for C₃₂H₂₇N₅O₂: 514.8 (MH⁺).
- 30 **[0699] N-(4-(2-([3-([4-(phenylcarbonyl)piperazin-1-yl]carbonyl]phenyl)amino]pyrimidin-4-yl)phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.25 (s, 1H), 9.83 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 8.04 (s, 1H), 7.86 (d, 1H), 7.76 (d, 2H), 7.44 (m, 5H), 7.39 (d, 2H), 7.00 (d, 1H), 3.56 (m, 8H) 2.09 (s, 3H). MS (EI) for
- 35 C₃₀H₂₈N₆O₃: 521.6 (MH⁺).
- [0700] N-[4-(2-([3-([4-(methyloxy)phenyl]piperazin-1-yl)carbonyl]phenyl)amino)-pyrimidin-4-yl]phenyl]acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.83 (s,

5 1H), 8.53 (d, 1H), 8.13 (m, 2H), 8.05 (m, 1H), 7.86 (m, 1H), 7.75 (d, 2H), 7.37-7.43 (m, 2H), 7.01 (m, 1H), 6.90 (m, 2H), 6.82 (m, 2H), 3.77 (m, 2H), 3.68 (s, 3H), 3.53 (m, 2H), 3.08 (m, 2H), 2.97 (m, 2H), 2.09 (s, 3H). MS (EI) for C₃₀H₃₀N₆O₃: 523.7 (MH⁺).

[0701] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-methyl-N-{[2-(methyloxy)-phenyl]methyl}benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H),
10 9.80 (s, 1H), 8.37 (m, 1H), 8.14 (d, 2H), 7.76 (m, 3H), 7.39 (d, 2H), 7.25-7.32 (m, 2H), 7.14-7.18 (m, 1H), 7.04 (m, 1H), 6.95 (d, 2H), 4.57 (d, 2H), 3.76 (d, 3H), 2.88 (s, 3H), 2.09 (s, 3H). MS (EI) for C₂₈H₂₇N₅O₃: 482.7 (MH⁺).

[0702] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-fluorophenyl)methyl]-N-methylbenzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H),
15 9.08 (s, 1H), 8.52 (m, 1H), 8.14 (d, 2H), 8.01 (m, 1H), 7.89 (m, 1H), 7.75 (d, 2H), 7.38 (m, 4H), 7.21 (m, 2H), 7.01 (m, 1H), 4.67 (d, 2H), 2.91 (s, 3H), 2.09 (s, 3H). MS (EI) for C₂₇H₂₄FN₅O₂: 470.6 (MH⁺).

[0703] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(2-pyridin-2-ylethyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 8.51-8.54 (m, 3H),
20 8.46 (s, 1H), 8.18 (d, 2H), 7.86-7.88 (m, 1H), 7.76 (d, 2H), 7.69-7.73 (m, 1H), 7.37-7.40 (m, 3H), 7.31 (m, 1H), 7.21-7.24 (m, 1H), 3.61-3.66 (m, 2H), 3.02 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₆H₂₄N₆O₂: 453.6 (MH⁺).

[0704] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(pyridin-2-ylmethyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.81 (s, 1H), 9.06 (m, 1H), 8.52
25 (m, 2H), 8.19 (d, 2H), 7.91 (m, 1H), 7.73-7.79 (m, 3H), 7.51 (d, 1H), 7.40 (m, 2H), 7.30 (d, 1H), 7.25-7.27 (m, 1H), 4.59 (d, 2H), 2.09 (s, 3H). MS (EI) for C₂₅H₂₂N₆O₂: 439.8 (MH⁺).

[0705] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(pyridin-3-ylmethyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 8.52 (d, 1H), 8.45 (s
30 br, 1H), 8.25 (d, 1H), 7.18 (d, 3H), 7.82-7.85 (m, 1H), 7.76 (d, 3H), 7.35-7.41 (m, 4H), 4.25 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₅H₂₂N₆O₂: 438.6 (MH⁺).

[0706] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(pyridin-4-ylmethyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.80 (s, 1H), 9.08 (t, 1H), 8.50-8.54 (m, 4H), 8.18 (d, 2H), 7.92 (d, 1H), 7.74 (d, 2H), 7.49 (d, 1H), 7.42 (t, 1H), 7.40 (d, 1H), 7.31 (d, 2H), 4.51 (d, 2H), 2.09 (s, 3H). MS (EI) for C₂₅H₂₂N₆O₂: 438.5 (MH⁺).

35 [0707] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-methyl-N-(phenylmethyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.81 (s, 1H),

5 8.53 (s br, 1H), 8.14 (d, 2H), 7.46 (d, 2H), 7.23-7.46 (m, 8H), 7.21 (s br, 1H), 7.02 (s br, 1H), 4.56 (d, 2H), 2.95 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{27}H_{25}N_5O_2$: 452.7 (MH^+).

[0708] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-cyclopentylbenzamide:
 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.75 (s, 1H), 8.52 (d, 1H), 8.45 (s br, 1H),
 8.25 (d, 1H), 8.17 (d, 2H), 7.84 (d, 1H) 7.75 (d, 2H), 7.35-7.42 (m, 3H), 4.22-4.27 (m, 1H),
 10 2.09 (s, 3H), 1.88-1.93 (m, 2H), 1.70 (m, 2H), 1.49-1.59 (m, 4H). MS (EI) for $C_{24}H_{25}N_5O_2$:
 416.8 (MH^+).

[0709] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(2-chlorophenyl)methyl]-benzamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.80 (s,
 1H), 9.00 (t, 1H), 8.53 (d, 1H), 8.50 (s br, 1H), 8.18 (d, 2H), 7.93 (d, 1H), 7.63 (d, 2H), 7.52
 15 (d, 1H), 7.28-7.48 (m, 6H), 4.56 (d, 2H), 2.09 (s, 3H). MS (EI) for $C_{26}H_{22}ClN_5O_2$: 473.0
 (MH^+).

[0710] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(4-chlorophenyl)methyl]-benzamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.79 (s,
 1H), 9.02 (t, 1H), 8.52 (d, 1H), 8.50 (s br, 1H), 8.18 (d, 2H), 7.90 (d, 1H), 7.74 (d, 2H), 7.46
 20 (d, 1H), 7.41 (m, 2H), 7.38 (s, 2H), 7.36 (m, 2H), 4.48 (d, 2H), 2.09 (s, 3H). MS (EI) for
 $C_{26}H_{22}ClN_5O_2$: 473.1 (MH^+).

[0711] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(furan-2-ylmethyl)-benzamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 8.91 (t, 1H), 8.52
 (d, 1H), 8.48 (s br, 1H), 8.17 (d, 2H), 7.88 (d, 1H), 7.76 (d, 2H), 7.58 (m, 1H), 7.44 (d, 1H),
 25 7.37-7.41 (m, 2H), 6.41 (m, 1H), 6.28 (dd, 1H), 4.48 (d, 2H), 2.09 (s, 3H). MS (EI) for
 $C_{24}H_{21}N_5O_3$: 428.6 (MH^+).

[0712] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(4-(methyloxy)phenyl)-methyl]benzamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H),
 9.77 (s, 1H), 8.92 (t, 1H), 8.52 (d, 1H), 8.49 (s br, 1H), 8.19 (d, 2H), 7.88 (d, 1H), 7.76 (d,
 30 2H), 7.45 (d, 1H), 7.37-7.41 (m, 2H), 7.27 (d, 2H), 6.89 (d, 2H), 4.42 (d, 2H), 3.72 (s, 3H),
 2.08 (s, 3H). MS (EI) for $C_{27}H_{25}N_5O_3$: 468.4 (MH^+).

[0713] N-[4-(2-({3-({4-[2-(methyloxy)phenyl]piperazin-1-yl}carbonyl)phenyl}amino)-pyrimidin-4-yl)phenyl]acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.82 (s,
 1H), 8.53 (d, 1H), 8.14 (d, 2H), 8.06 (s, 1H) 7.86 (d, 1H), 7.76 (d, 2H), 7.40 (m, 2H), 7.01 (d,
 35 1H), 6.59 (m, 2H), 6.88 (s, 2H), 3.78 (s, 3H), 3.55 (m, 2H), 3.03 (m, 2H), 2.94 (s, 2H), 2.79
 (s, 2H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{30}N_6O_3$: 523.5 (MH^+).

- 5 **[0714] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[3-(methoxy)propyl]-benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.76 (s, 1H), 8.52 (d, 1H), 8.46 (s br, 1H), 8.40 (t, 1H), 8.18 (d, 2H), 7.85-7.88 (m, 1H), 7.75 (d, 2H), 7.38-7.40 (m, 3H), 3.40 (m, 2H), 3.30 (m, 2H), 3.24 (s, 3H), 2.09 (s, 3H), 1.74-1.80 (m, 2H). MS (EI) for C₂₃H₂₅N₅O₃: 420.5 (MH⁺).
- 10 **[0715] N-(4-{2-[(3-{[(2R,6S)-2,6-dimethylmorpholin-4-yl]carbonyl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.23 (s, 1H), 9.81 (s, 1H), 8.53 (d, 1H), 8.13 (d, 2H), 7.99 (m, 1H), 7.86 (d, 1H), 7.76 (d, 2H), 7.37-7.41 (m, 2H), 6.98 (d, 1H), 2.94 (s, 2H), 2.79 (s, 2H), 2.09 (s, 3H), 1.96 (s, 1H), 1.16 (m, 3H), 0.99 (m, 3H). MS (EI) for C₂₅H₂₇N₅O₃: 445.5 (MH⁺).
- 15 **[0716] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[(6-chloropyridin-3-yl)methyl]benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.79 (s, 1H), 9.06 (t, 1H), 8.52 (d, 1H), 8.49 (s, 1H), 8.40 (d, 1H), 8.18 (d, 2H), 7.91 (m, 2H), 7.81 (dd, 1H), 7.74 (d, 2H), 7.49 (d, 1H), 7.41-7.46 (m, 2H), 7.39 (d, 1H), 4.50 (d, 2H), 2.09 (s, 3H). MS (EI) for C₂₅H₂₁ClN₆O₂: 474.1 (MH⁺).
- 20 **[0717] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-butylbenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.24 (s, 1H), 9.80 (s, 1H), 8.53 (d, 1H), 8.13 (d, 2H), 8.02 (s br, 1H), 7.78 (m, 1H), 7.75 (d, 2H), 7.39 (d, 1H), 7.35 (d, 1H), 6.91 (d, 2H), 3.26 (m, 2H), 2.09 (s, 3H), 1.62 (m, 2H), 1.32 (m, 2H), 1.08 (m, 3H). MS (EI) for C₂₃H₂₅N₅O₂: 404.5 (MH⁺).
- 25 **[0718] N-(4-{2-[(3-{[4-(2-chlorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.83 (s, 1H), 8.53 (d, 1H), 8.14 (d, 2H), 7.90 (s br, 1H), 7.85 (d, 1H), 7.76 (d, 2H), 7.38-7.43 (m, 3H), 7.28 (m, 1H), 7.13 (dd, 1H), 7.07 (dd, 1H), 7.03 (m, 1H), 3.81 (m, 4H), 3.58 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₇ClN₆O₂: 528.1 (MH⁺).
- 30 **[0719] 3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-ethyl-N-[2-(methoxy)-ethyl]benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.80 (s, 1H), 8.53 (d, 1H), 8.13 (d, 2H), 7.99 (m, 2H), 7.82 (m, 1H), 7.76 (d, 2H), 7.39 (d, 2H), 7.36 (d, 2H), 6.92 (d, 1H), 3.57 (m, 2H), 3.12 (m, 2H), 2.94 (s, 3H), 2.09 (s, 3H), 1.10 (m, 3H). MS (EI) for C₂₄H₂₇N₅O₃: 434.4 (MH⁺).
- 35 **[0720] N-{4-[2-({4-[4-(phenylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H),

5 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.35 (d, 4H), 7.27 (d, 2H), 6.92 (d, 2H), 3.54 (s, 2H), 3.9 (m, 4H), 2.53 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{30}N_6O$: 479.7 (MH^+).

[0721] N-(4-{2-[(4-{4-[(5-methyl-3-phenylisoxazol-4-yl)methyl]piperazin-1-

yl}phenyl)-amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):

10.20 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.93-7.96 (m, 2H), 7.74 (d, 2H), 7.66
10 (d, 2H), 7.50-7.53 (m, 3H), 7.26 (d, 1H), 6.93 (d, 2H), 3.43 (s, 2H), 3.09 (m, 4H), 2.56 (m,
4H), 2.48 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{33}H_{33}N_7O_2$: 560.4 (MH^+).

[0722] N-(4-{2-[(4-{4-[(5-methyl-1-phenyl-1H-pyrazol-4-yl)methyl]piperazin-1-yl}-

phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.20

(s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.56 (s, 1H), 7.53
15 (d, 4H), 7.40-7.45 (m, 1H), 7.26 (d, 1H), 6.92 (d, 2H), 3.43 (s, 2H), 3.09 (m, 4H), 2.56 (m,
4H), 2.31 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{33}H_{34}N_8O$: 559.7 (MH^+).

[0723] N-(4-{2-[(4-{4-[(2-phenyl-1,3-thiazol-4-yl)methyl]piperazin-1-

yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):

10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.94 (d, 2H), 7.74 (d, 2H), 7.65 (d,
20 2H), 7.57 (s, 1H), 7.48-7.53 (m, 3H), 7.26 (d, 1H), 6.93 (d, 1H), 3.73 (s, 2H), 3.11 (m, 4H),
2.65 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{32}H_{31}N_7OS$: 562.5 (MH^+).

[0724] N-[4-(2-{[4-(4-{[6-(phenyloxy)pyridin-3-yl]methyl]piperazin-1-

yl}phenyl]amino)-pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):

10.23 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 8.07 (d, 1H), 7.81 (dd, 1H), 7.74 (d,
25 2H), 7.65 (d, 2H), 7.43 (t, 2H), 7.26 (d, 1H), 7.19-7.23 (m, 1H), 7.14 (d, 2H), 7.01 (d, 1H),
6.92 (d, 2H), 3.21 (s, 2H), 3.08 (m, 4H), 2.48 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{34}H_{33}N_7O_2$:
572.4 (MH^+).

[0725] N-{4-[2-({4-[4-(cyclohexylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-

yl]-phenyl}acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d,

30 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.06 (m, 4H), 2.47
(m, 4H), 2.13 (d, 2H), 2.09 (s, 3H), 1.76 (d, 2H), 1.65 (m, 3H), 1.49-1.54 (m, 1H), 1.12-1.17
(m, 3H), 0.80-0.89 (m, 2H). MS (EI) for $C_{29}H_{36}N_6O$: 485.8 (MH^+).

[0726] N-(4-{2-[(4-{4-[(1S,4S)-bicyclo[2.2.1]hept-5-en-2-ylmethyl]piperazin-1-

yl}phenyl)-amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):

35 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d,
1H), 6.92 (d, 2H), 6.14-6.16 (m, 1H), 5.95-5.97 (m, 1H), 3.07 (m, 4H), 2.79 (d, 2H), 2.45 (m,

5 4H), 2.32-2.39 (m, 2H), 2.09 (s, 3H), 1.95-1.99 (m, 1H), 1.81-1.87 (m, 1H), 1.31 (m, 1H), 1.23 (m, 1H), 0.51 (m, 1H). MS (EI) for $C_{30}H_{34}N_6O$: 495.7 (MH^+)

[0727] N-[4-(2-[[4-(4-pentylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.07 (m, 4H), 2.55 (m, 4H), 2.307 (t, 2H), 2.09 (s, 3H), 1.43-1.49 (m, 2H), 1.22-1.34 (m, 4H), 0.88 (t, 3H). MS (EI) for $C_{27}H_{34}N_6O$: 459.7 (MH^+).

[0728] N-(4-{2-[[4-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.66 (d, 2H), 7.54 (dd, 1H), 7.45 (dd, 1H), 7.29-7.39 (m, 2H), 7.26 (d, 1H), 6.93 (d, 2H), 3.64 (s, 2H), 3.10 (m, 4H), 2.60 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{29}ClN_6O$: 514.1 (MH^+).

[0729] N-[4-(2-[[4-(4-{[3,5-bis(methyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.93 (d, 2H), 6.51 (d, 2H), 6.39 (t, 1H), 3.75 (s, 6H), 3.46 (s, 2H), 3.09 (m, 4H), 2.61 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{31}H_{34}N_6O_3$: 539.8 (MH^+).

[0730] N-(4-{2-[[4-{4-[(4-fluorophenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.36-7.39 (m, 2H), 7.26 (d, 1H), 7.16 (t, 2H), 6.92 (d, 2H), 3.51 (s, 2H), 3.08 (m, 4H), 2.28 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{29}FN_6O$: 497.8 (MH^+).

[0731] N-(4-{2-[[4-{4-[(1-methyl-1H-pyrrol-2-yl)methyl]piperazin-1-yl}phenyl]amino]-pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 6.68 (t, 1H), 5.88-5.91 (m, 2H), 3.61 (s, 2H), 3.4 (s, 3H), 3.06 (m, 4H), 2.58 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{28}H_{31}N_7O$: 482.8 (MH^+).

[0732] N-[4-(2-[[4-(4-{[5-(3-chlorophenyl)furan-2-yl]methyl}piperazin-1-yl)phenyl]-amino}pyrimidin-4-yl)phenyl]acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.20 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.72-7.76 (m, 3H), 7.63-7.66 (m, 3H), 7.45 (t, 1H), 7.33 (d, 1H), 7.26 (d, 1H), 7.06 (d, 1H), 6.92 (d, 2H), 6.48 (d, 1H), 3.64 (s, 2H), 3.10 (m, 4H), 2.60 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{33}H_{31}ClN_6O$: 580.3 (MH^+).

- 5 [0733] N-[4-(2-{[4-(4-{[4-fluoro-2-(trifluoromethyl)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.37 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.84-7.88 (m, 1H), 7.74 (d, 2H), 7.66 (d, 2H), 7.54-7.72 (m, 2H), 7.26 (d, 1H), 6.93 (d, 2H), 3.66 (s, 2H), 3.10 (m, 4H), 2.56 (m, 4H), 2.09 (s, 3H). MS (EI) for C₃₀H₂₈F₄N₆O: 565.3 (MH⁺).
- 10 [0734] N-[4-(2-{[4-(4-{[4-(1H-imidazol-1-yl)phenyl]methyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.25 (t, 1H), 7.11 (d, 2H), 7.74 (d, 3H), 7.61-7.67 (m, 4H), 7.48 (d, 2H), 7.26 (d, 1H), 7.11 (t, 1H), 6.93 (d, 2H), 3.58 (s, 2H), 3.10 (m, 4H), 2.55 (m, 4H), 20.9 (s, 3H). MS (EI) for C₃₂H₃₂N₈O: 545.8 (MH⁺).
- 15 [0735] N-[4-(2-{[4-(4-{[2,5-bis(trifluoromethyl)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.37 (s, 1H), 8.43 (d, 1H), 8.17 (s, 1H), 8.10 (d, 2H), 8.00 (d, 1H), 7.89 (d, 1H), 7.74 (d, 2H), 7.67 (d, 2H), 7.26 (d, 1H), 6.94 (d, 2H), 3.79 (s, 2H), 3.12 (m, 4H), 2.60 (m, 4H), 2.09 (s, 3H). MS (EI) for C₃₁H₂₈F₆N₆O: 615.7 (MH⁺).
- 20 [0736] N-(4-{2-[(4-{4-[(2,6-dimethylphenyl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.24 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 7.00-7.08 (m, 3H), 6.90 (d, 2H), 3.50 (s, 2H), 3.02 (m, 4H), 2.54 (m, 4H), 2.37 (s, 6H), 2.09 (s, 3H). MS (EI) for C₃₁H₃₄N₆O: 507.7 (MH⁺).
- 25 [0737] N-(4-{2-[(4-{4-[(2,3-dimethylphenyl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 7.01-7.09 (m, 3H), 6.91 (d, 2H), 3.45 (s, 2H), 3.05 (m, 4H), 2.53 (m, 4H), 2.24 (d, 6H), 2.09 (s, 3H). MS (EI) for C₃₁H₃₄N₆O: 507.8 (MH⁺).
- 30 [0738] N-[4-(2-{[4-(4-{[2,4-bis(ethyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.35 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 7.19 (d, 1H), 6.91 (d, 2H), 6.47-6.51 (m, 2H), 3.97-4.04 (m, 4H), 3.46 (s, 2H), 3.06 (m, 4H), 2.52 (m, 4H), 2.09 (s, 3H), 1.30-1.35 (m, 6H). MS (EI) for C₃₃H₃₈N₆O₃: 567.8 (MH⁺).
- 35 [0739] N-[4-(2-{[4-(4-{[3-(ethyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.21 (s, 1H), 9.36 (s, 1H), 8.45 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.24 (t, 1H), 7.11-7.19 (m, 3H), 7.02-7.09 (m,

5 3H), 6.91 (d, 2H), 3.96-4.00 (m, 2H), 3.40 (s, 2H), 3.07 (m, 4H), 2.59 (m, 4H), 2.09 (s, 3H), 1.32-1.38 (m, 3H). MS (EI) for $C_{31}H_{34}N_6O_2$: 523.8 (MH^+).

[0740] N-{4-[2-({4-[4-(3-methylbutanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.23 (s, 1H), 9.04 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 6.96 (d, 2H) 3.67 (m, 4H), 3.06
10 (m, 2H), 3.02 (m, 2H), 2.24 (d, 2H), 2.09 (s, 3H), 1.97-2.09 (m, 1H), 0.92 (d, 6H). MS (EI) for $C_{27}H_{32}N_6O_2$: 473.8 (MH^+).

[0741] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 6.95 (d, 2H), 3.59 (m, 2H), 3.46
15 (m, 2H), 3.39 (t, 1H), 3.02 (m, 4H), 2.11-2.23 (m, 4H), 2.09 (s, 3H), 1.89-1.92 (m, 1H), 1.72-1.78 (m, 1H). MS (EI) for $C_{27}H_{30}N_6O_2$: 470.7 (MH^+).

[0742] N-{4-[2-({4-[4-(cyclopentylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 6.96 (d, 2H), 3.63 (m, 4H),
20 3.04 (m, 4H), 2.98 (m, 1H), 2.09 (s, 3H), 1.74-2.09 (m, 2H), 1.51-1.72 (m, 6H). MS (EI) for $C_{28}H_{32}N_6O_2$: 485.5 (MH^+).

[0743] N-[4-(2-({4-(4-([2-(methyloxy)phenyl]carbonyl)piperazin-1-yl)phenyl]amino)-pyrimidin-4-yl)phenyl]acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.42 (m, 1H), 7.27 (d, 1H), 7.22
25 (dd, 1H), 7.10 (d, 1H), 7.02 (m, 1H), 6.96 (d, 2H), 3.81 (s, 3H), 3.77 (m, 2H), 3.27 (m, 2H), 3.12 (m, 2H), 3.01 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{30}N_6O_3$: 523.7 (MH^+).

[0744] N-(4-{2-[(4-{4-[(2-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO):
10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.30-7.36
30 (m, 2H), 7.28 (d, 2H), 7.19-7.24 (m, 1H), 6.96 (d, 2H), 3.82 (m, 2H), 3.28 (m, 2H), 3.16 (m, 2H), 3.00 (m, 2H), 2.25 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{30}N_6O_2$: 507.8 (MH^+).

[0745] N-[4-(2-({3-(4-([2-(methyloxy)phenyl]methyl)piperazin-1-yl)phenyl]amino)-pyrimidin-4-yl)phenyl]acetamide: 1H NMR (400 MHz, d_6 -DMSO): 11.21 (s, 1H), 9.46 (s, 1H), 8.49 (d, 1H), 8.12 (d, 2H), 7.73 (d, 2H), 7.63 (s, 1H), 7.36 (d, 1H), 7.32 (d, 1H), 7.21
35 (m, 2H), 7.10 (t, 1H), 6.98 (m, 2H), 6.55 (d, 1H), 3.80 (s, 3H), 3.54 (s, 2H), 3.16 (m, 4H), 2.57 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{32}N_6O_2$: 509.6 (MH^+).

- 5 [0746] **N-{4-[2-({4-[(2R,6S)-2,6-dimethylmorpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.44 (s, 1H), 10.07 (s, 1H), 8.74 (s, 1H), 8.57 (d, 1H), 8.21 (s, 2H), 7.90 (m, 4H), 7.73 (s, 2H), 7.48 (d, 1H), 4.49 (m, 1H), 4.24 (m, 2H), 4.02 (m, 1H), 3.57 (m, 3H), 3.29 (m, 2H), 1.99 (m, 4H), 1.18 (m, 7H). MS (EI) for C₂₇H₃₂N₆O₂: 473.5 (MH⁺).
- 10 [0747] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-5-oxo-D-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.33 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.14 (d, 2H), 7.93 (s, 1H), 7.79 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 4.23 (m, 1H), 3.75 (m, 4H), 3.05 (m, 4H), 2.35 (m, 1H), 2.21 (m, 2H), 2.03 (m, 1H). MS (EI) for C₂₅H₂₆N₆O₃: 459.5 (MH⁺).
- 15 [0748] **N¹-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-aspartamide:** ¹H NMR (400 MHz, d6-DMSO): 11.71 (s, 1H), 10.09 (s, 1H), 8.94 (s, 3H), 8.58 (d, 1H), 8.23 (d, 2H), 7.91 (m, 4H), 7.73 (s br, 2H), 7.49 (d, 1H), 4.52 (m, 4H), 4.05 (m, 5H), 3.33 (d, 1H), 3.29 (d, 1H). MS (EI) for C₂₄H₂₇N₇O₃: 462.5 (MH⁺).
- [0749] **N¹-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-glutamamide:** ¹H NMR (400 MHz, d6-DMSO): 9.39 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.83 (d, 2H), 7.67 (d, 2H), 7.35 (s, 1H), 7.29 (d, 1H), 6.93 (d, 2H), 6.77 (s, 1H), 3.75 (m, 4H), 3.37 (t, 1H), 3.05 (m, 4H), 1.88 (m, 2H), 1.70 (m, 1H). MS (EI) for C₂₅H₂₉N₇O₃: 476.5 (MH⁺).
- 20 [0750] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-threoninamide:** ¹H NMR (400 MHz, d6-DMSO): 11.26 (s, 1H), 9.89 (s, 1H), 8.54 (d, 1H), 8.31 (s, 2H), 8.19 (d, 2H), 7.86 (d, 3H), 7.44 (m, 3H), 4.11 (m, 2H), 3.96 (m, 4H), 3.39 (m, 4H), 1.23 (d, 3H). MS (EI) for C₂₄H₂₈N₆O₃: 449.5 (MH⁺).
- 25 [0751] **N-(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.43 (s, 1H), 9.99 (s, 1H), 8.55 (d, 1H), 8.20 (d, 2H), 8.03 (s, 1H), 7.89 (2H), 7.71 (dd, 1H), 7.46 (d, 2H), 7.20 (d, 1H), 4.49 (m, 1H), 3.76 (m, 4H), 3.28 (m, 2H), 2.96 (m, 4H), 1.97 (m, 4H). MS (EI) for C₂₅H₂₇ClN₆O₂: 479.9 (MH⁺).
- 30 [0752] **N-(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.55 (s, 1H), 10.15 (s, 1H), 8.56 (d, 1H), 8.21 (d, 2H), 8.01 (s, 1H), 7.89 (d, 2H), 7.71 (dd, 1H), 7.49 (d, 2H), 7.23 (d, 1H), 4.51 (m, 1H), 3.76 (m, 4H), 3.29 (m, 2H), 2.97 (m, 4H), 1.97 (m, 4H). MS (EI) for C₂₅H₂₇ClN₆O₂: 479.9 (MH⁺).
- 35 [0753] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-leucinamide:** ¹H NMR (400 MHz, d6-DMSO): 11.41 (s, 1H), 10.05 (s, 1H), 8.56 (d, 1H),

5 8.60 (m, 3H), 8.25 (d, 2H), 7.96 (m, 3H), 7.69 (m, 2H), 7.47 (d, 1H), 4.14 (m, 1H), 4.04 (m, 4H), 3.57 (m, 4H), 1.71 (m, 2H), 1.19 (m, 1H), 1.00 (s, 6H). MS (EI) for C₂₆H₃₂N₆O₂: 461.5 (MH⁺).

[0754] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-
isoleucinamide: ¹H NMR (400 MHz, d₆-DMSO): 11.41 (s, 1H), 10.10 (s, 1H), 8.56 (d, 1H),
10 8.47 (m, 2H), 8.20 (d, 2H), 7.92 (m, 3H), 7.73 (m, 2H), 7.47 (d, 1H), 4.12 (m, 4H), 3.57 (m, 4H), 1.65 (m, 2H), 1.18 (m, 2H), 1.00 (d, 3H), 0.89 (t, 3H). MS (EI) for C₂₆H₃₂N₆O₂: 461.5 (MH⁺).

[0755] (2R)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-
yl}phenyl)-butanamide: ¹H NMR (400 MHz, d₆-DMSO): 11.57 (s, 1H), 10.20 (s, 1H), 8.56
15 (m, 3H), 8.24 (m, 2H), 7.97 (m, 4H), 7.82 (s, 1H), 7.54 (s, 1H), 4.05 (m, 5H), 1.96 (m, 4H), 1.10 (m, 5H). MS (EI) for C₂₄H₂₈N₆O₂: 433.5 (MH⁺).

[0756] N-(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-
carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.5 (s, 1H), 9.40 (br s, 1H), 8.55 (d, 1H),
8.23 (d, 2H), 8.09 (dd, 1H), 7.96-7.91 (m, 5H), 7.53 (m, 1H), 7.45 (d, 1H), 7.32-7.25 (m, 3H),
20 6.74 (br s, 1H). MS (EI): 388.0 (MH⁺).

[0757] N-(3-{[4-(4-aminophenyl)pyrimidin-2-yl]amino}phenyl)-2,6-
dichlorobenzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.7 (s, 1H), 9.67 (s, 1H), 8.37-8.35
(m, 2H), 8.01 (d, 2H), 7.61-7.58 (m, 2H), 7.51-7.49 (m, 1H), 7.44 (dt, 1H), 7.31-7.22 (m,
3H), 6.69 (d, 2H), 5.95 (br, 2H)); MS (EI): 450.0 (MH⁺).

25 [0758] 4-[[2-chloro-4-(methyloxy)phenyl]oxy]-N-(4-morpholin-4-ylphenyl)pyrimidin-
2-amine: ¹H-NMR (400MHz, d₆-DMSO): 9.37 (s, 1H), 8.31 (d, 1H), 7.33-7.23 (m, 4H), 7.02
(dd, 1H), 7.00 (br d, 2H), 6.41 (d, 1H), 3.83 (s, 3H), 3.73-3.71 (m, 4H), 2.98-2.96 (m, 4H).
MS (EI): 412.8 (MH⁺).

[0759] N-[4-(2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-
30 yl)oxy)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.2 (s, 1H), 9.92 (br s, 1H),
8.33 (d, 1H), 7.66 (d, 2H), 7.42 (br s, 2H), 7.17 (d, 2H), 7.10 (br s, 2H), 6.50 (d, 1H), 3.86 (br
s, 4H), 3.24 (br s, 4H), 2.08 (s, 3H); MS (EI): 406.1 (MH⁺).

[0760] N-[4-(2-{[4-(4-D-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-3-
(methyloxy)propanamide: ¹H-NMR (400MHz, d₆-DMSO): 10.3 (s, 1H), 9.61 (br s, 1H),
35 8.46 (d, 1H), 8.16-8.12 (m, 4H), 7.78 (d, 2H), 7.72 (d, 2H), 7.34 (d, 1H), 7.12 (br s, 2H),
4.47-4.44 (m, 1H), 3.80 (br s, 4H), 3.64 (t, 2H), 3.64 (s, 3H), 3.07 (br s, 4H), 2.60 (t, 2H),
1.34 (d, 3H); MS (EI): 504.2 (MH⁺).

- 5 [0761] 3-(methoxy)-N-[4-(2-{{4-(4-L-prolylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]propanamide: ¹H-NMR (400MHz, d6-DMSO): 10.3 (s, 1H), 9.51 (s, 1H), 8.54 (m, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.71 (d, 2H), 7.31 (d, 1H), 7.04 (br d, 2H), 4.68 (m, 1H), 4.07 (br s, 4H), 3.64 (t, 2H), 3.25 (s, 3H), 3.17 (br s, 4H), 2.60 (t, 2H), 2.43-2.39 (m, 2H), 1.94-1.81 (m, 4H); MS (EI): 530.2 (MH+).
- 10 [0762] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide: ¹H-NMR (400MHz, d6-DMSO): 10.5 (s, 1H), 9.60 (br s, 1H), 8.46 (d, 1H), 8.12 (d, 2H), 7.78-7.76 (m, 4H), 7.34 (m, 1H), 7.13 (br s, 2H), 3.70 (m, 1H), 3.54 (br s, 4H), 3.41 (m, 1H), 3.16 (br s, 4H), 2.23-2.08 (m, 3H), 1.95-1.74 (m, 3H), 0.84-0.83 (m, 4H); MS (EI): 497.2 (MH+).
- 15 [0763] N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}cyclopropanecarboxamide: ¹H-NMR (400MHz, d6-DMSO): 10.5 (s, 1H), 9.69 (br s, 1H), 8.47 (d, 1H), 8.13 (d, 2H), 7.77 (d, 4H), 7.36 (d, 2H), 7.23 (br s, 1H), 3.76 (br s, 4H), 3.25 (br s, 4H), 2.94 (septet, 1H), 1.84 (p, 1H), 1.03 (d, 6H), 0.84-0.83 (m, 4H). MS (EI): 485.1 (MH+).
- 20 [0764] 2,6-dichloro-N-{3-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)-amino]phenyl}benzamide: ¹H-NMR (400MHz, d6-DMSO): 10.7 (s, 1H), 10.5 (s, 1H), 9.74 (s, 1H), 8.50 (d, 1H), 8.40 (s, 1H), 8.21 (d, 2H), 7.76 (d, 2H), 7.61-7.59 (m, 2H), 7.53-7.49 (m, 1H), 7.47-7.45 (m, 1H), 7.39 (d, 1H), 7.31-7.22 (m, 2H), 1.83 (p, 1H), 0.83-0.81 (m, 4H); MS (EI): 518.1 (MH+).
- 25 [0765] 2,6-dichloro-N-(3-{{4-(1H-indol-5-yl)pyrimidin-2-yl}amino}phenyl)benzamide: ¹H-NMR (400MHz, d6-DMSO): 11.3 (s, 1H), 10.7 (s, 1H), 9.56 (s, 1H), 8.45 (s, 1H), 8.39 (d, 1H), 8.31 (s, 1H), 7.96 (dd, 1H), 7.54-7.52 (m, 2H), 7.46-7.41 (m, 3H), 7.37-7.34 (m, 2H), 7.21 (d, 2H), 6.48-6.48 (m, 1H). MS (EI): 474.0 (MH+).
- [0766] N-(4-{2-[(3-aminophenyl)amino]pyrimidin-4-yl}phenyl)-2-morpholin-4-ylacetamide: ¹H-NMR (400MHz, d6-DMSO): 11.0 (s, 1H), 10.4 (br, 2H), 9.89 (s, 1H), 8.56 (d, 1H), 8.23 (d, 2H), 7.82-7.79 (m, 3H), 7.62 (br, 1H), 7.44 (d, 1H), 7.34 (br, 1H), 6.8 (br, 1H), 4.24 (s, 2H), 3.96-3.84 (m, 8H); MS (EI): 405.3 (MH+).
- [0767] N-(4-phenylpyrimidin-2-yl)benzene-1,3-diamine: ¹H-NMR (400MHz, d6-DMSO): 9.37 (s, 1H), 8.51 (d, 1H), 8.19-8.16 (m, 2H), 7.57-7.53 (m, 3H), 7.37-7.36 (d, 1H), 7.10 (t, 1H), 7.00-6.91 (m, 2H), 6.22-6.20 (m, 1H), 5.00 (s, 2H). MS (EI): 263.3 (MH+).
- 35 [0768] N-[3-({4-[4-(acetyl-amino)-2-chlorophenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide: ¹H-NMR (400MHz, d6-DMSO): 10.7 (s, 1H), 10.3 (s, 1H), 9.78 (s,

5 1H), 8.52 (d, 1H), 8.13-8.12 (m, 1H), 7.93 (s, 1H), 7.71 (d, 1H), 7.30-7.21 (m, 5H), 7.30-7.21 (m, 2H), 7.11 (d, 1H), 2.07 (s, 3H). MS (EI): 527.9 (MH+).

[0769] 2,6-dichloro-N-{3-[(4-phenylpyrimidin-2-yl)amino]phenyl}benzamide:

¹H-NMR (400MHz, d6-DMSO): 10.7 (s, 1H), 9.76 (s, 1H), 8.56 (d, 1H), 8.39 (s, 1H), 8.26-8.23 (m, 2H), 7.61-7.59 (m, 2H), 7.55-7.48 (m, 5H), 7.44 (d, 1H), 7.31-7.24 (m, 2H); MS
10 (EI): 437.0 (MH+).

[0770] 4-(2,4-dichlorophenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:

¹H-NMR (400MHz, d6-DMSO): 9.59 (s, 1H), 8.52 (d, 1H), 7.80 (d, 1H), 7.66 (d, 1H), 7.63-7.58 (m, 3H), 7.00 (d, 1H), 6.88 (d, 2H), 3.74-3.72 (m, 4H), 3.04-3.01 (m, 4H); MS
(EI): 401.0 (MH+).

15 **[0771] 4-(2,4-dichlorophenyl)-N-{3-[(4-ethylpiperazin-1-**

yl)carbonyl]phenyl}pyrimidin-2-amine: ¹H-NMR (400MHz, d6-DMSO): 10.0 (s, 1H), 8.62 (d, 1H), 7.92 (s, 1H), 7.82 (d, 1H), 7.76 (dd, 1H), 7.70-7.68 (m, 1H), 7.62-7.59 (m, 1H), 7.34 (t, 1H), 7.13 (d, 1H), 6.96-6.94 (m, 1H), 3.59 (br s, 2H), 3.32 (br s, 2H), 2.37 (br s, 2H), 2.30 (q, 2H), 2.22 (br s, 2H), 0.99 (t, 3H); MS (EI): 456.0 (MH+).

20 **[0772] 2,6-dichloro-N-(3-{[4-(2,4-dichlorophenyl)pyrimidin-2-**

yl]amino}phenyl)benzamide: ¹H-NMR (400MHz, d6-DMSO): 10.7 (s, 1H), 9.89 (s, 1H), 8.59 (d, 1H), 8.14-8.13 (m, 1H), 7.80 (d, 1H), 7.77 (d, 1H), 7.59-7.48 (m, 5H), 7.32-7.23 (m, 2H), 7.14 (d, 1H); MS (EI): 504.9 (MH+).

[0773] N-(2-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide:

25 ¹H-NMR (400MHz, d6-DMSO): 11.3 (s, 1H), 9.61 (s, 1H), 8.53 (d, 1H), 8.19 (d, 1H), 7.79 (d, 1H), 7.50 (d, 2H), 7.48-7.44 (m, 1H), 7.22 (td, 1H), 7.14 (d, 1H), 6.94 (d, 2H), 3.75-3.73 (m, 4H), 3.06-3.03 (m, 4H), 1.69 (s, 3H). MS (EI): 390.1 (MH+).

[0774] 4-[3-(methyloxy)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine:

30 ¹H-NMR (400MHz, d6-DMSO): 9.45 (s, 1H), 8.49 (d, 1H), 7.73-7.66 (m, 4H), 7.45 (t, 1H), 7.34 (d, 1H), 7.13-7.10 (m, 1H), 6.92 (d, 2H), 3.86 (s, 3H), 3.75-3.35 (m, 4H), 2.51-2.50 (m, 4H). MS (EI): 363.1 (MH+).

[0775] 4-(2,3-dihydro-1,4-benzodioxin-6-yl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-

amine: ¹H-NMR (400MHz, d6-DMSO): 9.36 (s, 1H), 8.41 (d, 1H), 7.69-7.64 (m, 4H), 7.25 (d, 1H), 6.99 (d, 1H), 6.92 (d, 2H), 4.33-4.30 (m, 4H), 3.75-3.73 (m, 4H), 3.06-3.03 (m, 4H);
35 MS (EI): 391.1 (MH+).

[0776] 3-(methyloxy)-N-{4-[2-({4-[4-(piperazin-1-ylacetyl)piperazin-1-

yl]phenyl}amino)-pyrimidin-4-yl]phenyl}propanamide: ¹H-NMR (400MHz, d6-

- 5 DMSO): 10.2 (s, 1H), 9.43 (s, 1H), 8.79 (br, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.98 (d, 2H), 3.64 (t, 2H), 3.59 (br s, 2H), 3.25 (s, 3H), 3.22 (br m, 8H), 3.13 (br m, 4H), 3.07 (br m, 4H), 2.60 (t, 2H). MS (EI): 559.3 (MH⁺).
- [0777] N²,N²-dimethyl-N-[4-(2-[[4-(4-L-prolylpiperazin-1-yl)phenyl]amino]pyrimidin-4-yl)phenyl]glycinamide • 1.3 AcOH: ¹H-NMR (400MHz, d₆-DMSO): 10.00 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.90 (m, 1H), 3.64 (m, 4H), 3.11 (s, 2H), 3.10-2.98 (m, 5H), 2.66 (m, 1H), 2.29 (s, 6H), 2.07-1.99 (m, 1H), 1.89 (s, 4H), 1.73-1.54 (m, 3H); MS (EI) C₂₉H₃₆N₈O₂: 529.2 (MH⁺).
- 10 [0778] N²,N²-dimethyl-N-[4-(2-[[4-(4-D-prolylpiperazin-1-yl)phenyl]amino]pyrimidin-4-yl)phenyl]glycinamide • 1.4 AcOH: ¹H-NMR (400MHz, d₆-DMSO): 10.00 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.97 (m, 1H), 3.64 (m, 4H), 3.12 (s, 2H), 3.11-3.0 (m, 5H), 2.70 (m, 1H), 2.29 (s, 6H), 2.07-1.99 (m, 1H), 1.90 (s, 4H), 1.75-1.55 (m, 3H); MS (EI) C₂₉H₃₆N₈O₂: 529.2 (MH⁺).
- 15 [0779] 2-(dimethylamino)-N-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide • 3 AcOH: ¹H-NMR (400MHz, d₆-DMSO): 10.01 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.96 (d, 1H), 3.71 (m, 2H), 3.59 (m, 2H), 3.15 (s, 2H), 3.12 (s, 2H), 3.10 (m, 2H), 3.02 (m, 2H), 2.71 (m, 4H), 2.35 (m, 4H), 2.29 (s, 6H), 1.84 (s, 9H); MS (EI) C₃₀H₃₉N₉O₂: 558.5 (MH⁺).
- 20 [0780] 1,1-dimethylethyl [(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)methyl]carbamate: ¹H NMR (400MHz, CDCl₃): 8.20-8.22 (b, 1H), 8.05 (d, 2H), 7.65(d,1H), 7.50 (d, 2H), 7.25 (s, 1H), 7.23 (d, 2H), 7.05 (d, 2H), 5.01 (d, 1H), 4.40-4.44 (b, 2H), 3.90 (t, 4H), 3.20 (t, 4H), 1.50 (s, 9H); MS (EI) for C₂₆H₃₁N₅O₃: 462 (MH⁺).
- 25 [0781] 4-(4-(aminomethyl)phenyl)-N-(4-morpholinophenyl)pyrimidin-2-amine: ¹H NMR (400MHz, CD₃CN): 10.10-10.20(b,1H), 8.40 (d, 1H), 8.20(d,2H), 7.80 (d, 2H), 7.60 (d, 2H), 7.50 (d, 2H), 7.45(d,1H),7.20-7.22 (b, 2H), 4.40-4.44 (b, 2H), 3.90 (t, 4H), 3.20 (t, 4H); MS (EI) for C₂₁H₂₃N₅O: 362 (MH⁺).
- 30 [0782] methyl 4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzoate: ¹H NMR (400MHz, CDCl₃):8.45(s,1H), 8.20-8.30 (m, 4H), 7.65(d,1H), 7.25 (d, 2H), 7.15 (d, 2H), 6.85 (d, 1H), 4.01(s,3H), 3.90 (t, 4H), 3.20 (t, 4H); MS (EI) for C₂₂H₂₂N₄O₃: 391 (MH⁺).
- 35

- 5 [0783] 1-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-ethylurea: ¹H NMR (400MHz, d6-DMSO): 9.50-9.45 (b, 1H), 8.80(s,1H), 8.40(s,1H), 8.05 (d, 2H), 7.75-7.70 (m,4H), 7.30 (d, 1H), 7.05 (d, 2H),, 6.01 (d, 1H), 4.01(q,2H), 3.90 (m, 2H), 3.20 (m, 6H), 1.20 (s, 9H),1.10(t,3H); MS (EI) for C₂₈H₃₅N₇O₂: 502 (MH⁺).
- 10 [0784] 1-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-ethylurea: ¹H NMR (400MHz, d6-DMSO): 9.45 (s, 1H), 8.80(s,1H), 8.40(s,1H), 8.05 (d, 2H), 7.75-7.70 (m, 4H), 7.30 (d, 1H), 7.05 (d, 2H),, 6.01 (d, 1H), 3.80 (m, 4H),3.50(q, 2H), 3.40 (m, 4H),3.30 (m, 1H),3.20 (m, 6H),1.10(t,3H); MS (EI) for C₂₈H₃₃N₇O₂: 500 (MH⁺).
- 15 [0785] 1-ethyl-3-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}urea: ¹H NMR (400MHz, d6-DMSO): 9.45 (s, 1H), 8.80(s,1H), 8.40(s,1H), 8.05 (d, 2H), 7.75-7.70 (m, 4H), 7.30 (d, 1H), 7.05 (d, 2H),, 6.01 (d, 1H), 3.80 (m, 4H),3.30(q, 2H), 3.20 (m, 4H),3.10 (m, 1H),1.20 (m, 6H),1.10(t,3H); MS (EI) for C₂₇H₃₃N₇O₂: 488 (MH⁺).
- 20 [0786] N-ethyl-4-(4-{[4-(4-{[(ethylamino)carbonyl]amino}phenyl)pyrimidin-2-yl]amino}-phenyl)piperazine-1-carboxamide: ¹H NMR (400MHz, d6-DMSO): 9.45 (s, 1H), 8.80(s,1H), 8.40(m,1H), 8.05 (d, 2H), 7.75-7.70 (m, 4H), 7.30 (d, 1H), 7.05 (d, 2H),6.50 (s, 1H), 6.20 (d, 1H),3.40-3.50(m, 4H),3.00-3.15 (m, 8H),1.10-1.20(m,6H); MS (EI) for C₂₆H₃₂N₈O₂: 489 (MH⁺).
- 25 [0787] 1-ethyl-3-[4-(2-{[4-(4-L-prolylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-urea: ¹H NMR (400MHz, d6-DMSO): 10.20-10.25 (b, 1H), 9.45 (s, 1H), 8.40(m,1H), 8.05 (d, 2H), 7.75-7.60 (m, 6H), 7.30 (d, 1H), 7.00-6.90 (m, 2H),3.80-3.81 (m, 1H), 3.70-3.65(m, 4H),3.20-3.25 (m, 2H), 3.15-3.10 (m, 4H), 1.60-1.50 (m, 6H),1.10-1.20(m,3H); MS (EI) for C₂₈H₃₄N₈O₂: 515 (MH⁺).
- 30 [0788] 1-[4-(2-{[4-(4-L-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-3-ethyl-urea: ¹H NMR (400MHz, d6-DMSO): 9.70-9.65 (b, 1H), 9.25 (s, 1H), 8.40(m,1H), 8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 6.80-6.75 (m, 1H),3.80-3.81 (m, 1H), 3.70-3.65(m, 4H),3.60-3.55 (b, 2H), 3.25-3.20 (m, 6H),,1.10-1.20(m,6H); MS (EI) for C₂₆H₃₂N₈O₂: 489 (MH⁺).
- 35 [0789] 1-ethyl-3-{4-[2-({4-[4-(piperazin-1-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}urea: ¹H NMR (400MHz, d4-MeOH): 8.40 (m,

5 1H), 8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 3.80-3.81 (m, 4H), 3.30-3.10(m, 16H), 2.80-2.90 (m, 4H), 1.20(t, 3H); MS (EI) for C₂₉H₃₇N₉O₂: 544 (MH⁺).

[0790] 1-ethyl-3-[4-(2-[[4-(4-D-prolylpiperazin-1-yl)phenyl]amino}pyrimidin-4-

yl)phenyl]-urea: ¹H NMR (400MHz, d6-DMSO): 10.20-10.25 (b, 1H), 9.40-9.35 (b, 1H), 9.20 (s, 1H), 8.40(m, 1H), 8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m,

10 2H), 6.80-6.75 (b, 1H), 3.80-3.81 (m, 1H), 3.70-3.65(m, 4H), 3.20-3.25 (m, 2H), 3.15-3.10 (m, 4H), 1.60-1.50 (m, 6H), 1.10-1.20(m, 3H); MS (EI) for C₂₈H₃₄N₈O₂: 515 (MH⁺).

[0791] 1-[4-(2-[[4-(4-D-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-3-

ethyl-urea: ¹H NMR (400MHz, d6-DMSO): 11.00-10.90 (b, 1H), 9.25 (s, 1H), 8.40(m, 1H),

8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 6.80-6.75 (m, 1H), 3.80-3.81 (m, 1H), 3.70-3.65(m, 4H), 3.60-3.55 (b, 2H), 3.25-3.20 (m, 6H), 1.10-1.20(m, 6H); MS (EI) for C₂₆H₃₂N₈O₂: 489 (MH⁺).

[0792] (R)-N-(4-(2-(4-(4-(2-ethoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)-

phenyl)pyrrolidine-2-carboxamide: ¹H NMR (400MHz, d6-DMSO): 10.20-10.25 (b, 1H),

9.40 (s, 1H), 8.50 (d, 1H), 8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 4.20 (s, 2H), 3.80-3.81 (m, 2H), 3.70-3.65(m, 1H), 3.20-3.25 (m, 2H), 3.25-2.85 (m, 6H), 2.20 (m, 1H), 1.80-1.60 (m, 6H), 1.20(t, 3H); MS (EI) for C₂₉H₃₅N₇O₃: 530 (MH⁺).

[0793] N-[4-(2-[[4-(4-formylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-

prolinamide: ¹H NMR (400MHz, CDCl₃): 10.10-10.00 (b, 1H), 8.30 (d, 2H), 8.05 (s, 1H),

8.00 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 6.10-6.00 (b, 1H), 4.20 (m, 1H), 3.80-3.60 (m, 4H), 3.20-3.25 (m, 6H), 2.20 (m, 2H), 1.90-1.80 (m, 2H); MS (EI) for C₂₆H₂₉N₇O₂: 472 (MH⁺).

[0794] N-(4-{2-[[4-{4-(4-(dimethylamino)butanoyl]piperazin-1-

yl}phenyl]amino}pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400MHz, d6-DMSO):

10.20-10.25 (b, 1H), 9.40 (s, 1H), 8.50 (d, 1H), 8.05 (d, 2H), 7.75-7.60 (m, 4H), 7.30 (d, 1H), 7.00-6.90 (m, 2H), 4.20 (m, 1H), 3.80-3.60 (m, 4H), 3.20-3.25 (m, 6H), 2.90-2.85 (m, 4H), 2.40 (s, 3H), 2.30-2.22 (m, 2H), 2.20 (m, 3H), 2.05 (s, 3H), 1.90-1.80 (m, 2H); MS (EI) for C₃₁H₄₀N₈O₂: 557 (MH⁺).

[0795] N-(4-(2-(3-aminophenylamino)pyrimidin-4-yl)phenyl)-2-phenoxyacetamide:

¹H NMR (400 MHz, d6-DMSO): 10.41 (s, 1H), 12.4 (s, br, 1H), 8.56 (s, 1H), 8.19 (s, 1H),

35 7.97-7.82 (m, 3H), 7.61-7.26 (m, 5H), 7.07-7.02 (m, 3H), 6.98 (m, 1H), 4.79 (s, 2H). MS (EI): 412 (MH⁺).

- 5 [0796] **N-(4-(2-(4-(4-acetylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide:** ¹H NMR (400 MHz, d₄-MeOH): 10.22 (s, 1H), 9.41 (m, 1H), 8.12 (m, 2H), 7.75 (m, 2H), 7.68 (m, 2H), 7.27 (m, 1H), 3.72 (m, 4H), 6.94 (m, 2H), 3.58 (m, 4H), 3.09 (m, 2H), 3.02 (m, 2H), 2.09 (s, 3H), 2.03 (s, 3H). MS (EI): 431 (MH⁺).
- [0797] **N-(4-(2-(3-amino-2,4,5,6-tetrafluorophenylamino)pyrimidin-4-yl)phenyl)acetamide:** ¹H NMR (400 MHz, d₄-MeOH): 8.26 (m, 1H), 8.07 (m, 2H), 7.82 (m, 2H), 7.41 (m, 1H), 2.18 (s, 3H). MS (EI): 392 (MH⁺).
- 10 [0798] **N-(4-(2-(4-(piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide:** MS (EI) for C₂₂H₂₄N₆O: 389 (MH⁺).
- [0799] **1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopropane-carboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.27 (s, 1H), 10.19 (s, 1H), 9.23 (m, 3H), 8.60 (m, 1H), 8.09 (m, 2H), 7.95 (m, 3H), 7.79 (m, 2H), 7.54 (m, 1H), 4.11 (m, 4H), 3.65 (m, 4H), 1.71 (m, 2H), 1.42 (m, 2H). MS (EI): 431 (MH⁺).
- 15 [0800] **(S)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)indoline-2-carboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 9.50 (m, 1H), 8.43 (m, 1H), 8.08 (m, 2H), 7.92 (m, 2H), 7.84 (m, 2H), 7.31 (m, 1H), 7.10-6.88 (m, 4H), 6.61 (m, 2H), 6.07 (m, 1H), 4.42 (m, 1H), 3.75 (m, 4H), 3.18-2.99 (m, 6H). MS (EI): 493 (MH⁺).
- 20 [0801] **N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.34 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 3.95 (t, 1H), 3.82-3.69 (m, 7H), 3.25-3.16 (m, 1H), 3.05 (t, 4H), 2.13-2.06 (m, 2H). MS (EI): 446 (MH⁺).
- 25 [0802] **N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyridin-3-yl)-acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.53 (s, 1H), 9.34 (s, 1H), 8.54 (s, 1H), 8.47 (d, 1H), 8.33 (d, 1H), 8.12 (d, 2H), 7.76 (d, 3H), 7.67 (d, 2H), 7.37 (m, 1H), 7.28 (d, 1H), 6.93 (d, 2H), 3.76-3.73 (m, 6H), 3.06-3.03 (m, 4H). MS (EI): 467 (MH⁺).
- 30 [0803] **1-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea:** ¹H NMR (400 MHz, d₆-DMSO): 9.33 (s, 1H), 9.00 (s, 1H), 8.40 (d, 1H), 8.04 (d, 2H), 7.66 (d, 2H), 7.55 (d, 2H), 7.23 (d, 1H), 6.93 (d, 2H), 6.39 (t, 1H), 3.17-3.08 (m, 10H), 2.85 (s, 6H), 2.74 (s, 4H), 1.08-1.04 (m, 9H). MS (EI): 531 (MH⁺).
- 35 [0804] **(R)-N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 12.64 (s, 1H), 9.35 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.69-7.63 (m, 4H), 7.28 (d,

5 1H), 6.89 (d, 2H), 3.30-3.23 (m, 1H), 3.15 (d, 2H), 3.10-3.04 (m, 4H), 2.62-2.58 (m, 4H), 2.34-2.28 (m, 1H), 2.21 (s, 6H), 2.18 (s, 2H), 2.10 (s, 2H), 1.82-1.64 (m, 4H), 0.84 (s, 6H). MS (EI): 557 (MH+).

[0805] (R)-2-amino-N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H NMR (400 MHz, d₆-DMSO):
10 9.35 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.80 (d, 2H), 7.63 (d, 2H), 7.27 (d, 1H), 6.89 (d, 2H), 3.49-3.43 (m, 1H), 3.05-3.01 (m, 4H), 2.62-2.58 (m, 4H), 2.19 (s, 6H), 2.15 (s, 2H), 2.07 (s, 2H), 1.21 (d, 3H), 0.82 (s, 6H). MS (EI): 531 (MH+).

[0806] N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide: ¹H NMR (400 MHz, d₆-DMSO):
15 10.31 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.78 (d, 2H), 7.66 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.63 (t, 2H), 3.34 (s, 4H), 3.25 (s, 3H), 3.13 (s, 4H), 2.84 (s, 6H), 2.74 (s, 4H), 2.60 (t, 2H), 1.05 (s, 6H). MS (EI): 546 (MH+).

[0807] 2-(dimethylamino)-N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO):
20 9.98 (s, 1H), 9.33 (s, 1H), 8.41 (d, 1H), 8.08 (d, 2H), 7.81 (d, 2H), 7.62 (d, 2H), 7.24 (d, 1H), 6.88 (d, 2H), 3.09 (s, 2H), 3.05-3.02 (m, 4H), 2.60-2.45 (m, 4H), 2.27 (s, 6H), 2.20 (s, 6H), 2.15 (s, 2H), 2.09 (s, 2H), 0.82 (s, 6H). MS (EI): 545 (MH+).

[0808] N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide: ¹H NMR (400 MHz, d₆-DMSO):
25 10.18 (s, 1H), 9.36 (s, 1H), 8.41 (d, 1H), 8.08 (d, 2H), 7.75 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 3.12 (s, 4H), 3.07 (s, 4H), 2.84 (s, 6H), 2.73 (s, 4H), 2.31 (t, 2H), 1.66-1.58 (m, 2H), 1.03 (s, 6H), 0.91 (t, 3H). MS (EI): 530 (MH+).

[0809] (3S,7S)-7-(hydroxymethyl)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)quinuclidine-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO):
30 10.25 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.74 (t, 4H), 3.65-3.58 (m, 2H), 3.45-3.37 (m, 3H), 3.05 (t, 4H), 2.94-2.88 (m, 3H), 2.71-2.67 (m, 1H), 2.17 (s, 1H), 1.68-1.63 (m, 2H), 1.50-1.46 (m, 1H), 1.27-1.21 (m, 1H). MS (EI): 515 (MH+).

[0810] (R)-N-(4-(2-(3-ethoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide: ¹H NMR (400 MHz, d₆-DMSO):
35 11.45 (s, 1H), 10.04 (s, 1H), 8.59 (d, 1H), 8.21 (d, 2H), 7.87 (d, 2H), 7.49 (d, 1H), 7.44-7.41 (m, 2H), 7.32 (s, 1H), 7.19 (s,

5 1H), 4.51-4.45 (m, 2H), 4.28-4.25 (m, 4H), 4.09-4.01 (m, 4H), 3.68-3.52 (m, 3H), 3.33-3.23 (m, 2H), 2.49-2.42 (m, 1H), 2.03-1.91 (m, 3H), 1.49 (t, 3H). MS (EI): 489 (MH+).

[0811] N-{4-[2-({4-morpholin-4-yl-3-[(phenylmethyl)oxy]phenyl}amino)pyrimidin-4-yl]-phenyl}-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 11.57 (s, 1H), 10.17 (s, 1H), 8.59 (d, 1H), 8.23 (d, 2H), 8.09 (s, 1H), 7.90 (m, 3H), 7.59 (m, 2H), 7.48 (m, 5H), 5.34 (s, 10 2H), 4.51 (m, 4H), 4.06 (m, 5H), 3.29 (m, 3H), 1.98 (m, 3H). MS (EI) for C₃₂H₃₄N₆O₃: 551.7 (MH+).

[0812] 4-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-piperazine-1-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.35 (s, 1H), 8.88 (s, 1H), 8.41 (d, 1H), 8.15 (s, 1H), 8.06 (d, 1H), 7.66 (m, 3H), 7.25 (d, 1H), 6.93 (d, 2H), 3.74 (m, 15 4H), 3.53 (m, 8H), 3.04 (m, 4H), 3.32 (m, 3H). MS (EI) for C₂₆H₃₁N₇O₂: 474.6 (MH+).

[0813] 1-[3-(dimethylamino)propyl]-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea: ¹H NMR (400 MHz, d6-DMSO): 9.34 (s, 1H), 9.22 (s, 1H), 8.40 (d, 1H), 8.05 (d, 2H), 7.67 (d, 2H), 7.57 (d, 2H), 7.24 (d, 1H), 6.93 (d, 2H), 6.66 (t, 1H), 3.74 (m, 4H), 3.17 (m, 2H), 3.05 (m, 4H), 2.90 (t, 2H), 2.62 (s, 6H), 1.78 20 (m, 2H). MS (EI) for C₂₆H₃₃N₇O₂: 476.6 (MH+).

[0814] 1-[3-(methyloxy)propyl]-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea: ¹H NMR (400 MHz, d6-DMSO): 9.3 (s, 1H), 8.87 (s, 1H), 8.40 (d, 1H), 8.05 (d, 2H), 7.67 (d, 2H), 7.55 (d, 2H), 7.24 (d, 1H), 6.93 (d, 2H), 6.37 (t, 1H), 3.74 (m, 4H), 3.38 (d, 2H), 3.25 (s, 3H), 3.15 (m, 2H), 3.04 (m, 4H), 1.68 (m, 2H). MS (EI) for 25 C₂₅H₃₀N₆O₃: 463.6 (MH+).

[0815] 1-(2-morpholin-4-ylethyl)-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea: ¹H NMR (400 MHz, d6-DMSO): 9.34 (s, 1H), 9.00 (s, 1H), 8.40 (d, 1H), 8.17 (s, 1H), 8.05 (d, 2H), 7.68 (d, 2H), 7.55 (d, 2H), 7.24 (d, 1H), 6.93 (d, 2H), 6.25 (t, 1H), 3.74 (m, 4H), 3.60 (m, 4H), 3.22 (m, 2H), 3.04 (m, 4H), 2.40 (m, 5H). MS (EI) for 30 C₂₇H₃₃N₇O₃: 504.5 (MH+).

[0816] 1-[2-(dimethylamino)ethyl]-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea: ¹H NMR (400 MHz, d6-DMSO): 9.33 (s, 1H), 9.06 (s, 1H), 8.40 (d, 1H), 8.21 (s, 1H), 8.04 (d, 2H), 7.67 (d, 2H), 7.55 (d, 2H), 7.23 (d, 1H), 6.93 (d, 2H), 6.33 (t, 1H), 3.74 (d, 4H), 3.21 (m, 2H), 3.05 (m, 1H), 2.38 (t, 2H), 2.12 35 (s, 6H). MS (EI) for C₂₅H₃₁N₇O₂: 462.5 (MH+).

[0817] 1-ethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 9.53 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.14

5 (d, 2H), 7.85 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 1H), 3.75 (m, 4H), 3.21 (m, 1H), 3.09 (m, 4H), 2.65 (m, 1H), 2.54 (m, 2H), 2.35 (m, 1H), 2.14 (m, 1H), 1.79 (m, 3H), 1.08 (t, 3H). MS (EI) for $C_{27}H_{32}N_6O_2$: 473.6 (MH⁺).

[0818] 1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)-D-prolinamide: ¹H NMR (400 MHz, d6-DMSO): 10.31 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.81 (d, 2H), 7.67 (d, 2H), 7.30 (d, 1H), 6.93 (d, 1H), 5.05 (s, br, 1H), 3.75 (m, 4H), 3.05 (m, 4H), 2.75 (m, 2H), 2.63 (m, 1H), 2.40 (m, 2H), 2.18 (m, 2H), 1.80 (m, 3H). MS (EI) for $C_{27}H_{32}N_6O_3$: 489.5 (MH⁺).

[0819] N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.39 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.79 (d, 2H), 7.66 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 3.94 (t, 1H), 3.75 (m, 4H), 3.15 (m, 8H), 2.80 (m, 9H), 2.09 (m, 2H), 1.03 (s, 7H). MS (EI) for $C_{32}H_{43}N_7O_2$: 558.7 (MH⁺).

[0820] (2R)-N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.88 (s, 1H), 9.34 (s, 1H), 8.52 (s, 1H), 8.38 (s, 1H), 8.00 (s, 1H), 7.56 (m, 2H), 7.22 (d, 1H), 6.96 (d, 2H), 4.43 (m, 1H), 4.00 (m, 1H), 3.86 (m, 1H), 3.05 (m, 7H), 2.60 (m, 7H), 2.20 (m, 10H), 0.85 (s, 6H). $C_{32}H_{43}N_7O_2$. MS (EI) for $C_{32}H_{43}N_7O_2$: 558.7 (MH⁺).

[0821] N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl)cyclopropanecarboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.47 (s, 1H), 9.36 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.75 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.90 (d, 2H), 3.05 (m, 4H), 2.61 (m, 4H), 2.21 (s, 6H), 2.17 (s, 2H), 2.10 (s, 2H), 1.90 (s, 1H), 1.82 (m, 1H), 0.84 (m, 9H). MS (EI) for $C_{31}H_{41}N_7O$: 528.6 (MH⁺).

[0822] (S)-N-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenyl)-amino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.95 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.88 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 4.44 (t, 1H), 4.01 (m, 1H), 3.85 (m, 1H), 3.14 (m, 6H), 2.86 (s, 6H), 2.77 (m, 4H), 2.21 (m, 2H), 2.01 (m, 2H), 1.90 (m, 2H), 1.05 (s, 6H). MS (EI) for $C_{32}H_{43}N_7O_2$: 558.7 (MH⁺).

35 [0823] N-(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)-phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.35 (s, 1H), 9.41 (s, 1H), 8.44 (s, 1H), 8.14 (d, 2H), 7.78 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d,

5 2H), 3.96 (m, 1H), 3.73 (m, 3H), 3.62 (m, 4H), 3.20 (m, 1H), 3.04 (m, 5H), 2.79 (m, 1H), 2.62 (t, 2H), 2.11 (m, 2H), 2.79 (m, 3H), 1.55 (m, 3H). MS (EI) for $C_{31}H_{37}N_7O_3$: 556.6 (MH⁺).

[0824] 1-(1-methylethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 9.98 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.14 (m, 1H), 3.05 (m, 4H), 2.81 (1H), 2.54 (m, 2H), 2.08 (m, 1H), 1.77 (m, 1H), 1.75 (m, 2H), 1.05 (m, 6H). MS (EI) for $C_{28}H_{34}N_6O_2$: 487.6 (MH⁺).

[0825] 1-ethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 9.95 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.85 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.74 (m, 4H), 3.20 (m, 1H), 3.07 (m, 5H), 2.64 (m, 1H), 2.54 (m, 1H), 2.35 (m, 1H), 2.14 (m, 1H), 1.79 (m, 3H), 1.08 (t, 3H). MS (EI) for $C_{27}H_{32}N_6O_2$: 473.5 (MH⁺).

[0826] 2-(2-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.50 (s, 1H), 9.32 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.76 (d, 2H), 7.67 (d, 1H), 7.41 (m, 1H), 7.34 (m, 1H), 7.28 (d, 1H), 7.18 (m, 2H), 6.93 (d, 2H), 3.79 (s, 2H), 3.74 (m, 4H), 3.05 (m, 4H). MS (EI) for $C_{28}H_{26}FN_5O_2$: 484.5 (MH⁺).

[0827] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.76 (s, 1H), 9.42 (s, 1H), 8.82 (d, 2H), 8.47 (d, 1H), 8.20 (d, 2H), 7.96 (s, 2H), 7.89 (d, 2H), 7.68 (d, 2H), 7.32 (d, 1H), 6.95 (d, 2H), 3.75 (m, 4H), 3.07 (m, 4H). MS (EI) for $C_{26}H_{24}N_6O_2$: 453.5 (MH⁺).

[0828] (R)-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)-phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.19 (s, br, 1H), 9.25 (s, br, 1H), 8.31 (s, 1H), 7.81-7.58 (m, 6H), 7.09-6.76 (br m, 3H), 3.79 (m, 3H), 3.66 (s, 3H), 3.02-2.92 (m, 4H), 2.20 (m, 4H), 2.09 (m, 2H), 2.00 (m, 1H), 1.82 (m, 1H), 1.70 (m, 1H), 12.4 (s, 3H). MS (EI): 552 (MH⁺).

[0829] (R)-2-amino-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, br, 1H), 9.24 (s, br, 1H), 8.31 (s, 1H), 7.80 (d, 2H), 7.66-7.69 (m, 4H), 7.09 (s, 1H), 6.85 (m, 2H), 6.76 (s, 1H), 3.80 (m, 3H), 3.66 (m, 3H), 3.00 (m, 4H), 2.95 (m, 4H), 2.22 (s, 3H), 1.24 (s, 3H). MS (EI): 526 (MH⁺).

- 5 [0830] (S)-2-amino-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, br, 1H), 9.24 (s, br, 1H), 8.31 (s, 1H), 7.79 (d, 2H), 7.67-7.58 (m, 4H), 7.09 (s, 1H), 6.86 (m, 2H), 6.76 (s, 1H), 3.80 (m, 3H), 3.66 (m, 3H), 3.01 (m, 4H), 2.95 (m, 4H), 2.22 (s, 3H), 1.23 (s, 3H). MS (EI) for C₂₉H₃₅N₉O: 526 (MH⁺).
- 10 [0831] N-[3-({2-[(4-morpholin-4-yl)phenyl]amino}-7H-pyrrolo[2,3-d]pyrimidin-4-yl)-amino]phenyl]acetamide: ¹H NMR (400MHz, d₆-DMSO): 11.14 (s, 1H), 9.87 (s, 1H), 9.15 (s, 1H), 8.47 (s, 1H), 8.04 (s, 1H), 7.80-7.73 (m, 1H), 7.70-7.63 (m, 2H), 7.25-7.18 (m, 2H), 6.89-6.81 (m, 3H), 6.67-6.64 (m, 1H), 3.77-3.71 (m, 4H), 3.04-2.98 (m, 4H), 2.06 (s, 3H). MS (EI) for C₂₄H₂₅N₇O₂: 444 (MH⁺).
- 15 [0832] N-(4-{2-[(2-methyl-4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.36 (br s, 1H), 9.51 (br s, 1H), 8.34 (d, 1H), 8.08 (d, 2H), 7.74 (d, 2H), 7.39 (d, 2H), 7.05 (br d, 2H), 3.80 (s, 4H), 3.22 (s, 4H), 2.21 (s, 3H), 2.07 (s, 3H). MS (EI): (MH⁺).
- [0833] N-(4-{2-[(4-pyrrolidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.33 (br s, 1H), 9.72 (br s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.69 (br s, 2H), 7.34 (d, 1H), 3.37 (m, 4H), 2.10 (s, 3H), 2.03 (s, 4H). MS (EI): 374 (MH⁺).
- 20 [0834] N-[4-(2-{[4-(diethylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.34 (br s), 9.99 (br s, 1H), 8.56 (d, 1H), 8.12 (d, 2H), 8.06 (d, 2H), 7.79 (d, 2H), 7.72 (d, 2H), 7.44 (d, 1H), 3.49 (q, 4H), 2.10 (s, 3H), 1.05 (dt, 6H). MS (EI): 376 (MH⁺).
- 25 [0835] N-(4-{2-[(4-azepan-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.34 (br s), 9.99 (br s, 1H), 8.56 (d, 1H), 8.12 (d, 2H), 8.06 (d, 2H), 7.79 (d, 2H), 7.72 (d, 2H), 7.44 (d, 1H), 3.54 (m, 4H), 2.09 (s, 3H), 1.91 (m, 2H), 1.70 (m, 2H), 1.64 (m, 4H), 1.44 (m, 2H), 1.37 (m, 2H). MS (EI): 402 (MH⁺).
- 30 [0836] N-{4-[2-({4-[methyl(2-phenylethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl} acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.34 (br s, 1H), 9.95 (br s, 1H), 8.54 (d, 1H), 8.15 (d, 2H), 7.98 (m, 1H), 7.79 (d, 2H), 7.43 (d, 1H), 7.31 (m, 3H), 7.23 (m, 4H), 3.71 (m, 2H), 3.13 (m, 2H), 2.10 (s, 1H), 1.99 (s, 3H). MS (EI): 438 (MH⁺).
- 35 [0837] N-[4-(2-{[4-(1,4-dioxo-8-azaspiro[4.5]dec-8-yl)phenyl]amino}pyrimidin-4-yl)phenyl] acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.34 (br s), 9.99 (br s, 1H), 8.56

5 (d, 1H), 8.12 (d, 2H), 8.06 (d, 2H), 7.79 (d, 2H), 7.72 (d, 2H), 7.44 (d, 1H), 3.90 (s, 4H), 2.70 (t, 4H), (2.10 (s, 3H), 1.76 (t, 4H). MS (EI): 446 (MH+).

[0838] N-[4-(2-[[4-(2-oxopiperidin-1-yl)phenyl]amino]pyrimidin-4-

yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.32 (br s, 1H), 9.86 (br s, 1H), 8.51 (d, 1H), 8.14 (d, 2H), 7.85 (t, 4H), 7.40 (d, 1H), 7.22 (d, 2H), 3.59 (m, 2H), 2.38 (t, 2H),
10 2.09 (s, 3H), 1.85 (m, 4H). MS (EI): 402 (MH+).

[0839] N-[4-(2-[[4-(2-methylpiperidin-1-yl)phenyl]amino]pyrimidin-4-yl)phenyl]-

acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (br s, 1H), 9.36 (br s, 1H), 8.43 (s, 1H), 8.27 (s, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (s, 1H), 6.92 (d, 2H), 2.09 (s, 3H), 3.41 (m, 3H), 1.60 (m, 6H), 0.88 (d, 3H). MS (EI): 402 (MH+).

15 [0840] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-

valinamide: ¹H NMR (400 MHz, d6-DMSO): 11.51 (br s, 1H), 10.16 (br s, 1H), 8.57 (s, 1H), 8.48 (m, 2H), 8.20 (m, 2H), 7.93 (m, 3H), 7.78 (m, 1H), 7.50 (s, 1H), 5.45 (br s, 4H), 4.07 (s, 4H), 3.53 (s, 4H), 3.35 (m, 1H), 2.25 (m, 1H), 1.03 (m, 6H). MS (EI): 447 (MH+).

[0841] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-

20 valinamide: ¹H NMR (400 MHz, d6-DMSO): 11.51 (br s, 1H), 10.16 (br s, 1H), 8.57 (s, 1H), 8.48 (m, 2H), 8.20 (m, 2H), 7.93 (m, 3H), 7.78 (m, 1H), 7.50 (s, 1H), 5.45 (br s, 4H), 4.07 (s, 4H), 3.53 (s, 4H), 3.35 (m, 1H), 2.25 (m, 1H), 1.03 (m, 6H). MS (EI): 447 (MH+).

[0842] 2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-

alaninamide: ¹H NMR (400 MHz, d6-DMSO): 10.68 (br s, 1H), 10.02 (br s, 1H), 8.53 (m, 2H), 8.18 (d, 2H), 7.95 (d, 2H), 7.89 (d, 2H), 7.66 (m, 1H), 7.47 (d, 1H), 5.20 (br s, 4H), 4.01 (s, 4H), 3.44 (s, 4H), 1.66 (6H). MS (EI): 433 (MH+).

[0843] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-

yl}phenyl)tryptophanamide: ¹H NMR (400 MHz, d6-DMSO): 11.37 (s, 1H), 10.07 (s, 1H), 10.03 (s, 1H), 8.56 (d, 1H), 8.42 (d, 2H), 8.19 (d, 2H), 7.91 (d, 2H), (d, 2H), 7.73 (d, 1H),
30 7.66 (1H), 7.46 (d, 1H), 7.35 (d, 1H), 7.28 (d, 1H), 7.07 (t, 1H), 6.95 (t, 1H), 4.70 (br s, 4H), 4.34 (m, 1H), 4.03 (s, 4H), 3.49 (s, 4H), 3.36 (dq, 2H). MS (EI): 534 (MH+).

[0844] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-

prolinamide: ¹H NMR (400 MHz, d6-DMSO): 11.43 (br s, 1H), 10.07 (br s, 2H), 8.73 (d, 1H), 8.57 (d, 1H), 8.21 (d, 2H), 7.91 (d, 2H), 7.98 (d, 2H), 7.71 (br s, 2H), 7.48 (d, 1H), 4.48
35 (m, 1H), 4.08 (s, 4H), 3.74 (m, 4H), 3.42 (m, 1H), 3.36 (m, 1H), 3.04 (m, 4H), 2.22 (m, 1H), 1.90 (m, 2H), 1.82 (m, 2H). MS (EI): 445 (MH+).

- 5 [0845] **N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1,2,3,4-tetrahydro-isoquinoline-1-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 11.30 (br d, 1H), 10.04 (br s, 1H), 8.56 (d, 1H), 8.39 (s, 3H), 8.20 (d, 2H), 7.90 (m, 2H), 7.87 (m, 2H), 7.67 (m, 3H), 7.47 (d, 1H), 5.00 (br s, 3H), 4.65 (s, 1H), 4.20 (m, 2H), 4.03 (s, 4H), 3.97 (m, 1H), 3.94 (m, 2H), 3.80 (m, 1H), 3.49 (s, 4H). MS (EI): 507 (MH+).
- 10 [0846] **O-(1,1-dimethylethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)-L-serinamide:** ¹H NMR (400 MHz, d6-DMSO): 12.11 (br s, 1H), 10.65 (br s, 1H), 10.12 (s, 1H), 9.60 (s, 1H), 8.58 (d, 1H), 8.23 (d, 2H), 7.95 (d, 2H), 7.79 (s, 1H), 7.56 (d, 1H), 7.49 (d, 1H), 7.31 (s, 2H), 5.14 (br s, 4H), 4.06 (s, 4H), 3.79 (m, 1H), 3.54 (s, 4H), 3.45 (m, 1H), 3.15 (q, 1H), 1.21 (s, 9H). MS (EI): 491 (MH+).
- 15 [0847] **3-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydro-furan-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.89 (br s, 1H), 9.92 (br s, 1H), 8.83 (s, 2H), 8.55 (d, 1H), 8.21 (d, 2H), 7.94 (d, 2H), 7.87 (d, 1H), 7.53 (s, 1H), 7.43 (d, 1H), 4.30 (br s, 4H), 4.21 (d, 1H), 4.07 (d, 1H), 4.05 (m, 1H), 4.02 (m, 1H), 3.97 (s, 4H), 3.42 (s, 4H), 2.79 (m, 1H), 2.28 (m, 1H). MS (EI): 461 (MH+).
- 20 [0848] **bis(1,1-dimethylethyl) (2R)-2-[[[4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]carbonyl]piperazine-1,4-dicarboxylate:** ¹H NMR (400 MHz, d6-DMSO): 10.41 (br s, 1H), 9.35 (s, 1H), 8.42 (d, 1H), 8.14 (d, 2H), 8.76 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 4.51 (m, 1H), 3.90 (m, 2H), 3.74 (m, 4H), 3.66 (t, 4H), 3.04 (t, 4H), 1.41 (s, 3H), 1.33 (s, 9H), 1.17 (s, 6H). MS (EI): 660 (MH+).
- 25 [0849] **N-(4-{2-[(4-{4-[2-(2-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.70 (d, 2H), 7.29 (m, 3H), 7.16 (m, 2H), 6.98 (d, 2H), 3.80 (s, 2H), 3.69 (m, 2H), 3.63 (m, 2H), 3.09 (m, 2H), 3.04 (m, 2H), 2.09 (s, 3H). MS (EI): 525.5 (MH+).
- 30 [0850] **N-(4-{2-[(4-{4-[2-(2-methylphenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 7.12 (m, 4H), 6.97 (d, 2H), 3.74 (s, 2H), 3.65 (m, 4H), 3.05 (m, 4H), 2.20 (s, 3H), 2.09 (s, 3H). MS (EI): 521.6 (MH+).
- 35 [0851] **N-(4-{2-[(4-{4-[2-(3-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.69 (d, 2H), 7.35 (m,

5 1H), 7.28 (d, 1H), 7.08 (m, 3H), 6.96 (d, 2H), 3.81 (s, 2H), 3.64 (m, 4H), 3.02 (m, 4H), 2.09 (s, 3H). MS (EI): 525.4 (MH⁺).

[0852] N-{4-[2-({4-[4-(3-thienylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.83 (m, 1H), 7.75 (d, 2H), 7.70 (d, 2H), 7.64 (m, 1H), 7.25 (m, 2H), 6.98 (d, 2H), 3.68 (m, 4H), 3.11 (m, 4H), 2.09 (s, 3H). MS (EI): 499.4 (MH⁺).

[0853] N-(4-{2-[(4-{4-[(6-chloropyridin-3-yl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.53 (m, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.98 (dd, 1H), 7.75 (d, 2H), 7.70 (d, 2H), 7.65 (d, 1H), 7.28 (d, 1H), 6.98 (d, 2H), 3.78 (m, 2H), 3.48 (m, 2H), 3.17 (m, 2H), 3.08 (m, 2H), 2.09 (s, 3H). MS (EI): 529.1 (MH⁺).

[0854] N-(4-{2-[(4-{4-[(3-methylfuran-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75-7.68 (m, 5H), 7.28 (d, 1H), 6.98 (d, 2H), 6.52 (d, 1H), 3.73 (m, 4H), 3.11 (m, 4H), 2.17 (s, 3H), 2.09 (s, 3H). MS (EI): 497.6 (MH⁺).

20 [0855] N-(4-{2-[(4-{4-[(3-fluoro-2-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for C₃₀H₂₉FN₆O₂: 525.5 (MH⁺).

[0856] N-(4-{2-[(4-{4-[(imidazol-4-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for C₂₆H₂₆N₈O₂: 483.5 (MH⁺).

25 [0857] N-(4-{2-[(4-{4-[(2-methoxypyridin-3-yl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.26 (dd, 1H), 8.12 (d, 2H), 7.75-7.67 (m, 5H), 7.28 (d, 1H), 7.09 (dd, 1H), 6.97 (d, 2H), 3.90 (s, 3H), 3.77 (m, 2H), 3.29 (m, 2H), 3.14 (m, 2H), 3.04 (m, 2H), 2.09 (s, 3H). MS (EI): 524.6 (MH⁺).

[0858] N-(4-{2-[(4-{4-[(4-fluoro-3-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for C₃₀H₂₉FN₆O₂: 525.5 (MH⁺).

35 [0859] N-{4-[2-({4-[4-(naphthalen-2-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.20 (s, 1H), 9.37 (s, 1H), 8.50 (d, 1H), 8.42 (d, 1H), 8.22 (dd, 2H), 8.09 (dd, 3H), 7.82-7.71

5 (m, 5H), 7.64 (d, 2H), 7.26 (d, 1H), 6.89 (d, 2H), 3.14 (m, 4H), 3.11 (m, 4H), 2.08 (s, 3H). MS (EI): 579.6 (MH⁺).

[0860] N-{4-[2-({4-[4-(quinolin-8-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.37 (s, 1H), 9.10 (dd, 1H), 8.56 (dd, 1H), 8.42 (m, 2H), 8.34 (dd, 1H), 8.10 (d, 2H), 7.79 (m, 1H), 7.73 (m, 10 3H), 7.65 (d, 2H), 7.27 (d, 1H), 6.90 (d, 2H), 3.45 (m, 4H), 3.08 (m, 4H), 2.09 (s, 3H). MS (EI): 580.8 (MH⁺).

[0861] N-[4-(2-{{4-(4-{[4-(1,1-dimethylethyl)phenyl]sulfonyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.20 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.70 (m, 4H), 7.66 (d, 15 2H), 7.27 (d, 1H), 6.91 (d, 2H), 3.16 (m, 4H), 3.01 (m, 4H), 2.08 (s, 3H), 1.32 (s, 9H). MS (EI): 585.5 (MH⁺).

[0862] N-[4-(2-{{4-(4-{[5-bromo-2-(methyloxy)phenyl]sulfonyl}piperazin-1-yl)phenyl]-amino}pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.09 (dd, 2H), 7.84 (m, 2H), 7.74 (d, 2H), 7.68 (d, 20 2H), 7.28 (d, 2H), 6.94 (d, 2H), 3.92 (s, 3H), 3.27 (m, 4H), 3.11 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₉BrN₆O₄S: 638.6 (MH⁺).

[0863] N-(4-{2-[(4-{4-[(phenylmethyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.69 (d, 2H), 7.42 (m, 25 5H), 7.28 (d, 1H), 6.96 (d, 2H), 4.50 (s, 2H), 3.20 (m, 4H), 3.06 (m, 4H), 2.08 (s, 3H). MS (EI): 543.6 (MH⁺).

[0864] N-[4-(2-{{4-(4-{[3-(trifluoromethyl)phenyl]sulfonyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.18-8.08 (m, 4H), 8.02 (s, 1H), 7.97 (d, 1H), 7.74 30 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.92 (d, 2H), 3.15 (m, 4H), 3.10 (m, 4H), 2.09 (s, 3H). MS (EI): 597.7 (MH⁺).

[0865] N-(4-{2-[(4-{4-[(2-methylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.09 (dd, 2H), 7.85 (dd, 1H), 7.74 (d, 2H), 7.67-7.59 35 (m, 3H), 7.47 (m, 2H), 7.27 (d, 1H), 6.94 (d, 2H), 3.17 (m, 4H), 3.13 (m, 4H), 2.61 (s, 3H), 2.09 (s, 3H). MS (EI): 543.7 (MH⁺).

- 5 **[0866]** N-(4-{2-[(4-{4-[(3-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.10 (dd, 2H), 7.73 (m, 3H), 7.67-7.62 (m, 5H), 7.27 (d, 1H), 6.92 (d, 2H), 3.14 (m, 4H), 3.08 (m, 4H), 2.09 (s, 3H). MS (EI): 547.7 (MH⁺).
- [0867]** N-(4-{2-[(4-{4-[(2,4-difluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for C₂₈H₂₆F₂N₆O₃S: 565.6 (MH⁺).
- 10 **[0868]** N-{4-[2-({3-[4-({4-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.47 (s, 1H), 8.49 (d, 1H), 8.14 (dd, 2H), 7.75 (d, 2H), 7.64 (s, 1H), 7.48 (dd, 2H), 7.33 (m, 3H), 7.22 (m, 1H), 7.13 (m, 1H), 6.56 (dd, 1H), 3.57 (s, 2H), 3.16 (m, 4H), 2.54 (m, 4H), 2.09 (s, 3H). MS (EI): 563.6 (MH⁺).
- 15 **[0869]** N-(4-{2-[(3-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.45 (s, 1H), 8.49 (d, 1H), 8.13 (dd, 2H), 7.75 (d, 2H), 7.58 (s, 1H), 7.33 (d, 1H), 7.25 (dd, 1H), 7.15-7.09 (m, 2H), 6.77 (d, 1H), 6.56 (dd, 1H), 3.68 (s, 3H), 3.58 (s, 2H), 3.12 (m, 4H), 2.54 (m, 4H), 2.09 (s, 3H). MS (EI): 483.5 (MH⁺).
- 20 **[0870]** N-{4-[2-({3-[4-({2-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)-amino]pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.47 (s, 1H), 8.50 (d, 1H), 8.13 (dd, 2H), 7.75 (d, 2H), 7.64-7.62 (m, 2H), 7.44-7.36 (m, 4H), 7.23 (dd, 1H), 7.14 (m, 1H), 6.55 (dd, 1H), 3.62 (s, 2H), 3.16 (m, 4H), 2.57 (m, 4H), 2.09 (s, 3H). MS (EI): 563.6 (MH⁺).
- 25 **[0871]** N-(4-{2-[(3-{4-[(3-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.69 (d, 2H), 7.44-7.38 (m, 4H), 7.28 (d, 1H), 6.96 (d, 2H), 4.48 (s, 2H), 3.27 (m, 4H), 3.09 (m, 4H), 2.09 (s, 3H). MS (EI): 514.1 (MH⁺).
- 30 **[0872]** N-{4-[2-({3-[4-(2,3-dihydroxypropyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.46 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.63 (s, 1H), 7.33 (d, 1H), 7.22 (d, 1H), 7.13 (m, 1H), 6.56 (dd, 1H), 3.67 (s, 2H), 3.14 (m, 5H), 2.60 (m, 4H), 2.45 (m, 1H), 2.30 (m, 1H), 2.09 (s, 3H). MS (EI): 463.6 (MH⁺).
- 35 **[0873]** N-{4-[2-({3-[4-(1,3-benzodioxol-5-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO):

5 10.21 (s, 1H), 9.46 (s, 1H), 8.49 (d, 1H), 8.12 (m, 2H), 7.75 (d, 2H), 7.62 (s, 1H), 7.33 (d, 1H), 7.23 (dd, 1H), 7.12 (t, 1H), 6.90-6.85 (m, 2H), 6.79 (m, 1H), 6.55 (dd, 1H), 5.99 (s, 2H), 3.44 (s, 2H), 3.15 (m, 4H), 2.52 (m, 4H), 2.09 (s, 3H). MS (EI): 523.5 (MH+).

[0874] N-{4-[2-({3-[4-(pyridin-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.47 (s, 1H), 8.52-
10 8.49 (m, 2H), 8.13 (m, 2H), 7.81-7.72 (m, 3H), 7.63 (s, 1H), 7.50 (d, 1H), 7.33 (d, 1H), 7.30-7.21 (m, 2H), 7.13 (t, 1H), 6.57 (dd, 1H), 3.67 (s, 2H), 3.17 (m, 4H), 2.60 (m, 4H), 2.09 (s, 3H). MS (EI): 480.6 (MH+).

[0875] N-{4-[2-({3-[4-(pyridin-3-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.47 (s, 1H), 8.54 (d,
15 1H), 8.49 (d, 2H), 8.12 (m, 2H), 7.78-7.73 (m, 3H), 7.63 (s, 1H), 7.40-7.37 (m, 1H), 7.33 (d, 1H), 7.23 (d, 1H), 7.13 (t, 1H), 6.55 (dd, 1H), 3.58 (s, 2H), 3.16 (m, 4H), 2.55 (m, 4H), 2.10 (s, 3H). MS (EI): 480.5 (MH+).

[0876] N-{4-[2-({3-[4-(pyridin-4-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.47 (s, 1H), 8.53
20 (dd, 2H), 8.49 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.64 (s, 1H), 7.38 (dd, 2H), 7.33 (d, 1H), 7.23 (d, 1H), 7.13 (t, 1H), 6.56 (dd, 1H), 3.59 (s, 2H), 3.18 (m, 4H), 2.56 (m, 4H), 2.09 (s, 3H). MS (EI): 480.7 (MH+).

[0877] N-{4-[2-({3-[4-(1H-pyrrol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.45 (s, 1H), 8.49
25 (d, 1H), 8.12 (m, 2H), 7.75 (d, 2H), 7.61 (s, 1H), 7.33 (d, 1H), 7.21 (d, 1H), 7.12 (t, 1H), 6.64 (m, 1H), 6.55 (dd, 1H), 5.92 (m, 2H), 3.46 (s, 2H), 3.14 (m, 4H), 2.51 (m, 4H), 2.09 (s, 3H). MS (EI): 468.6 (MH+).

[0878] 4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-(phenylmethyl)-piperazine-1-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s,
30 1H), 9.40 (s, 1H), 8.62 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.70 (d, 2H), 7.48 (dd, 2H), 7.28-7.21 (m, 3H), 7.00 (d, 2H), 6.94 (t, 1H), 4.41 (s, 2H), 3.60 (m, 4H), 3.10 (m, 4H), 2.09 (s, 3H). MS (EI): 522.4 (MH+).

[0879] N-[4-(2-({3-[4-({2-(methyloxy)phenyl}carbonyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.50 (s,
35 1H), 8.50 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.64 (s, 1H), 7.41 (m, 1H), 7.34 (d, 1H), 7.29 (d, 1H), 7.23 (dd, 1H), 7.16 (t, 1H), 7.10 (d, 1H), 7.01 (t, 1H), 6.59 (dd, 1H), 3.79 (s, 3H), 3.21 (m, 4H), 3.07 (m, 4H), 2.09 (s, 3H). MS (EI): 523.5 (MH+).

- 5 [0880] N-{4-[2-({3-[4-(1H-pyrazol-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 13.22 (s, 1H), 10.22 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 3H), 7.76 (d, 3H), 7.66 (s, 1H), 7.34 (d, 1H), 7.29 (d, 1H), 7.17 (t, 1H), 6.59 (dd, 1H), 3.78 (m, 4H), 3.20 (m, 4H), 2.09 (s, 3H). MS (EI): 483.5 (MH⁺).
- 10 [0881] N-{4-[2-({3-[4-(3-pyridin-3-ylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.24 (s, 1H), 9.50 (s, 1H), 8.50 (d, 2H), 8.39 (dd, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.69 (dd, 2H), 7.34-7.24 (m, 3H), 7.15 (t, 1H), 6.58 (dd, 1H), 3.61 (m, 4H), 3.10 (m, 4H), 2.86 (t, 2H), 2.74 (t, 2H), 2.09 (s, 3H). MS (EI): 522.7 (MH⁺).
- 15 [0882] N-(4-{2-[(3-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.50 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.66 (s, 1H), 7.34 (d, 1H), 7.27 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.63 (m, 4H), 3.58 (t, 2H), 3.23 (s, 3H), 3.12 (m, 4H), 2.63 (t, 2H), 2.09 (s, 3H). MS (EI): 475.6 (MH⁺).
- 20 [0883] N-[4-(2-({3-[4-(2-[(4-fluorophenyl)oxy]acetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl)phenyl]acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.66 (s, 1H), 7.34 (d, 1H), 7.29 (d, 1H), 7.19-7.10 (m, 3H), 6.96 (m, 2H), 6.60 (dd, 1H), 4.88 (s, 2H), 3.64 (m, 4H), 3.17 (m, 4H), 2.09 (s, 3H). MS (EI): 541.5 (MH⁺).
- 25 [0884] N-{4-[2-({3-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.24 (s, 1H), 9.50 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.69 (s, 1H), 7.34 (d, 1H), 7.24 (d, 1H), 7.15 (t, 1H), 6.58 (dd, 1H), 3.61 (m, 2H), 3.48 (m, 2H), 3.41 (t, 1H), 3.10 (m, 4H), 2.18 (m, 2H), 2.09 (s, 3H), 1.92 (m, 2H), 1.75 (m, 2H). MS (EI): 471.4 (MH⁺).
- 30 [0885] N-{4-[2-({3-[4-(pyridin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.22 (s, 1H), 9.51 (s, 1H), 8.69 (dd, 2H), 8.50 (d, 1H), 8.14 (d, 2H), 7.74 (d, 2H), 7.66 (s, 1H), 7.45 (dd, 2H), 7.34 (d, 1H), 7.28 (d, 1H), 7.16 (t, 1H), 6.60 (d, 1H), 3.81 (m, 2H), 3.43 (m, 2H), 3.27 (m, 2H), 3.14 (m, 2H), 2.10 (s, 3H). MS (EI): 494.6 (MH⁺).
- 35 [0886] N-{4-[2-({3-[4-(pyridin-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.50 (s, 1H), 8.61 (d, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.95 (t, 1H), 7.74 (d, 2H), 7.66 (d, 2H), 7.50 (t, 1H), 7.34

5 (d, 1H), 7.29 (d, 1H), 7.16 (t, 1H), 6.60 (d, 1H), 3.84 (m, 2H), 3.59 (m, 2H), 3.26 (m, 2H), 3.14 (m, 2H), 2.09 (s, 3H). MS (EI): 494.6 (MH⁺).

[0887] N-(4-{2-[(3-{4-[(2-methylphenyl)carbonyl]piperazin-1-

yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO):

10.23 (s, 1H), 9.49 (s, 1H), 8.49 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.63 (s, 1H), 7.34-7.16
10 (m, 7H), 6.59 (d, 1H), 3.84 (m, 2H), 3.28 (m, 2H), 3.25 (m, 2H), 3.06 (m, 2H), 2.24 (s, 3H),
2.09 (s, 3H). MS (EI): 507.6 (MH⁺).

[0888] N-{4-[2-[(3-{4-(2,2-dimethylpropanoyl)piperazin-1-

yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO):

10.21 (s, 1H), 9.51 (s, 1H), 8.50 (dd, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.68 (s, 1H), 7.34 (d,
15 1H), 7.26 (d, 1H), 7.16 (t, 1H), 6.59 (d, 1H), 3.72 (m, 4H), 3.13 (m, 4H), 2.09 (s, 3H), 1.23 (s,
9H). MS (EI): 473.5 (MH⁺).

[0889] N-{4-[2-[(3-{4-(pyridin-3-ylcarbonyl)piperazin-1-yl}phenyl)amino]pyrimidin-

4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.51 (s, 1H), 8.67

(m, 2H), 8.50 (d, 1H), 8.13 (m, 2H), 7.90 (m, 1H), 7.75 (d, 2H), 7.66 (s, 1H), 7.50 (m, 1H),
20 7.34 (d, 1H), 7.29 (dd, 1H), 7.17 (t, 1H), 6.60 (dd, 1H), 3.82 (m, 2H), 3.50 (m, 2H), 3.29 (m,
2H), 3.16 (m, 2H), 2.09 (s, 3H). MS (EI): 494.7 (MH⁺).

[0890] N-{4-[2-[(3-{4-(2-methylpropanoyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-

yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d,

1H), 8.14 (dd, 2H), 7.76 (d, 2H), 7.69 (s, 1H), 7.34 (d, 1H), 7.26 (dd, 1H), 7.16 (t, 1H), 6.60
25 (dd, 1H), 3.66 (m, 4H), 3.11 (m, 4H), 2.90 (m, 1H), 2.09 (s, 3H), 1.03 (s, 6H). MS (EI): 459.6
(MH⁺).

[0891] N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-

yl)phenyl]-tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.32 (s,

1H), 9.48 (s, 1H), 8.48 (d, 1H), 8.16 (d, 2H), 7.78 (d, 2H), 7.67 (s, 1H), 7.32 (d, 1H), 7.29
30 (dd, 1H), 6.87 (d, 1H), 3.95 (t, 2H), 3.81 (s, 3H), 3.78 (m, 2H), 3.71 (m, 4H), 3.29-3.17 (m,
1H), 2.91 (m, 4H), 2.13-2.07 (m, 2H). MS (EI): 476.5 (MH⁺).

[0892] (2R)-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)-

phenyl]tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.94 (s, 1H),

9.48 (s, 1H), 8.48 (d, 1H), 8.16 (d, 2H), 7.89 (d, 2H), 7.66 (s, 1H), 7.33 (d, 1H), 7.30 (dd,
35 1H), 6.87 (d, 1H), 4.43 (dd, 1H), 4.01 (m, 1H), 3.86 (m, 1H), 3.81 (s, 3H), 3.72 (m, 4H), 2.91
(m, 4H), 2.22-2.19 (m, 1H), 2.03-1.98 (m, 1H), 1.89 (m, 2H). MS (EI): 476.4 (MH⁺).

- 5 [0893] (2S)-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)-phenyl]tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 9.94 (s, 1H), 9.48 (s, 1H), 8.48 (d, 1H), 8.16 (d, 2H), 7.89 (d, 2H), 7.66 (s, 1H), 7.33 (d, 1H), 7.30 (dd, 1H), 6.87 (d, 1H), 4.43 (dd, 1H), 4.01 (m, 1H), 3.86 (m, 1H), 3.81 (s, 3H), 3.72 (m, 4H), 2.91 (m, 4H), 2.24-2.19 (m, 1H), 2.03-1.98 (m, 1H), 1.89 (m, 2H). MS (EI): 476.5 (MH⁺).
- 10 [0894] N-(4-{2-[(4-{[4-(2-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.85-7.77 (m, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.56-7.46 (m, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 3.18-3.16 (m, 8H), 2.09 (s, 3H). MS (EI): 547.7 (MH⁺).
- 15 [0895] N-(4-{2-[(3-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.73 (m, 3H), 7.64 (s, 1H), 7.54 (d, 2H), 7.34 (d, 1H), 7.29 (d, 1H), 7.16 (t, 1H), 6.60 (dd, 1H), 3.79 (m, 2H), 3.46 (m, 2H), 3.26 (m, 2H), 3.15 (m, 2H), 2.09 (s, 3H). MS (EI): 562.5 (MH⁺).
- 20 [0896] ethyl 3-(4-(2-(4-morpholinophenylamino(pyrimidin-4-yl)phenylamino)-3-oxopropanoate: ¹H NMR (400 MHz, d6-DMSO): 10.46 (s, 1H), 9.35 (s, 1H), 8.42 (d, 1H), 8.11 (d, 2H), 7.71 (d, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.92 (d, 2H), 4.11 (q, 2H), 3.72 (m, 2H), 3.37 (m, 4H), 3.02 (m, 4H), 1.19 (t, 3H). MS (EI): 462 (MH⁺).
- [0897] N-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)penylamino)pyrimidin-4-yl)phenyl)-tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.30 (s, 1H), 9.43 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.97 (m, 2H), 3.96 (t, 1H), 3.76 (m, 2H), 3.63 (m, 4H), 3.19 (m, 2H), 3.06 (m, 4H), 2.93 (m, 1H), 2.10 (m, 2H), 1.02 (d, 6H). MS (EI): 515 (MH⁺).
- 25 [0898] N-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.30 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.96 (d, 2H), 3.96 (t, 1H), 3.76 (m, 3H), 3.59 (m, 2H), 3.41 (m, 3H), 3.19 (m, 1H), 3.03 (m, 4H), 2.41 (m, 6H), 1.90 (m, 1H), 1.75 (m, 1H). MS (EI): 527 (MH⁺).
- [0899] N-ethyl-4-(4-(4-(4-(tetrahydrofuran-3-carboxamido)phenyl)pyrimidin-2-yl-amino(phenyl)piperazine-1-carboxamide: ¹H NMR (400 MHz, d6-DMSO): 10.30 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H),
- 30
- 35

5 6.59 (t, 1H), 3.96 (t, 1H), 3.75 (m, 3H), 3.42 (m, 4H), 3.19 (m, 1H), 3.05 (m, 6H), 2.10 (q, 2H), 1.02 (t, 3H). MS (EI): 516 (MH⁺).

[0900] N-(4-(2-(4-(4-((R)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.97 (d, 10 2H), 3.96 (dd, 1H), 3.89 (m, 1H), 3.76 (m, 4H), 3.63 (m, 4H), 3.18 (m, 2H), 3.07 (m, 4H), 2.09 (m, 2H), 1.13 (d, 3H). MS (EI): 516 (MH⁺).

[0901] N-(4-(2-(4-(4-((S)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.97 (d, 15 2H), 3.96 (t, 1H), 3.85 (q, 1H), 3.76 (m, 3H), 3.63 (m, 4H), 3.19 (m, 1H), 3.07 (m, 4H), 2.10 (m, 2H), 1.11 (d, 3H). MS (EI): 516 (MH⁺).

[0902] N-(4-(2-(4-(4-((R)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.69 20 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.97 (m, 2H), 3.76 (m, 3H), 3.64 (m, 4H), 3.19 (m, 1H), 3.06 (m, 6H), 2.73 (m, 1H), 2.09 (m, 2H), 1.67 (m, 4H). MS (EI): 542 (MH⁺).

[0903] N-(4-(2-(4-(4-((S)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.31 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.77 (d, 2H), 7.69 25 (d, 2H), 7.29 (d, 1H), 6.97 (d, 2H), 3.96 (t, 1H), 3.86 (m, 1H), 3.76 (m, 3H), 3.64 (m, 4H), 3.18 (m, 1H), 3.05 (m, 6H), 2.64 (m, 1H), 2.10 (m, 1H), 2.00 (m, 1H), 1.62 (m, 4H). MS (EI): 542 (MH⁺).

[0904] N-{4-[2-(1H-benzimidazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.25 (s, 1H), 8.97 (s, 1H), 8.78 (s, 30 1H), 8.21(d, 2H), 7.86 (d, 2H), 7.80 (d, 2H), 6.96 (s, 1H), 6.78 (dd, 2H), 2.44 (s, 3H), 2.11 (s, 3H); MS (EI) C₂₀H₁₈N₆O: 359.3 (M+H)⁺.

[0905] 4-(4-furan-2-ylphenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine: ¹H-NMR (400MHz, d₆-DMSO): 9.46 (s, 1H), 8.49 (d, 1H), 8.22 (d, 2H), 7.87 (d, 2H), 7.84 (dd, 1H), 7.68 (d, 2H), 7.37 (d, 1H), 7.12 (t, 1H), 6.94 (d, 2H), 6.66 (dd, 1H), 3.75 (t, 4H), 3.05 (t, 35 4H); MS (EI) C₂₄H₂₂N₄O₂: 399.3 (M+H)⁺.

[0906] N-(4-morpholin-4-ylphenyl)-4-[4-(pyrimidin-2-ylamino)phenyl]pyrimidin-2-amine: ¹H-NMR (400MHz, d₆-DMSO): 10.01 (s, 1H), 9.35 (s, 1H), 8.56 (d, 2H), 8.42 (d,

5 1H), 8.11 (dd, 2H), 7.95 (dd, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 6.96-6.92 (m, 3H), 3.75 (t, 4H), 3.06 (t, 4H); MS (EI) C₂₄H₂₃N₇O: 426.3 (M+H)⁺.

[0907] N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}-5-methylpyrimidin-4-yl)phenyl]-cyclopropanecarboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.40 (s, 1H), 9.41 (bs, 1H), 9.30 (s, 1H), 8.30 (s, 1H), 7.23-7.70 (m, 2H), 7.65-7.62 (m, 3H), 6.91 (d, 2H), 3.70-3.50 (bs, 2H), 3.21-2.87 (m, 8H), 2.20 (s, 3H), 1.80 (p, 1H), 1.18 (bs, 3H), 0.81 (d, 4H); MS (EI) C₂₄H₂₃N₇O: 457.4 (M+H)⁺.

[0908] N-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-cyclopropanecarboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.48 (s, 1H), 9.37 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.76 (d, 2H), 7.65 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.09 (bs, 4H), 2.60-2.35 (m, 6H), 1.83 (p, 1H), 1.06 (t, 3H), 0.84-0.82 (m, 4H); MS (EI) C₂₆H₃₀N₆O: 443.4 (M+H)⁺.

[0909] N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-methylpyrimidin-4-yl}phenyl)-N²,N²-dimethylglycinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.04 (s, 1H), 9.25 (s, 1H), 8.37 (s, 1H), 7.81 (d, 2H), 7.74 (d, 2H), 7.12 (s, 2H), 6.11 (s, 1H), 3.73 (t, 8H), 3.20 (bs, 2H), 3.06 (t, 8H), 2.34 (s, 6H), 2.28 (s, 3H); MS (EI) C₂₉H₃₇N₇O₃: 532.4 (M+H)⁺.

[0910] N²,N²-dimethyl-N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)glycinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.43 (s, 1H), 9.28 (s, 1H), 8.33 (s, 1H), 7.77 (d, 2H), 7.69 (d, 2H), 7.63 (d, 2H), 6.88 (d, 2H), 3.73 (t, 4H), 3.35 (bs, 2H), 3.01 (t, 4H), 2.65 (s, 6H), 2.21 (s, 3H); MS (EI) C₂₅H₃₀N₆O₂: 447.4 (M+H)⁺.

25 [0911] N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.15 (s, 1H), 9.28 (s, 1H), 8.32 (s, 1H), 7.82-7.79 (m, 2H), 7.64 (t, 4H), 6.88 (d, 2H), 3.75-3.72 (t, 5H), 3.01 (t, 4H), 2.91 (t, 2H), 2.22 (s, 3H), 2.11-2.02 (m, 1H), 1.84-1.75 (m, 1H), 1.70-1.63 (m, 2H); MS (EI) C₂₆H₃₀N₆O₂: 459.4 (M+H)⁺.

30 [0912] N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.25 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.83 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 3.80-3.77 (m, 1H), 3.65-3.41 (m, 4H), 3.08-3.02 (m, 4H), 2.96-2.89 (m, 3H), 2.13-2.08 (m, 1H), 1.84-1.78 (m, 1H), 1.73-1.68 (m, 2H), 1.02 (d, 6H); MS (EI) C₂₉H₃₅N₇O₂: 514.4 (M+H)⁺.

35 [0913] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.85 (s, 1H), 9.41 (s, 1H), 8.47 (d, 1H), 8.18 (d, 2H), 7.78 (d, 2H), 7.68 (d, 2H),

5 7.31 (d, 1H), 6.96 (d, 2H), 4.40-4.34 (m, 1H), 3.70 (t, 4H), 3.32-3.25 (m, 2H), 3.05 (t, 4H), 2.44-2.38 (m, 1H), 2.05-1.94 (m, 3H), 1.23 (s, 9H); MS (EI) $C_{30}H_{37}N_7O_2$: 528.4 (M+H)⁺.

[0914] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.19 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 3.74-3.70 (m, 1H), 3.60-3.46 (m, 4H), 3.04-3.00 (m, 4H), 2.91 (t, 2H), 2.22-2.02 (m, 6H), 1.95-1.87 (m, 1H), 1.82-1.73 (m, 2H), 1.70-1.64 (m, 2H); MS (EI) $C_{30}H_{35}N_7O_2$: 526.2 (M+H)⁺.

[0915] N-ethyl-4-[4-({4-[4-(D-prolylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-piperazine-1-carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.19 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.96 (d, 2H), 6.59 (t, 1H), 15 3.74-3.71 (m, 1H), 3.42 (t, 4H), 3.10-3.05 (m, 2H), 3.01 (t, 4H), 2.91 (t, 2H), 2.22 (s, 3H), 2.09-2.02 (m, 1H), 1.84-1.76 (m, 1H), 1.70-1.63 (m, 2H), 1.02 (t, 3H); MS (EI) $C_{28}H_{34}N_8O_2$: 515.5 (M+H)⁺.

[0916] N-[4-(2-({4-(4-D-prolylpiperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 20 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.97 (d, 2H), 3.92-3.89 (m, 1H), 3.75-3.71 (m, 1H), 3.65-3.59 (m, 4H), 3.09-2.98 (m, 5H), 2.91 (t, 2H), 2.69-2.63 (m, 1H), 2.09-2.02 (m, 2H), 1.84-1.76 (m, 1H), 1.70-1.54 (m, 5H); MS (EI) $C_{30}H_{36}N_8O_2$: 541.4 (M+H)⁺.

[0917] N-[4-(2-({4-(4-L-prolylpiperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]-D-prolinamide: ¹H-NMR (400MHz, d₆-DMSO): 10.20 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 25 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.96 (d, 2H), 3.92-3.89 (m, 1H), 3.75-3.71 (m, 1H), 3.65-3.59 (m, 4H), 3.09-2.98 (m, 5H), 2.91 (t, 2H), 2.69-2.63 (m, 1H), 2.09-2.02 (m, 2H), 1.84-1.76 (m, 1H), 1.70-1.56 (m, 5H); MS (EI) $C_{30}H_{36}N_8O_2$: 541.4 (M+H)⁺.

[0918] 1-Methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 9.93 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.87 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.12 (m, 1H), 30 3.05 (m, 4H), 2.95 (m, 1H), 2.36 (m, 4H) 2.17 (m, 1H), 1.80 (m, 3H); MS (EI) for $C_{26}H_{30}N_6O_2$: 459 (MH⁺).

[0919] 1-Methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-2-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 9.97 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.85 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.05 (m, 4H), 2.92 (m, 1H), 2.60 (dd, 1H), 2.16 (s, 3H) 2.03 (m, 1H), 1.76 (m, 2H), 35 1.60 (m, 3H), 1.25 (m, 1H); MS (EI) for $C_{27}H_{32}N_6O_2$: 473 (MH⁺).

- 5 **[0920] N-{4-[2-({4-[4-(Piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.25 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.36 (s, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 3.64 (m, 4H), 3.17 (m, 2H), 3.06 (m, 4H), 2.93 (m, 1H), 2.81 (m, 2H), 2.09 (s, 3H) 1.60-1.75 (m, 4H); MS (EI) for C₂₈H₃₃N₇O₂: 500 (MH⁺).
- 10 **[0921] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-4-ylacetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.53 (s, 1H), 9.38 (s, 1H), 8.53 (d, 2H), 8.44 (d, 1H), 8.12 (d, 2H), 7.95 (d, 1H), 7.76 (d, 2H), 7.67 (d, 2H), 7.35 (d, 1H), 6.92 (d, 2H), 3.75 (m, 6H), 3.04 (m, 4H). MS (EI): 467 (MH⁺).
- [0922] 2-(3-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.44 (s, 1H), 9.36 (s, 1H), 8.42 (m, 1H), 8.09 (d, 2H), 7.74 (d, 2H), 7.64 (d, 2H), 7.35 (m, 1H), 7.24 (m, 1H), 7.15 (d, 2H), 7.07 (m, 1H), 6.90 (d, 2H), 3.71 (m, 6H), 3.04 (m, 4H). MS (EI): 484 (MH⁺).
- 15 **[0923] 3-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)propanamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.19 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.73 (d, 2H), 7.68 (d, 2H), 7.35 (d, 2H), 7.26 (m, 3H), 6.93 (d, 2H), 3.74 (m, 4H), 3.04 (m, 4H), 2.92 (t, 2H), 2.67 (t, 2H). MS (EI): 515 (MH⁺).
- [0924] 2-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.47 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.66 (d, 2H), 7.43 (s, 1H), 7.32 (m, 4H), 6.93 (d, 2H), 3.74 (m, 6H), 3.04 (m, 4H). MS (EI): 500 (MH⁺).
- 25 **[0925] 2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenyl-propanamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.12 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.09 (d, 2H), 7.72 (d, 2H), 7.67 (d, 2H), 7.25 (m, 4H), 7.17 (m, 2H), 6.93 (d, 2H), 3.74 (m, 4H), 3.06 (m, 4H), 2.99 (m, 1H), 2.81 (m, 1H), 2.64 (m, 1H), 1.12 (d, 3H). MS (EI): 494 (MH⁺).
- 30 **[0926] trans-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenyl-cyclopropanecarboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.53 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.77 (d, 2H), 7.67 (d, 2H), 7.28 (m, 3H), 7.21 (m, 3H), 6.93 (d, 2H), 3.74 (m, 4H), 3.04 (m, 4H), 2.39 (m, 1H), 2.11 (m, 1H), 1.53 (m, 1H), 1.42 (m, 1H). MS (EI): 492 (MH⁺).
- 35 **[0927] 2-(4-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.44 (s, 1H), 9.38 (s, 1H), 8.43 (d,

- 5 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.65 (d, 2H), 7.38 (m, 2H), 7.27 (d, 1H), 7.18 (dd, 2H), 6.93 (d, 2H), 3.71 (m, 4H), 3.69 (s, 2H), 3.04 (m, 4H). MS (EI): 484 (MH+).
- [0928] 3-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)propanamide:** ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.45 (dd, 1H), 7.40 (dd, 1H), 7.25 (m, 3H), 6.93 (d, 2H), 3.74 (m, 4H), 3.04 (m, 6H), 2.70 (t, 2H). MS (EI): 515 (MH+).
- 10 **[0929] 3-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)propanamide:** ¹H NMR (400 MHz, d6-DMSO): 10.19 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.27 (m, 5H), 6.93 (d, 2H), 3.74 (m, 4H), 3.04 (m, 4H), 2.94 (t, 2H), 2.69 (t, 2H). MS (EI): 515 (MH+).
- 15 **[0930] 3-(2-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)propanamide:** ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.73 (d, 2H), 7.67 (d, 2H), 7.35 (t, 1H), 7.26 (d, 2H), 7.11 (m, 2H), 6.93 (d, 2H), 3.74 (m, 4H), 3.04 (m, 4H), 2.95 (t, 2H), 2.68 (t, 2H). MS (EI): 498 (MH+).
- [0931] Nalpha,Nalpha-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-phenylalaninamide:** ¹H NMR (400 MHz, d6-DMSO): 10.04 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.09 (d, 2H), 7.74 (d, 2H), 7.67 (d, 2H), 7.24 (m, 5H), 7.17 (m, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.48 (dd, 1H), 3.06 (m, 5H), 2.86 (dd, 1H), 2.49 (s, 6H). MS (EI): 523 (MH+).
- 20 **[0932] 2-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.54 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.76 (d, 2H), 7.67 (d, 2H), 7.46 (m, 2H), 7.33 (m, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 3.89 (s, 2H), 3.74 (m, 4H), 3.04 (m, 4H). MS (EI): 500 (MH+).
- 25 **[0933] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-2-yl-acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.51 (s, 1H), 9.35 (s, 1H), 8.49 (d, 1H), 8.41 (d, 1H), 8.10 (d, 2H), 7.77 (m, 3H), 7.64 (d, 2H), 7.39 (d, 1H), 7.26 (m, 2H), 6.92 (d, 2H), 3.87 (s, 2H), 3.71 (m, 4H), 3.02 (m, 4H). MS (EI): 467 (MH+).
- 30 **[0934] 2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.47 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.66 (d, 2H), 7.39 (m, 4H), 7.26 (m, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.70 (s, 2H), 3.04 (m, 4H). MS (EI): 500 (MH+).
- 35 **[0935] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-{4-[(trifluoro-methyl)oxy]phenyl}acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.50 (s, 1H),

5 9.38 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.75 (d, 2H), 7.66 (d, 2H), 7.47 (d, 2H), 7.35 (d, 2H), 7.27(d, 1H), 6.93 (d, 2H), 3.74 (m, 6H), 3.04 (m, 4H). MS (EI): 550 (MH+).

[0936] 2-[2-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.34 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.76 (d, 2H), 7.67 (d, 2H), 7.26 (m, 3H), 7.00 (d, 1H), 6.91 (m, 3H),
10 3.77 (s, 3H), 3.74 (m, 4H), 3.67 (s, 2H), 3.04 (m, 4H). MS (EI): 496 (MH+).

[0937] 2-[3-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.44 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.76 (d, 2H), 7.67 (d, 2H), 7.26 (m, 2H), 6.93 (m, 4H), 6.82 (dd, 1H), 3.74 (m, 7H), 3.55 (s, 2H), 3.04 (m, 4H). MS (EI): 496 (MH+).

15 [0938] 2-[4-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.38 (s, 1H), 9.38 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.26 (m, 3H), 6.93 (m, 4H), 3.74 (m, 7H), 3.60 (s, 2H), 3.04 (m, 4H). MS (EI): 496 (MH+).

[0939] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-alaninamide: ¹H NMR (400 MHz, d6-DMSO): 9.35 (s, 1H), 8.43 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.63 (d, 2H), 7.28 (d, 1H), 6.92 (d, 2H), 3.48 (m, 1H), 2.35 (q, 2H), 1.86 (br s, 8H), 1.24 (d, 3H), 1.03 (t, 3H). MS (EI): 446 (MH+).

[0940] N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H NMR (400 MHz, d6-DMSO): 10.21 (s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.68 (m, 4H), 3.10 (s, 2H), 3.07 (m, 4H), 2.18 (s, 6H), 2.09 (s, 3H). MS (EI): 474 (MH+).

[0941] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-3-(methyloxy)propanamide: ¹H NMR (400 MHz, d6-DMSO): 10.23 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.65 (d, 2H), 7.25 (d, 1H), 6.92 (d, 2H), 3.63 (t, 2H),
30 3.25 (s, 3H), 3.09 (m, 4H), 2.60 (t, 2H), 2.58 (m, 6H), 1.05 (t, 3H). MS (EI): 461 (MH+).

[0942] (2R)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide: ¹H NMR (400 MHz, d6-DMSO): 9.37 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.80 (d, 2H), 7.66 (d, 2H), 7.49 (d, 2H), 7.34 (t, 2H), 7.26 (m, 2H), 6.92 (d, 2H), 4.56 (s, 1H), 3.75 (m, 4H), 3.04 (m, 4H). MS (EI): 481 (MH+).

35 [0943] N'2',N'2'-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)-D-alaninamide: ¹H NMR (400 MHz, d6-DMSO): 10.02 (s, 1H), 9.38 (s, 1H), 8.44

- 5 (d, 1H), 8.11 (d, 2H), 7.84 (d, 2H), 7.66 (d, 2H), 7.28 (d, 1H), 6.92 (d, 2H), 3.75 (m, 4H), 3.21 (q, 1H), 3.04 (m, 4H), 2.25 (s, 6H), 1.19 (d, 3H). MS (EI): 447 (MH⁺).
- [0944] 1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide:** ¹H NMR (400 MHz, d₆-DMSO): 9.94 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.87 (d, 2H), 7.67 (d, 2H), 7.30 (d, 1H), 6.94 (d, 2H), 3.76 (m, 4H), 3.12 (m, 1H),
- 10 3.05 (m, 4H), 2.95 (m, 1H), 2.36 (s, 3H), 2.30 (m, 1H), 2.18 (m, 1H), 1.78 (m, 3H). MS (EI): 459 (MH⁺).
- [0945] N'2',N'2'-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)-L-alaninamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.03 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.28 (d, 1H), 6.92 (d, 2H), 3.75 (m, 4H),
- 15 3.21 (q, 1H), 3.05 (m, 4H), 2.25 (s, 6H), 1.19 (d, 3H). MS (EI): 447 (MH⁺).
- [0946] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1-phenyl-cyclopropanecarboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 9.38 (s, 1H), 8.43 (d, 1H), 8.09 (d, 2H), 7.72 (d, 2H), 7.65 (d, 2H), 7.40 (m, 4H), 7.28 (m, 2H), 6.92 (d, 2H), 3.74 (m, 4H), 3.04 (m, 4H), 1.74 (dd, 2H), 1.15 (dd, 2H). MS (EI): 492 (MH⁺).
- 20 **[0947] 2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-butanamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.12 (s, 1H), 9.38 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.78 (d, 2H), 7.67 (d, 2H), 7.27 (d, 1H), 6.93 (d, 2H), 3.75 (m, 4H), 3.04 (m, 4H), 2.46 (q, 1H), 1.65 (m, 1H), 1.41 (m, 1H), 1.10 (d, 3H), 0.87 (t, 3H). MS (EI): 432 (MH⁺).
- [0948] (2S)-1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-azetidine-2-carboxamide:** ¹H NMR (400 MHz, d₆-DMSO): 9.88 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.89 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.93 (d, 2H), 3.74 (m, 4H), 3.56 (t, 1H), 3.36 (m, 1H), 3.04 (m, 4H), 2.93 (q, 1H), 2.33 (s, 3H), 2.30 (m, 1H), 2.12 (m, 1H). MS (EI): 445 (MH⁺).
- 25 **[0949] 2,4,6-trichloro-N-(3-{[4-(4-methyl-2-thienyl)pyrimidine-2-yl]amino}propyl)-benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 8.68 (br s, 1H), 8.24 (d, 1H), 7.72-7.70 (m, 3H), 7.29 (s, 1H), 7.17 (t, 1H), 6.98 (d, 1H), 3.37-3.35 (m, 2H), 3.28-3.27 (m, 2H), 2.22 (s, 3H), 1.77 (br t, 2H). MS (EI): 457.0 (MH⁺).
- 30 **[0950] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide:** MS (EI) C₂₉H₃₄N₆O₂: 499 (MH⁺)
- 35 **[0951] 4-{4-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide:** MS (EI) C₂₇H₃₁N₇O₂: 486 (MH⁺)

- 5 **[0952] N-[3-({4-[3,4-bis(methoxy)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichloro-benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.67 (br s, 1H), 8.26 (d, 1H), 7.69-7.67 (m, 2H), 7.49-7.35 (m, 3H), 7.14-7.09 (m, 2H), 7.03 (d, 1H), 3.82 (s, 3H), 3.80 (s, 3H), 3.42 (m, 2H), 3.32 (m, 2H), 1.80 (m, 2H). MS (EI): 461.2 (MH+).
- [0953] 2,6-dichloro-N-[3-({4-[(4-morpholino-4-ylphenyl)amino]pyrimidin-2-yl}amino)-propyl]benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.85 (br s, 1H), 8.63 (t, 1H), 7.69 (d, 1H), 7.48 (d, 2H), 7.44 (d, 1H), 7.42 (s, 1H), 7.37-7.33 (m, 1H), 6.81 (d, 2H), 6.59 (br s, 1H), 5.83 (d, 1H), 3.67-3.65 (m, 4H), 3.23-3.20 (m, 4H), 2.97-2.94 (m, 4H), 1.70 (t, 2H). MS (EI): 501.2 (MH+).
- 10 **[0954] 2,6-dichloro-N-[3-({4-(2,3-dihydro-1,4-benzodioxin-6-yl)-5-fluoropyrimidin-2-yl}amino)propyl]benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.62 (t, 1H), 8.29 (d, 1H), 7.49-7.42 (m, 4H), 7.37-7.33 (m, 1H), 7.15 (t, 1H), 6.94-6.92 (m, 1H), 4.27-4.21 (m, 4H), 3.33-3.22 (m, 4H), 1.74 (t, 2H); MS (EI): 477.1 (MH+).
- [0955] 2,6-dichloro-N-[3-({4-[3-[(dimethylamino)methyl]phenyl]pyrimidin-2-yl}amino)propyl]benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.68 (t, 1H), 8.31 (d, 1H), 8.01-7.95 (m, 2H), 7.49-7.38 (m, 5H), 7.23 (br s, 1H), 7.10 (d, 1H), 3.43 (m, 2H), 3.33-3.29 (m, 4H), 2.14 (s, 6H), 1.82 (t, 2H); MS (EI): 460.2 (MH+).
- 20 **[0956] 2,6-dichloro-N-[3-({4-[3-(1-methylethyl)phenyl]pyrimidin-2-yl}amino)-propyl]-benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.66 (t, 1H), 8.31 (d, 1H), 7.95-7.88 (m, 2H), 7.55 (br s, 1H), 7.45-7.43 (m, 2H), 7.40-7.34 (m, 3H), 7.20 (br s, 1H), 3.30 (br s, 2H), 3.29-3.25 (m, 2H), 2.92 (septet, 1H), 1.78 (m, 2H), 1.18 (d, 6H); MS (EI): 443.0 (MH+).
- 25 **[0957] 2,6-dichloro-N-[3-({4-[4-[(1-methylethyl)oxy]phenyl]pyrimidin-2-yl}amino)propyl]-benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.68 (t, 1H), 8.25 (d, 1H), 8.03 (d, 2H), 7.49-7.47 (m, 2H), 7.42-7.38 (m, 1H), 7.10 (t, 1H), 7.03 (d, 1H), 6.98 (d, 2H), 4.69 (septet, 1H), 3.42 (m, 2H), 3.30 (m, 2H), 1.80 (t, 2H), 1.27 (d, 6H); MS (EI): 459.0 (MH+).
- 30 **[0958] N-[3-({4-[3-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichloro-benzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.1 (s, 1H), 8.70 (t, 1H), 8.33-8.27 (m, 2H), 7.70 (m, 2H), 7.49-7.46 (m, 2H), 7.42-7.37 (m, 2H), 7.37 (br s, 1H), 7.01 (d, 1H), 3.43 (m, 4H), 2.02 (s, 3H), 1.97 (m, 2H). MS (EI): 458.2 (MH+).
- 35 **[0959] 2,6-dichloro-N-[3-({4-[(E)-2-phenylethenyl]pyrimidin-2-yl}amino)propyl]-benzamide:** ¹H-NMR (400MHz, d6-DMSO): 8.71 (t, 1H), 8.27 (d, 1H), 7.76 (d, 1H), 7.67-

- 5 7.65 (m, 2H), 7.51-7.49 (m, 2H), 7.44-7.34 (m, 4H), 7.12-7.06 (m, 2H), 6.72 (d, 1H), 3.36 (m, 2H), 3.33 (m, 2H), 1.81 (t, 2H). MS (EI): 427.0 (MH⁺).
- [0960] phenyl (4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)carbamate:** ¹H-NMR (400MHz, d6-DMSO): 8.59 (d, 1H), 7.90 (d, 2H), 7.69 (d, 1H), 7.44-7.40 (m, 2H), 7.28-7.20 (m, 5H), 6.97 (d, 2H), 6.62 (d, 2H), 5.90 (s, 2H), 3.74-3.72 (m, 4H), 3.13-3.11 (m, 4H). MS (EI): 468.1 (MH⁺).
- 10 **[0961] phenylmethyl (4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-carbamate:** ¹H-NMR (400MHz, d6-DMSO): 8.55 (d, 1H), 7.85 (d, 2H), 7.64 (d, 1H), 7.33-7.30 (m, 5H), 7.13 (d, 2H), 6.92 (d, 2H), 6.62 (d, 2H), 5.88 (s, 2H), 5.88 (s, 2H), 3.74-3.71 (m, 4H), 3.11-3.09 (m, 4H). MS (EI): 428.3 (MH⁺).
- 15 **[0962] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.2 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.95 (d, 2H), 3.71-3.69 (m, 4H), 3.65 (t, 2H), 3.25 (s, 3H), 3.06-3.03 (m, 4H), 2.59 (t, 2H), 1.23 (s, 9H). MS (EI): 517.4 (MH⁺).
- 20 **[0963] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methyloxy)propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.2 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.65-3.60 (m, 4H), 3.47-3.37 (m, 4H), 3.25 (s, 3H), 3.03-3.02 (m, 3H), 2.60 (t, 2H), 2.21-2.07 (m, 4H), 1.94-1.87 (m, 1H), 1.78-1.73 (m, 1H). MS (EI): 515.2 (MH⁺).
- 25 **[0964] 3-(methyloxy)-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)-pyrimidin-4-yl]phenyl}propanamide:** ¹H-NMR (400MHz, d6-DMSO): 10.2 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.76 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.96 (d, 2H), 3.65-3.62 (m, 6H), 3.25 (s, 3H), 3.08-3.02 (m, 4H), 2.92 (m, 1H), 2.59 (t, 2H), 1.02 (d, 6H). MS (EI): 503.4 (MH⁺).
- 30 **[0965] N-ethyl-4-(4-{[4-(4-{[3-(methyloxy)propanoyl]amino}phenyl)pyrimidin-2-yl]-amino}phenyl)piperazine-1-carboxamide:** ¹H-NMR (400MHz, d6-DMSO): 10.2 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 6.59 (t, 1H), 3.64 (t, 2H), 3.43 (m, 4H), 3.25 (s, 3H), 3.10-3.03 (m, 6H), 2.61 (t, 2H), 1.02 (t, 3H). MS (EI): 504.4 (MH⁺).
- 35 **[0966] N-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-2-phenyl-acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.45 (s, 1H), 9.36 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.76 (d, 2H), 7.64 (d, 2H), 7.38-7.33 (m, 3H), 7.27 (d, 1H), 6.92 (d, 2H),

- 5 3.69 (s, 2H), 3.10-3.04 (m, 4H), 2.35 (q, 3 H), 1.89 (s, 2H), 1.03 (t, 2H); MS (EI): 493.1 (MH⁺).
- [0967] 1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidin-2-one: ¹H-NMR (400MHz, d6-DMSO): 8.26 (d, 1H), 8.14 (d, 2H), 7.77 (d, 2H), 7.65 (d, 2H), 7.36 (d, 1H), 7.25 (d, 2H), 3.92-3.84 (m, 5H), 3.82-3.74 (m, 1H), 3.74-3.60 (m, 1H), 3.42-3.30 (m, 10 4H), 3.06-3.02 (m, 1H), 2.16-2.06 (m, 2H); MS (EI): 416.1 (MH⁺).
- [0968] (R)-2-amino-N-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.63-3.56 (m, 2H), 3.43-3.37 (m, 3H), 3.18 (d, 1H), 3.07-2.98 (m, 4H), 2.25-2.02 (m, 4H), 1.98-1.83 (m, 15 1H), 1.82-1.70 (m, 1H), 1.23 (d, 3H); MS (EI): 500.2 (MH⁺).
- [0969] (R)-2-amino-N-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide: ¹H-NMR (400MHz, d6-DMSO): 9.41 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.73-3.67 (m, 4H), 3.52-4.42 (m, 1H), 3.08-3.02 (m, 4H), 1.25 (s, 3H), 1.23 (d, 3H); MS (EI): 502.4 (MH⁺).
- 20 [0970] (S)-2-hydroxy-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)-phenyl)butanamide: ¹H-NMR (400MHz, d6-DMSO): 9.90 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (d, 1H), 3.86 (dd, 1H), 3.78-3.73 (m, 4H), 3.08-3.02 (m, 4H), 0.96 (d, 3H), 0.87 (d, 3H); MS (EI): 448.3 (MH⁺).
- 25 [0971] (R)-2-hydroxy-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)-phenyl)butanamide: ¹H-NMR (400MHz, d6-DMSO): 9.90 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.94 (d, 2H), 5.76 (d, 1H), 3.86 (dd, 1H), 3.78-3.73 (m, 4H), 3.08-3.02 (m, 4H), 0.96 (d, 3H), 0.87 (d, 3H); MS (EI): 448.3 (MH⁺).
- 30 [0972] N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}-D-alaninamide: ¹H NMR (400 MHz, d6-DMSO): 11.17 (s, 1H), 10.04 (s, 1H), 8.58 (s, 1H), 8.40 (s, 2H), 8.11-8.09 (m, 2H), 7.96-7.82 (m, 3H), 7.76-7.65 (m, 1H), 7.45 (d, 1H), 4.05 (t, 4H), 3.75-3.70 (m, 1H), 3.50 (t, 4H), 2.10-2.00 (m, 1H), 1.45 (d, 3H), 0.82-0.72 (m, 4H). MS (EI): 486 (MH⁺).
- 35 [0973] (2S)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide: ¹H NMR (400 MHz, d6-DMSO): 11.73 (s, 1H), 10.08 (s,

5 1H), 8.99 (s, br, 3H), 8.58 (s, 1H), 8.19 (d, 2H), 7.96-7.83 (m, 3H), 7.76-7.65 (m, 3H), 7.45-7.40 (m, 3H), 5.40 (s, br, 1H), 3.85 (s, br, 4H), 3.50 (s, br, 4H). MS (EI): 481 (MH⁺).

[0974] 2-amino-2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 11.80 (s, 1H), 10.00 (s, 1H), 9.00 (s, 2H), 8.57 (d, 1H), 8.20 (d, 2H), 7.95-7.83 (m, 4H), 7.80-7.60 (m, 3H), 7.58 (d, 2H), 7.43 (d, 1H), 5.50 (s, 1H), 4.00 (t, 4H), 3.50 (t, 4H). MS (EI): 515 (MH⁺).

[0975] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)morpholine-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 11.60 (s, 1H), 10.20 (s, 1H), 10.00 (s, 1H), 9.40 (s, br, 1H), 8.58 (d, 1H), 8.20 (d, 2H), 7.95-7.88 (m, 3H), 7.60-7.20 (m, 4H), 4.42-4.30 (m, 2H), 4.05-3.90 (m, 2H), 3.85-3.70 (m, 4H), 3.60-3.45 (m, 4H), 3.25-3.10 (m, 3H). MS (EI): 461 (MH⁺).

[0976] 1-ethyl-3-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]urea: ¹H NMR (400 MHz, d₆-DMSO): 9.40 (s, 1H), 8.90 (s, 1H), 8.42 (d, 1H), 8.20 (s, 1H), 8.05 (d, 2H), 7.56 (d, 2H), 7.28 (d, 2H), 6.83 (d, 1H), 6.36 (t, 1H), 3.80 (s, 3H), 3.12 (q, 2H), 2.98 (s, br, 4H), 2.58 (s, br, 4H), 2.42 (q, 2H), 1.08-1.00 (m, 6H). MS (EI): 476 (MH⁺).

[0977] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)-phenyl]-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 11.23 (s, 1H), 11.05 (s, 1H), 10.18 (s, br, 1H), 10.00 (s, 1H), 8.75 (s, br, 1H), 8.57 (d, 1H), 8.21 (d, 2H), 7.85 (d, 2H), 7.63 (s, 1H), 7.44 (d, 1H), 7.33 (dd, 1H), 7.03 (d, 1H), 4.55-4.50 (m, 1H), 3.82 (s, 3H), 3.80-3.60 (m, 4H), 3.35-3.05 (m, 7H), 2.50-2.45 (m, 2H), 2.02-1.95 (m, 3H), 1.30 (t, 3H). MS (EI): 502 (MH⁺).

[0978] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)-phenyl]acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (s, 1H), 8.15 (d, 2H), 7.80-7.60 (m, 3H), 7.33 (d, 2H), 6.85 (d, 1H), 3.80 (s, 3H), 2.90 (s, br, 4H), 2.35 (q, 2H), 2.05 (s, 4H), 1.95 (s, 3H), 1.00 (s, 3H). MS (EI): 447 (MH⁺).

[0979] 1-(2,6-dichlorophenyl)-3-(3-{[4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino}propyl)-urea: ¹H NMR (400 MHz, d₆-DMSO): 8.26 (d, 1H), 8.02 (br, 1H), 7.71 (s, 1H), 7.48 (d, 2H), 7.31 (s, 1H), 7.26 (t, 1H), 7.16 (t, 1H), 6.99 (d, 1H), 6.39 (t, 1H), 3.38 (t, 2H), 3.15 (t, 2H), 2.21 (s, 3H), 1.65 (m, 2H). MS (EI) for C₁₉H₁₉Cl₂N₅OS : 436 (MH⁺)

[0980] 1-[2-fluoro-5-(trifluoromethyl)phenyl]-3-(3-{[4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino}propyl)urea: ¹H NMR (400 MHz, d₆-DMSO): 8.65 (d, 2H), 8.23 (s, 1H), 7.7 (s,

- 5 1H), 7.4 (t, 1H), 7.38-7.15 (m, 3H), 7.0 (s, 1H), 6.8 (t, 1H), 3.38 (t, 2H), 3.2 (t, 2H), 2.21 (s, 3H), 1.75 (m, 2H). MS (EI) for $C_{20}H_{19}F_4N_5OS$: 454 (MH^+)
- [0981] 2,6-dichloro-N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-benzenesulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.16 (d, 1H), 8.12 (t, 1H), 7.94 (d, 2H), 7.58 (d, 2H), 7.48 (t, 1H), 6.97 (d, 1H), 6.92 (t, 6.76 (d, 2H) 3.28 (m, 2H), 3.022.96 (m, 8H), 1.68 (m, 2H). MS (EI) for $C_{21}H_{23}Cl_2N_5O_2S$: 480 (MH^+)
- 10 [0982] N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-difluoro-benzenesulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.25 (T, 1H), 8.16 (d, 1H), 7.94 (d, 2H), 7.66 (m, 1H), 7.24 (t, 2H), 6.98 (d, 1H), 6.95 (t, 1H), 6.76 (d, 2H), 3.33 (t, 2H), 3.0 (t, 2H), 2.98 (s, 6H), 1.68 (m, 2H). MS (EI) for $C_{21}H_{23}F_2N_5O_2S$: 448 (MH^+)
- 15 [0983] N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]naphthalene-2-sulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.42 (br, 1H), 8.15-8.06 (m, 3H), 8.02 (d, 1H), 7.94 (d, 2H), 7.8 (dd, 1H), 7.74-7.62 (m, 3H), 6.96 (d, 1H), 6.92 (t, 1H), 6.74 (d, 2H), 3.3 (t, 2H), 2.98 (s, 6H), 2.83 (t, 2H), 1.63 (m, 2H). MS (EI) for $C_{25}H_{27}N_5O_2S$: 462 (MH^+)
- 20 [0984] N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]-3,4-bis(methyloxy)benzenesulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.17 (d, 1H), 7.94 (d, 2H), 7.46 (t, 1H), 7.34 (dd, 1H), 7.27 (d, 1H), 7.06 (d, 1H), 6.97 (d, 1H), 6.93 (t, 1H), 6.76 (d, 2H), 3.8 (s, 6H), 3.3 (t, 2H), 2.98 (s, 6H), 2.8 (t, 2H), 1.65 (m, 2H). MS (EI) for $C_{23}H_{29}N_5O_4S$: 472 (MH^+)
- 25 [0985] 3-chloro-N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]propane-1-sulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.2 (s, 1H), 7.98 (d, 2H), 7.2 (t, 1H), 7.0 (t, 2H), 6.8-6.7 (m, 2H), 3.7 (t, 2H), 3.1-2.9 (m, 10H), 2.05 (t, 2H), 1.7 (m, 2H), 1.2 (m, 2H). MS (EI) for $C_{18}H_{26}ClN_5O_2S$: 412 (MH^+)
- [0986] N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]propane-1-sulfonamide: 1H NMR (400 MHz, d_6 -DMSO): 8.2 (d, 1H), 7.96 (d, 2H), 7.0-6.95 (m, 3H), 6.76 (d, 2H), 3.38 (t, 2H), 3.0-2.9 (m, 10H), 1.75 (t, 2H), 1.6 (q, 2H), 0.95 (t, 3H). MS (EI) for $C_{18}H_{27}N_5O_2S$: 378 (MH^+).
- 30 [0987] methyl (3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)propyl)carbamate: 1H MR (400 MHz, d_6 -DMSO): 8.4 (d, 1H), 7.75 (s, 1H), 7.63-7.55 (m, 2H), 7.35 (t, 1H), 7.12 (t, 1H), 6.8 (d, 1H), 3.5 (s, 3H), 3.28 (t, 2H), 3.03 (t, 2H), 1.65 (m, 2H). MS (EI) for $C_{15}H_{16}Cl_2N_4O_2$: 355 (MH^+).
- 35

- 5 **[0988] 1-methylethyl (3-{{4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino}propyl)-
arbamate:** ¹H NMR (400 MHz, d₆-DMSO): 8.38 (d, 1H), 7.75 (s, 1H), 7.63-7.55 (m,
2H), 7.35 (t, 1H), 7.0 (t, 1H), 6.8 (d, 1H), 4.72 (m, 1H), 3.28 (q, 2H), 3.0 (q, 2H), 1.65 (p, 2H),
1.12 (d, 6H). MS (EI) for C₁₇H₂₀Cl₂N₄O₂: 383 (MH⁺).
- [0989] phenylmethyl (3-{{4-(2,4-dichlorophenyl)pyrimidin-2-
10 yl}amino}propyl)carbamate:** ¹H NMR (400 MHz, d₆-DMSO): 8.46 (d, 1H), 8.2 (br, 1H),
7.8 (d, 1H), 7.66 (br, 1H), 7.6 (dd, 1H), 7.4-7.28 (m, 5H), 7.04 (br, 1H), 5.0 (s, 2H), 3.4 (t,
2H), 3.1 (t, 2H), 1.7 (m, 2H). MS (EI) for C₂₁H₂₀Cl₂N₄O₂: 431 (MH⁺).
- [0990] N-{4-[2-({[3-(3-chlorophenyl)isoxazol-5-yl]methyl}amino)pyrimidin-4-yl]-
15 henyl}acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 8.37 (d, 1H), 8.06 (d, 2H),
7.92 (t, 1H), 7.88-7.82 (m, 2H), 7.7 (d, 2H), 7.56-7.48 (m, 2H), 7.2 (d, 1H), 7.0 (s, 1H), 4.7
(s, 2H), 2.05 (s, 3H). MS (EI) for C₂₂H₁₈ClN₅O₂: 420 (MH⁺).
- [0991] ethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)piperidine-1-
20 carboxylate:** ¹H MR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 8.3 (d, 1H), 8.04 (d, 2H), 7.7 (d,
2H), 7.13 (d, 1H), 7.06 (d, 1H), 4.05 (q, 3H), 3.95 (br, 2H), 2.96 (br, 2H), 2.08 (s, 3H), 1.9
(br, 2H), 1.4 (q, 2H), 1.2 (t, 3H). MS (EI) for C₂₀H₂₅N₅O₃: 384 (MH⁺).
- [0992] 1,1-dimethylethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-
yl}amino)piperidine-1-carboxylate:** ¹H NMR (400 MHz, d₆-DMSO): 10.18 (s, 1H), 8.3 (d,
1H), 8.05 (d, 2H), 7.7 (d, 2H), 7.2 (br, 1H), 7.1 (d, 1H), 3.92 (br, 3H), 2.9 (br, 2H), 2.08 (s,
3H), 1.87 (br, 2H), 1.46-1.36 (m, 11H). MS (EI) for C₂₂H₂₉N₅O₃: 412 (MH⁺).
- 25 **[0993] N-{4-[2-(3,5-diamino-1H-1,2,4-triazol-1-yl)pyrimidin-4-yl]phenyl}acetamide:**
¹H MR (400 MHz, d₆-DMSO): 10.28 (s, 1H), 8.7 (d, 1H), 8.16 (d, 2H), 7.78 (d, 2H), 7.7 (d,
1H), 7.58 (s, 2H), 2.03 (s, 3H). MS (EI) for C₁₄H₁₄N₈O: 311 (MH⁺).
- [0994] N-{4-[2-({5-[(4-ethylpiperazin-1-yl)carbonyl]pyridin-2-yl}amino)pyrimidin-4-
30 yl]-henyl}acetamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.26 (s, 1H), 10.12 (s, 1H), 8.6 (d,
1H), 8.46 (d, 1H), 8.36 (d, 1H), 8.18 (d, 2H), 7.9 (dd, 1H), 7.77 (d, 2H), 7.54 (d, 1H), 2.46-
2.32 (m, 6H), 2.1 (s, 3H), 1.0 (t, 3H). MS (EI) for C₂₄H₂₇N₇O₂: 446 (MH⁺).
- [0995] N-(4-{2-[(4-cyanophenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H MR (400
MHz, d₆-DMSO): 10.24 (d, 2H), 8.6 (d, 1H), 8.17 (d, 2H), 8.06 (d, 2H), 7.78 (d, 4H), 7.5 (d,
1H), 2.05 (s, 3H). MS (EI) for C₁₉H₁₅N₅O: 330 (MH⁺).
- 35 **[0996] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyridin-4-yl}phenyl)acetamide:**
¹H MR (400 MHz, d₆-DMSO): 10.14 (s, 1H), 8.82 (s, 1H), 8.12 (d, 1H), 7.72 (d, 2H), 7.62

5 (d, 2H), 7.53 (d, 2H), 6.97-6.92 (m, 2H), 6.9 (d, 2H), 3.74 (t, 4H), 3.02 (t, 4H), 2.07 (s, 3H). MS (EI) for $C_{23}H_{24}N_4O_2$: 389 (MH^+).

[0997] **N-[4-(2-([4-(4-ethylpiperazin-1-yl)phenyl]amino)-5-methylpyrimidin-4-yl)phenyl]-3-(methyloxy)propanamide:** 1H NMR (400 MHz, d_6 -DMSO): 10.18 (s, 1H), 9.25 (s, 1H), 8.3 (s, 1H), 7.75 (d, 2H), 7.63 (d, 2H), 7.61 (d, 2H), 6.86 (d, 2H), 3.64 (t, 2H), 10 3.25 (s, 3H), 3.03 (t, 4H), 2.6 (t, 2H), 2.38 (br, 2H), 2.2 (s, 3H), 1.03 (t, 3H). MS (EI) for $C_{22}H_{34}N_6O_2$: 475 (MH^+).

[0998] **tert-butyl 1-(4-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)piperidin-4-ylcarbamate:** 1H NMR (400 MHz, d_6 -DMSO): 10.21 (s, 1H), 9.38 (s, br, 1H), 8.43 (s, 1H), 8.08 (m, 2H), 7.74 (m, 2H), 7.62 (m, 2H), 7.23 (m, 1H), 6.98 (m, 3H), 3.77 (m, 2H), 3.63 (m, 15 2H), 2.09 (s, 3H), 1.80 (m, 2H), 1.49 (m, 2H), 1.30 (s, 9H). MS (EI) for $C_{28}H_{34}N_6O_3$: 503 (MH^+).

[0999] **4-(4-aminophenyl)-N-(4-(4-aminopiperidin-1-yl)phenyl)pyrimidin-2-amine:** 1H MR (400 MHz, d_6 -DMSO): 9.19 (s, 1H), 8.3 (m, 1H), 7.87 (m, 2H), 7.63 (m, 2H), 7.14 (m, 1H), 6.92 (m, 2H), 6.62 (m, 2H), 5.74 (m, 2H), 3.57 (m, 2H), 2.67 (m, 2H), 1.81 (m, 2H), 20 1.38 (m, 2H). MS (EI) for $C_{21}H_{24}N_6$: 361 (MH^+).

[01000] **N-(1-(4-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)piperidin-4-yl)-acetamide:** 1H NMR (400 MHz, d_6 -DMSO): 10.26 (s, 1H), 9.34 (s, 1H), 8.43 (d, 1H), 8.1 (d, 2H), 7.85 (d, 1H), 7.75 (d, 2H), 7.64 (d, 2H), 7.27 (d, 1H), 6.94 (d, 2H), 3.67 (m, 1H), 3.55 (m, 2H), 2.72 (t, 2H), 2.09 (s, 3H), 1.88-1.76 (m, 5H), 1.48 (m, 2H). MS (EI) for $C_{25}H_{28}N_6O_2$: 25 445 (MH^+).

[01001] **N-(4-(2-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)-phenyl)cyclopropanecarboxamide:** 1H NMR (400 MHz, d_6 -DMSO): 10.51 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.87 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 3.82 (m, 2H), 3.61 (m, 2H), 3.25-2.99 (m, 4H), 2.04 (m, 1H), 1.83 (m, 1H), 0.89-0.68 (m, 30 8H).. MS (EI) for $C_{28}H_{30}N_6O_2$: 483 (MH^+).

[01002] **N-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-tetrahydrofuran-2-carboxamide:** 1H NMR (400 MHz, d_6 -DMSO): 9.94 (s, 1H), 9.42 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.87 (d, 2H), 7.68 (d, 2H), 7.31 (d, 1H), 6.96 (d, 2H), 4.42 (m, 1H), 3.99 (m, 1H), 3.85 (m, 1H), 3.62 (m, 4H), 3.08 (m, 2H), 3.02 (m, 2H), 2.92 (m, 1H) 35 2.21 (m, 1H), 2.01 (m, 1H), 2.73 (m, 2H), 1.03 (d, 6H). MS (EI) for $C_{29}H_{34}N_6O_3$: 515 (MH^+).

[01003] **N-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide:** 1H NMR (400 MHz, d_6 -DMSO): 9.94 (s, 1H),

5 9.41 (s, 1H), 8.44 (d, 1H), 8.13 (d, 2H), 7.87 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 4.43 (m, 1H), 3.99 (m, 1H), 3.84 (m, 1H), 3.59 (m, 2H), 3.43 (m, 2H), 3.01 (m, 4H), 2.28-1.69 (m, 10H). MS (EI) for C₃₀H₃₄N₆O₃: 527 (MH⁺).

[01004] N-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 9.95 (s, 1H), 9.42 (s, 10 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.88 (d, 2H), 7.69 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 4.43 (m, 1H), 4.0 (m, 1H), 3.84 (m, 1H), 3.7 (m, 4H), 3.04 (m, 4H), 2.22 (m, 1H), 2.02 (m, 1H), 1.86 (m, 2H), 1.23 (s, 9H). MS (EI) for C₃₀H₃₆N₆O₃: 529 (MH⁺).

[01005] N-cyclopropyl-4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzamide: ¹H NMR (400 MHz, d₆-DMSO): 9.51 (s, br, 1H), 8.58 (s, br, 1H), 8.52 (d, 1H), 8.21 (d, 2H), 15 7.96 (d, 2H), 7.67 (d, 2H), 7.39 (d, 1H), 6.93 (d, 2H), 3.76 (m, 4H), 3.05 (m, 4H), 2.89 (m, 1H), 0.71 (m, 2H), 0.60 (m, 2H). MS (EI): 416 (MH⁺).

[01006] N-(2-methoxyethyl)-4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)benzamide: ¹H NMR (400 MHz, d₆-DMSO): 9.51 (s, br, 1H), 8.68 (s, br, 1H), 8.52 (d, 1H), 8.23 (d, 2H), 8.00 (d, 2H), 7.67 (d, 2H), 7.40 (d, 1H), 6.94 (d, 2H), 3.75 (m, 4H), 3.47 20 (m, 4H), 3.28 (s, 3H), 3.05 (m, 4H). MS (EI): 434 (MH⁺).

[01007] 2,6-dichloro-n-{3-[(4-pyridin-3-ylpyrimidin-2-yl)amino]propyl}-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 9.25 (br s, 1H), 8.68-8.66 (m, 2H), 8.37 (m, 2H), 7.49-7.47 (m, 3H), 7.42-7.38 (m, 1H), 7.32 (m, 1H), 7.21-7.20 (m, 1H), 3.44 (m, 2H), 3.29 (m, 2H), 1.82 (m, 2H). MS (EI): 402.0 (MH⁺).

25 **[01008] 2,6-dichloro-n-(3-{[4-(4-methyl-3,4-dihydro-2h-1,4-benzoxazin-7-yl)pyrimidin-2-yl]amino}propyl)benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 8.69 (t, 1H), 8.18 (d, 1H), 7.58 (dd, 1H), 7.51-7.40 (m, 4H), 7.02-6.97 (m, 2H), 6.73 (d, 1H), 4.25-4.23 (m, 2H), 3.41 (m, 2H), 3.32-3.29 (m, 4H), 2.91 (s, 3H), 1.81 (t, 2H). MS (EI): 472.3 (MH⁺).

[01009] 2,6-dichloro-n-(3-{[4-(2,3-dihydro-1,4-benzodioxin-6-yl)-6-methyl-pyrimidin-2-yl]amino}propyl)benzamide: ¹H-NMR (400MHz, d₆-DMSO): 8.68 (t, 1H), 7.62-7.59 (m, 2H), 7.51-7.49 (m, 2H), 7.44-7.40 (m, 1H), 7.02 (t, 1H), 6.97-6.92 (m, 2H), 4.30-4.28 (m, 4H), 3.44-3.43 (m, 2H), 3.32-3.29 (m, 2H), 2.27 (s, 3H), 1.80 (t, 2H); MS (EI): 473.3 (MH⁺).

[01010] N-(4-{2-[(3-{[(2,6-dichlorophenyl)carbonyl]amino}propyl)-amino]pyrimidin-4-yl}phenyl)morpholine-4-carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 8.69 (m, 2H), 35 8.33 (d, 1H), 8.18 (m, 1H), 7.60 (m, 2H), 7.51-7.49 (m, 2H), 7.44-7.34 (m, 2H), 7.20 (m, 1H), 7.01 (d, 1H), 3.62-3.61 (m, 4H), 3.43 (m, 6H), 3.32 (m, 2H), 1.83 (m, 2H). MS (EI): 529.1 (MH⁺).

- 5 **[01011] 2,6-dichloro-n-{3-[(4-{4-[(cyclopropylcarbonyl)amino]-phenyl}pyrimidin-2-yl)amino]propyl}benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.4 (s, 1H), 8.72 (t, 1H), 8.36-8.33 (m, 2H), 7.73 (m, 2H), 7.51-7.39 (m, 4H), 7.24 (m, 1H), 7.03 (d, 1H), 3.45 (m, 2H), 3.33 (m, 4H), 1.84-1.78 (m, 2H), 0.81-0.78 (m, 3H). MS (EI): 484.0 (MH⁺).
- [01012] N-(4-{2-[(3-{[(2,6-dichlorophenyl)carbonyl]amino}propyl)-amino]pyrimidin-**
10 **4-yl}phenyl)thiophene-2-carboxamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.4 (s, 1H), 8.72 (t, 1H), 8.44 (t, 1H), 8.37 (d, 1H), 8.05 (s, 1H), 7.90-7.81 (m, 3H), 7.50-7.39 (m, 4H), 7.25-7.23 (m, 2H), 7.07 (d, 1H), 3.47 (m, 2H), 3.34 (m, 2H), 1.85 (m, 2H). MS (EI): 526.0 (MH⁺).
- [01013] 2,6-dichloro-n-(3-{[4-(4-{[n-(2-morpholin-4-ylethyl)glycyl]-**
15 **amino}phenyl)pyrimidin-2-yl]amino}propyl)benzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.0 (br s, 1H), 8.72 (t, 1H), 8.35-8.32 (m, 2H), 7.82-7.75 (m, 2H), 7.51-7.40 (m, 4H), 7.22 (s, 1H), 7.05 (d, 1H), 3.56 (m, 4H), 3.45 (m, 2H), 3.30 (m, 3H), 2.64 (m, 2H), 2.41-2.35 (m, 8H), 1.84 (br s, 2H). MS (EI): 586.1 (MH⁺).
- [01014] 1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-ethanone:**
20 ¹H-NMR (400MHz, d₆-DMSO): 8.50 (d, 1H), 8.26-8.24 (m, 2H), 8.12-8.10 (m, 2H), 7.30 (s, 1H), 7.64-7.62 (m, 2H), 7.31 (d, 1H), 7.00-6.98 (m, 2H), 3.82-3.80 (m, 4H), 3.12-3.10 (m, 4H), 2.64 (s, 3H). MS (EI): 375.1 (MH⁺).
- [01015] (1e)-1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)ethanone**
25 **oxime:** ¹H-NMR (400MHz, d₆-DMSO): 11.4 (s, 1H), 9.82 (br s, 1H), 8.55 (d, 1H), 8.20 (d, 2H), 7.85-7.82 (m, 4H), 7.45 (d, 1H), 7.36 (br s, 1H), 4.69 (br, 1H), 3.91 (m, 4H), 3.34 (m, 4H), 2.21 (s, 3H). MS (EI): 388.1 (MH⁻).
- [01016] N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}-amino)pyrimidin-**
30 **4-yl]phenyl}-2-phenylacetamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.4 (s, 1H), 9.43 (br s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.75 (d, 2H), 7.67 (d, 2H), 7.33-7.20 (m, 2H), 7.07 (s, 2H), 6.97-6.95 (m, 4H), 3.82 (m, 4H), 3.67 (s, 2H), 3.03 (m, 4H), 2.07 (m, 1H), 0.75-0.69 (m, 4H). MS (EI): 533.2 (MH⁺).
- [01017] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-**
35 **bromobenzamide:** ¹H-NMR (400MHz, d₆-DMSO): 10.14 ppm (s, 1H), 8.42 ppm (t, 1H), 8.29 ppm (d, 1H), 8.06 ppm (d, 1H), 7.70 ppm (d, 2H), 7.65 ppm (m, 1H), 7.39 ppm (m, 3H), 7.12 ppm (t, 1H), 7.07 ppm (d, 1H), 3.33 ppm (br. m, 2H), 3.31 ppm (m, 2H), 2.07 ppm (s, 3H), 1.81 ppm (m, 2H); MS (EI) C₂₂H₂₂BrN₅O₂: 468 (MH⁺).
- [01018] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-**
fluorobenzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.17 ppm (s, 1H), 8.37 ppm (t, 1H),

5 8.30 ppm (d, 1H), 8.06 ppm (d, 2H), 7.70 ppm (d, 2H), 7.61 ppm (m, 1H), 7.50 ppm (m, 1H), 7.26 ppm (m, 2H), 7.16 ppm (t, 1H), 7.08 ppm (d, 1H), 3.41 ppm (br. m, 2H), 3.33 ppm (m, 2H), 2.08 ppm (s, 3H), 1.81 ppm (br. m, 2H); MS (EI) $C_{22}H_{22}FN_5O_2$: 408 (MH^+).

[01019] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-chlorobenzamide: 1H -NMR (400MHz, d_6 -DMSO): 10.14 ppm (s, 1H), 8.45 ppm (t, 1H),
10 8.28 ppm (d, 1H), 8.04 ppm (d, 2H), 7.68 ppm (d, 2H), 7.68 ppm (d, 2H), 7.47 ppm (m, 1H), 7.41 ppm (m, 2H), 7.35 ppm (m, 1H), 7.13 ppm (t, 1H), 7.06 ppm (d, 1H), 3.42 ppm (br. m, 2H), 3.29 ppm (m, 2H), 2.05 ppm (s, 3H), 1.78 ppm (br m, 2H); MS (EI) $C_{22}H_{22}BrN_5O_2$: 468 (MH^+).

[01020] N-[4-(2-({3-(morpholin-4-ylsulfonyl)phenyl}amino)pyrimidin-4-yl)phenyl]-acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.23 ppm (s, 1H), 10.12 ppm (s, 1H), 8.74 ppm (s, 1H), 8.58 ppm (d, 1H), 8.21 ppm (d, 2H), 7.93 ppm (m, 1H), 7.76 ppm (d, 2H), 7.60 ppm (t, 1H), 7.47 ppm (d, 1H), 7.30 ppm (m, 1H), 3.64 ppm (m, 4H), 2.90 ppm (m, 4H), 2.10 ppm (s, 3H); MS (EI) $C_{22}H_{23}N_5O_4S$: 454 (MH^+).

[01021] N-{4-[2-({3-[(cyclohexylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}-acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.43 ppm (s, 1H), 9.32 ppm (s, 1H), 8.46 ppm (d, 1H), 8.13 ppm (d, 2H), 7.77 ppm (d, 2H), 7.30 ppm (d, 2H), 7.17 ppm (s, 1H), 6.94 ppm (m, 2H), 6.22 ppm (m, 1H), 5.56 ppm (t, 1H), 2.87 ppm (t, 2H), 2.09 ppm (s, 3H), 1.85 ppm (br d, 2H), 1.69 ppm (br m, 2H), 1.64 ppm (br m, 1H), 1.19 ppm (m, 3H), 0.94 ppm (m, 2H); MS (EI) $C_{25}H_{29}N_5O$: 416 (MH^+).

25 [01022] N-(4-{2-[(3-[(5-bromo-2-fluorophenyl)methyl]amino)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.21 ppm (s, 1H), 9.37 ppm (s, 1H), 8.45 ppm (d, 1H), 8.12 ppm (d, 2H), 7.74 ppm (d, 2H), 7.55 ppm (m, 1H), 7.47 ppm (m, 1H), 7.31 ppm (d, 1H), 7.20 ppm (m, 1H), 7.05 ppm (m, 1H), 7.00 ppm (t, 1H), 6.28 ppm (t, 1H), 6.22 ppm (m, 1H), 4.32 ppm (d, 2H), 2.09 ppm (s, 3H); MS (EI) $C_{25}H_{21}BrFN_5O$: 507 (MH^+).

[01023] N-(4-{2-[(3-[(2,5-dimethylphenyl)methyl]amino)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.20 ppm (s, 1H), 9.33 ppm (s, 1H), 8.45 ppm (d, 1H), 8.13 ppm (d, 2H), 7.73 ppm (d, 2H), 7.30 ppm (d, 1H), 7.16 ppm (m, 2H), 7.06 ppm (d, 1H), 6.97 ppm (m, 3H), 6.23 ppm (m, 1H), 5.99 ppm (t, 1H), 4.17 ppm (d, 2H), 2.28 ppm (s, 3H), 2.21 ppm (s, 3H), 2.08 ppm (s, 3H); MS (EI) $C_{27}H_{27}N_5O$: 438 (MH^+).

35 [01024] N-(4-{2-[(3,4-dimorpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: 1H -NMR (400MHz, d_6 -DMSO): 10.22 ppm (s, 1H), 9.43 ppm (s,

5 1H), 8.46 ppm (d, 1H), 8.14 ppm (d, 2H), 7.74 ppm (d, 2H), 7.65 ppm (s, 1H), 7.34 ppm (m, 2H), 7.29 ppm (d, 2H), 6.88 ppm (d, 1H), 3.75 ppm (m, 8H), 3.15 ppm (br s, 4H), 3.05 ppm (br s, 4H), 2.09 ppm (s, 3H); MS (EI) C₂₆H₃₀N₆O₃: 475 (MH⁺).

[01025] N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.47 ppm (s, 1H), 9.41 ppm (s, 1H), 8.67 ppm (m, 2H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.89 ppm (m, 1H), 7.76 ppm (d, 2H), 7.69 ppm (d, 2H), 7.51 ppm (m, 1H), 7.28 ppm (d, 1H), 6.97 ppm (d, 2H), 3.80 ppm (s, 2H), 3.49 ppm (s, 2H), 3.13 ppm (br d, 4H), 1.83 ppm (m, 1H), 0.84 ppm (m, 4H); MS (EI) C₃₀H₂₉N₇O₂: 520 (MH⁺).

[01026] N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}butanamide: ¹H-NMR (400MHz, d₆-DMSO): 10.16 ppm (s, 1H), 9.41 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.77 ppm (d, 2H), 7.68 ppm (d, 2H), 7.28 ppm (d, 1H), 6.96 ppm (d, 2H), 3.63 ppm (m, 4H), 3.05 ppm (m, 4H), 2.92 ppm (m, 1H), 2.33 ppm (t, 1H), 1.63 ppm (m, 2), 1.02 ppm (d, 6H), 0.93 ppm (t, 3H); MS (EI) C₂₈H₃₄N₆O₂: 487 (MH⁺).

[01027] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide: ¹H-NMR (400MHz, d₆-DMSO): 10.15 ppm (s, 1H), 9.41 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.77 ppm (d, 2H), 7.68 ppm (d, 2H), 7.28 ppm (d, 1H), 6.95 ppm (d, 2H), 3.70 ppm (m, 4H), 3.05 ppm (m, 4H), 2.33 ppm (t, 2H), 1.63 ppm (m, 2H), 1.23 ppm (s, 9H), 0.93 ppm (t, 3H); MS (EI) C₂₉H₃₆N₆O₂: 501 (MH⁺).

25 [01028] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide: ¹H-NMR (400MHz, d₆-DMSO): 10.28 ppm (s, 1H), 9.41 ppm (s, 1H), 8.44 ppm (d, 1H), 8.11 ppm (d, 2H), 7.79 ppm (d, 2H), 7.68 ppm (d, 2H), 7.28 ppm (d, 1H), 6.95 ppm (d, 2H), 3.59 ppm (m, 2H), 3.46 ppm (m, 2H), 3.40 ppm (m, 1H), 3.02 ppm (m, 4H), 2.35 ppm (t, 2H), 2.14 ppm (m, 4H), 1.91 ppm (m, 1H), 1.75 ppm (m, 1H), 1.63 ppm (m, 2H), 0.93 ppm (t, 3H); MS (EI) C₂₉H₃₄N₆O₂: 499 (MH⁺).

[01029] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-2-carboxamide: ¹H-NMR (400MHz, d₆-DMSO): 10.91 (s, 1H), 9.41 (s, 1H), 8.78 (d, 1H), 8.47 (d, 2H), 8.17 (m, 4H), 7.69 (m, 2H), 7.33 (d, 1H), 6.95 (d, 2H), 6.80 (s, 2H), 3.75 (m, 4H), 3.06 (m, 4H). MS (EI) for C₂₆H₂₄N₆O₂: 453.5(MH⁺).

35 [01030] 2-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-benzamide: ¹H-NMR (400MHz, d₆-DMSO): 10.39 (s, 1H), 9.51 (s, 1H), 8.47 (s, 1H), 8.16

- 5 (d, 2H), 7.91 (d, 2H), 7.17 (m, 3H), 7.53 (t, 1H), 7.34 (s, 2H) 7.20 (d, 1H), 7.07 (m, 3H), 3.91 (m, 4H), 3.12 (m, 4H). MS (EI) for $C_{27}H_{25}N_5O_3$: 468.5(MH⁺).
- [01031] 3-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-benzamide:** 1H-NMR (400MHz, d6-DMSO): 10.49 (s, 2H), 8.19 (m, 3H), 7.97 (m, 3H), 7.55 (m, 2H), 7.50 (m, 3H), 7.20 (dd, 3H), 3.86 (m, 8H), 3.77 (s, 3H). MS (EI) for $C_{28}H_{27}N_5O_3$: 482.6(MH⁺).
- 10 **[01032] 4-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-benzamide:** 1H-NMR (400MHz, d6-DMSO): 10.35 (s, 1H), 9.40 (s, 1H), 8.45 (s, 1H), 8.16 (d, 2H), 7.98 (m, 4H), 7.68 (d, 2H), 7.31 (d, 1H), 7.09 (d, 2H), 6.94(d, 2H), 3.83 (s, 3H), 3.75 (m, 4H), 3.05 (m, 4H). MS (EI) for $C_{28}H_{27}N_5O_3$: 482.6(MH⁺).
- 15 **[01033] 4-chloro-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-benzamide:** 1H-NMR (400MHz, d6-DMSO):10.58 (s, 1H), 9.41 (s, 1H), 8.46 (s, 1H), 8.17 (d, 2H), 8.01 (d, 2H), 7.95 (d, 2H), 7.66 (m, 4H), 7.32 (d, 1H), 6.94 (d, 2H), 3.75 (m, 4H), 3.05 (m, 4H). MS (EI) for $C_{27}H_{24}ClN_5O_2$: 487.0(MH⁺).
- [01034] (2R)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide:** 1H-NMR (400MHz, d6-DMSO):9.95 (s, 1H), 9.38 (s, 1H), 8.45 (s, 1H), 8.12 (d, 2H), 7.89 (d, 2H), 7.66 (d, 2H), 7.29 (s, 1H), 6.92 (d, 2H), 4.44 (t, 2H), 4.00 (m, 2H), 3.85 (m, 2H), 2.41 (m, 4H), 2.20 (m, 2H), 2.02 (m, 2H), 1.88 (m, 2H), 1.05 (m, 4H). MS (EI) for $C_{27}H_{32}N_6O_2$: 473.6(MH⁺).
- 20 **[01035] (2S)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide:** 1H-NMR (400MHz, d6-DMSO):9.94 (s, 1H, 9.38 (s, 1H), 8.44 (s, 1H), 8.12 (d, 2H), 7.87 (d, 2H), 7.65 (d, 2H), 7.29 (s, 1H), 6.99 (d, 2H), 4.43 (m, 2H), 3.99 (m, 2H), 3.86 (m, 2H), 2.43 (m, 2H), 2.22 (m, 2H), 2.02 (m, 2H), 1.88 (m, 3H), 1.05 (m, 5H) . MS (EI) for $C_{27}H_{32}N_6O_2$: 473.6(MH⁺).
- 25 **[01036] 1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}-phenyl)-L-prolinamide:** 1H-NMR (400MHz, d6-DMSO): 10.31 (s, 1H), 9.39 (s, 1H), 8.44 (d, 1H), 8.14 (d, 2H), 7.81 (d, 2H), 7.67 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 5.05 (s, br, 1H), 3.74 (m, 4H), 3.59 (m, 1H), 3.49 (m, 1H), 3.23 (m, 2H), 3.05 (m, 4H), 2.77 (m, 1H), 2.63 (m, 1H), 2.41 (m, 1H), 2.16 (m, 1H), 1.80 (m, 3H). MS (EI) for $C_{27}H_{32}N_6O_3$: 489.6(MH⁺).
- 30 **[01037] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-carboxamide:** 1H-NMR (400MHz, d6-DMSO): 10.47 (s, 1H), 9.45 (s, 1H), 8.47 (s, 1H), 8.17 (d, 2H), 8.08 (s, 1H), 7.92 (m, 3H), 7.70 (s, 2H), 7.32 (s, 1H), 7.26 (t, 1H), 6.99 (s, 2H), 3.76 (m, 4H), 3.09 (m, 4H). MS (EI) for $C_{25}H_{23}N_5O_2S$: 458.6(MH⁺).
- 35

- 5 **[01038] N-[4-(2-([4-(4-ethylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-tetrahydrofuran-3-carboxamide:** ¹H-NMR (400MHz, d6-DMSO): 10.31 (s, 1H), 9.37 (s, 1H), 8.44 (s, 1H), 8.14 (m, 2H), 7.79 (d, 2H), 7.65 (d, 2H), 7.27 (d, 1H), 6.92 (d, 2H), 3.96 (t, 1H), 3.76 (m, 3H), 3.19 (m, 1H), 3.09 (m, 4H), 2.55 (m, 4H), 2.42 (m, 2H), 2.10 (m, 2H), 1.05 (t, 3H). MS (EI) for C₂₇H₃₂N₆O₂: 473.6(MH⁺).
- 10 **[01039] N-(4-{2-[4-(4-nicotinoylpiperazin-1-yl)phenylamino]pyrimidin-4-yl}phenyl)-2-phenylacetamide:** ¹H NMR (400MHz, d6-DMSO): 8.8 (d, 2H), 8.40 (s, 1H), 8.05 (d, 2H), 7.80 (d, 1H), 7.60-7.2 (m, 10H), 7.1-6.90 (m, 5H), 3.80-3.60 (m, 4H), 3.7 (s, 2H) 3.20-3.25 (m, 4H); MS (EI) for C₃₄H₃₁N₇O₂: 570 (MH⁺).
- [01040] 3-([4-[4-(acetylamino)phenyl]pyrimidin-2-yl]amino)-N-(diphenylmethyl)benzamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.78 (s, 1H), 9.24 (d, 1H), 8.52 (d, 1H), 8.39 (s br, 1H), 8.16 (d, 2H), 7.95 (d, 1H), 7.75 (d, 2H), 7.53 (d t, 1H), 7.39-7.43 (m, 3H), 7.38 (d, 3H), 7.32-7.36 (m, 4H), 7.24-7.28 (m, 2H), 6.43 (d, 1H), 2.10 (s, 3H). MS (EI) for C₃₂H₂₇N₅O₂: 514.3 (MH⁺).
- [01041] N-[4-(2-([4-(4-methylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.24 (s, 1H), 9.35 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.65 (d, 2H), 7.26 (d, 1H), 6.93 (d, 2H), 3.07 (m, 4H), 2.45 (m, 4H), 2.22 (s, 3H), 2.09 (s, 3H). MS (EI) for C₂₃H₂₆N₆O: 403.4(MH⁺).
- 20 **[01042] N-{4-[2-([4-[4-(phenylcarbonyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]-phenyl}acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.45 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.73-7.79 (m, 5H), 7.28 (d, 1H), 6.98 (d, 2H), 3.78 (m, 2H), 3.48 (m, 2H), 3.15 (m, 2H), 3.07 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₈N₆O₂: 493.4 (MH⁺).
- 25 **[01043] N-{4-[2-([4-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]-phenyl}acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.27 (s, 1H), 6.96 (d, 2H), 3.60 (m, 4H), 3.04 (m, 4H), 2.37 (d, 2H), 2.15 (m, 1H), 2.09 (s, 3H), 1.61-1.78 (m, 2H), 1.53-1.59 (m, 2H), 1.46-1.52 (m, 2H), 1.09-1.77 (m, 2H). MS (EI) for C₂₉H₃₄N₆O₂: 499.3 (MH⁺).
- 30 **[01044] N-{4-[2-([4-[4-(cyclohexylcarbonyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]-phenyl}acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.96 (d, 2H), 3.61 (m, 4H), 3.04 (m, 4H), 2.09 (s, 3H), 1.62-1.70 (m, 6H), 1.26-1.38 (m, 5H). MS (EI) for C₂₉H₃₄N₆O: 499.2 (MH⁺).
- 35

- 5 **[01045] N-(4-{2-[(4-{4-[(2-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.56 (d, 1H), 7.41-7.49 (m, 3H), 7.27 (d, 1H), 6.96 (d, 2H), 3.81 (m, 2H), 3.28 (m, 2H), 3.16 (m, 2H), 3.05 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₇ClN₆O₂: 527.8 (MH⁺).
- 10 **[01046] N-(4-{2-[(4-{4-[(3-fluorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.53 (m, 1H), 7.27-7.35 (m, 4H), 6.97 (d, 2H), 3.77 (m, 2H), 3.46 (m, 2H), 3.16 (m, 2H), 3.07 (m, 2H), 2.09 (s, 3H). MS(EI) for C₂₉H₂₇FN₆O₂: 511.5 (MH⁺).
- 15 **[01047] N-(4-{2-[(4-{4-[(3-fluoro-4-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.38 (m, 1H), 7.28 (d, 1H), 7.25 (d, 1H), 7.18 (d, 1H), 6.56 (d, 2H), 3.75 (m, 2H), 3.49 (m, 2H), 3.14 (m, 2H), 3.07 (m, 2H), 2.28 (d, 3H), 2.09 (s, 3H). MS(EI) for C₃₀H₂₉FN₆O₂: 525.7 (MH⁺).
- 20 **[01048] N-(4-{2-[(4-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.45 (s, 1H), 8.11 (d, 2H), 7.75 (m, 3H), 7.69 (d, 2H), 7.45 (dd, 1H), 7.28 (d, 1H), 6.97 (d, 2H), 3.77 (m, 2H), 3.47 (m, 2H), 3.16 (m, 2H), 3.07 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₆Cl₂N₆O₂: 562.3(MH⁺).
- 25 **[01049] N-(4-{2-[(4-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.54 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 3.77 (m, 2H), 3.45 (m, 2H), 3.17 (m, 2H), 3.08 (m, 2H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₆Cl₂N₆O₂: 562.6(MH⁺).
- 30 **[01050] N-[4-(2-{[4-(4-{[3-(methyloxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino}-pyrimidin-4-yl)phenyl)acetamide:** ¹H-NMR(400MHz, d6-DMSO): 10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.38 (t, 1H), 7.28 (d, 1H), 7.04 (dd, 1H), 6.95-6.99 (m, 4H), 3.79 (s, 3H), 3.77 (m, 2H), 3.47 (m, 2H), 3.15 (m, 2H), 3.06 (m, 2H), 2.09 (s, 3H). MS(EI) for C₃₀H₃₀N₆O₃: 523.5(MH⁺).
- 35 **[01051] N-(4-{2-[(4-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.53 (d,

5 2H), 7.49 (d, 2H), 7.27 (d, 1H), 6.97 (d, 2H), 3.77 (m, 2H), 3.47 (m, 2H), 3.15 (m, 2H), 3.06 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{27}ClN_6O_2$: 527.8(MH⁺).

[01052] N-(4-{2-[(4-{4-[(4-methylphenyl)carbonyl]piperazin-1-

yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO):

10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.34 (d, 10 2H), 7.27 (m, 3H), 6.97 (d, 2H), 3.75 (m, 2H), 3.51 (m, 2H), 3.10 (m, 4H), 2.36 (s, 3H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{30}N_6O_2$: 507.3 (MH⁺).

[01053] N-(4-{2-[(4-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-

yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR (400MHz, d₆-DMSO):

10.21 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 15 1H), 6.98 (d, 2H), 6.92 (t, 1H), 6.37 (dd, 1H), 6.05 (dd, 1H), 3.76 (m, 4H), 3.69 (s, 3H), 3.11 (m, 4H), 2.09 (s, 3H), MS (EI) for $C_{28}H_{29}N_7O_2$: 496.4(MH⁺).

[01054] N-{4-[2-({4-[4-(furan-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-

yl]-phenyl}acetamide: ¹H-NMR(400MHz,d₆-DMSO):10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.87 (d, 1H), 7.74 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 7.04 (d, 1H), 6.98 (d, 20 2H), 6.66 (dd, 1H), 3.82 (m, 4H), 3.14 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{27}H_{26}N_6O_3$: 483.3 (MH⁺).

[01055] N-[4-(2-({4-(4-{2-[(4-fluorophenyl)oxy]acetyl}piperazin-1-yl)phenyl}amino)-

pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR (400MHz, d₆-DMSO): 10.22 (s, 1H), 9.41 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 7.12 (m, 2H), 6.94- 25 6.99 (m, 4H), 4.87 (s, 2H), 3.61 (m, 4H), 3.13 (m, 2H), 3.06 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{29}FN_6O_3$:541.4(MH⁺).

[01056] N-(4-{2-[(4-{4-[(3-methylphenyl)sulfonyl]piperazin-1-

yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: MS (EI) for $C_{29}H_{30}N_6O_3S$: 543.5 (MH⁺).

30 **[01057] N-{4-[2-({4-[4-(phenylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-**
phenyl}acetamide: ¹H-NMR(400MHz,d₆-DMSO): 10.21 (s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.80 (m, 1H), 7.78 (d, 1H), 7.74-7.77 (m, 2H), 7.70-7.73 (m, 2H), 7.67-7.69 (m, 1H), 7.65 (d, 2H), 7.27 (d, 1H), 6.90 (d, 2H), 3.15 (m, 4H), 3.02 (m, 4H), 2.09 (s,3H). MS (EI) for $C_{26}H_{26}N_6O_3S_2$: 529.4(MH⁺).

35 **[01058] N-{4-[2-({4-[4-(2-thienylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-**
yl]-phenyl}acetamide: ¹H-NMR(400MHz,d₆-DMSO):10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d,

- 5 1H), 8.10 (m, 3H), 7.74 (d, 2H), 7.70 (dd, 1H), 7.66 (d, 2H), 7.33 (dd, 1H), 7.27 (d, 1H), 6.93 (d, 2H), 3.19 (m, 4H), 3.07 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₆H₂₆N₆O₃S₂: 535.6(MH⁺).
- [01059] N-(4-{2-[(4-{4-[(4-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO):10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.10 (d, 2H), 7.87 (m, 2H), 7.74 (d, 2H), 7.66 (d, 2H), 7.53 (m, 2H), 7.27 (d, 1H), 6.91 (d, 2H), 3.15 (m, 4H), 3.04 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₈H₂₇FN₆O₃S: 547.4 (MH⁺).
- [01060] N-[4-(2-{[4-(4-{[4-(methyloxy)phenyl]sulfonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO): MS (EI) for C₂₉H₃₀N₆O₄S: 559.9 (MH⁺).
- 15 **[01061] N-(4-{2-[(4-{4-[(4-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO):10.21 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.72-7.82 (d, 6H), 7.66 (d, 2H), 7.27 (d, 1H), 6.91 (d, 2H), 3.15 (m, 4H), 3.05 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₈H₂₇ClN₆O₃S: 563.9 (MH⁺).
- 20 **[01062] N-(4-{2-[(4-{4-[(3-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO):10.21 (s, 1H), 9.40 (s, 1H), 8.43 (d, 1H), 8.10 (d, 2H), 7.85 (m, 1H), 7.76-7.81 (m, 2H), 7.73 (d, 3H), 7.66 (d, 2H), 7.27 (d, 1H), 6.91 (d, 2H), 3.14 (m, 4H), 3.09 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₈H₂₇ClN₆O₃S: 5640 (MH⁺).
- 25 **[01063] N-{4-[2-({4-[4-(biphenyl-4-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR(400MHz,d6-DMSO):10.21(s, 1H), 9.39 (s, 1H), 8.43 (d, 1H), 8.08 (d, 2H), 7.96 (d, 2H), 7.86 (d, 2H), 7.77 (d, 2H), 7.73 (d, 2H), 7.66 (d, 2H), 7.51-7.55 (m, 2H), 7.46 (m, 1H), 7.26 (d, 1H), 6.91 (d, 2H), 3.18 (m, 4H), 3.08 (m, 4H), 2.09 (s, 3H). MS (EI) for C₃₄H₃₂N₆O₃S: 605.8 (MH⁺).
- 30 **[01064] N-{4-[2-({4-[4-(naphthalen-1-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.21 (s, 1H), 9.38 (s, 1H), 8.72 (d, 1H), 8.42 (d, 1H), 8.33 (d, 1H), 8.20 (dd, 1H), 8.14 (d, 1H), 8.09 (d, 2H), 7.74-7.79 (m, 1H), 7.73 (d, 2H), 7.62-7.70 (m, 2H), 7.64 (d, 2H), 7.26 (d, 1H), 6.88 (d, 2H), 3.21 (m, 4H), 3.09 (m, 4H), 2.09 (s, 3H). MS (EI) for C₃₂H₃₀N₆O₃S: 579.6 (MH⁺).
- 35 **[01065] N-(4-{2-[(3-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.21

- 5 (s, 1H), 9.46 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.74 (d, 2H), 7.64 (s br, 1H), 7.36 (dd, 1H), 7.33 (d, 1H), 7.20-7.27 (m, 2H), 7.13 (t, 1H), 6.92-7.00 (m, 2H), 6.55 (d, 1H), 3.54 (s, 2H), 3.16 (m, 4H), 2.57 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{29}ClN_6O$: 513.8 (MH⁺).
- [01066]** N-[4-(2-({3-(4-({3-(methyloxy)phenyl)methyl}piperazin-1-yl)phenyl}amino)-pyrimidin-4-yl)phenyl]acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.21 (s, 1H), 9.46 (s, 1H), 8.48 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.62 (s, 1H), 7.33 (d, 1H), 7.22-7.27 (m, 2H), 7.13 (t, 1H), 6.19 (m, 2H), 6.83 (d, 1H), 6.55 (d, 1H), 3.74 (s, 3H), 3.52 (s, 2H), 3.16 (m, 4H), 2.55 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{30}H_{32}N_6O_2$: 509.8 (MH⁺).
- [01067]** N-{4-[2-({3-[4-(3-methylbutyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.22 (s, 1H), 9.47 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.68 (s, 1H), 7.33 (d, 1H), 7.19 (d, 1H), 7.13 (t, 1H), 6.55 (d, 1H), 3.15 (m, 4H), 2.54 (m, 4H), 2.34 (t, 2H), 2.09 (s, 3H), 1.57-1.62 (m, 1H), 1.34-1.40 (m, 2H), 0.90 (d, 6H). MS (EI) for $C_{27}H_{34}N_6O$: 459.7 (MH⁺).
- [01068]** N-{4-[2-({3-[4-(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): MS
20 (EI) for $C_{31}H_{32}N_6O_3$: 537.5 (MH⁺).
- [01069]** N-{4-[2-({3-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.22 (s, 1H), 9.47 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.69 (s, 1H), 7.33 (d, 1H), 7.19 (d, 1H), 7.13 (t, 1H), 6.56 (d, 1H), 3.17 (m, 4H), 2.60 (m, 4H), 2.24 (d, 2H), 2.09 (s, 3H), 0.88 (m, 1H), 0.47-0.51 (m, 2H),
25 0.11-0.13 (m, 2H). MS (EI) for $C_{28}H_{30}N_6O$: 443.8 (MH⁺).
- [01070]** N-(4-{2-[(3-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.22 (s, 1H), 9.47 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.68 (s, 1H), 7.33 (d, 1H), 7.20 (d, 1H), 7.14 (t, 1H), 6.56 (d, 1H), 3.15 (m, 4H), 2.56 (m, 4H), 2.41 (t, 4H), 2.09 (s, 3H), 2.06 (s, 3H), 1.71-1.78 (m, 2H).
30 MS (EI) for $C_{26}H_{32}N_6OS$: 477.5 (MH⁺).
- [01071]** N-(4-{2-[(3-{4-[(4-({3-(dimethylamino)propyl}oxy)phenyl)methyl]piperazin-1-yl}-phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.22 (s, 1H), 9.46 (s, 1H), 8.48 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.62 (s, 1H), 7.33 (d, 1H), 7.23 (m 3H), 7.13 (t, 1H), 6.88 (d, 2H), 6.55 (d, 1H), 3.97 (t, 2H), 3.46 (s, 2H), 3.14 (m, 4H), 2.55 (m, 4H), 2.34 (t, 2H), 2.14 (s, 6H), 2.10 (s, 3H), 1.80-1.85 (m, 2H). MS (EI) for $C_{34}H_{41}N_7O_2$:
35 580.5 (MH⁺).

- 5 [01072] N-{4-[2-({3-[4-({3-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)-amino)pyrimidin-4-yl}phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO):10.21 (s, 1H), 9.47 (s, 1H), 8.49 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.63 (s, 1H), 7.47 (t, 1H), 7.41 (d, 1H), 7.33 (d, 2H), 7.22-7.28 (m, 2H), 7.13 (t, 1H), 6.56 (dd, 1H), 3.62 (s, 2H), 3.16 (m, 4H), 2.56 (m, 4H), 2.09 (s, 3H). MS (EI) for C₃₀H₂₉F₃N₆O₂: 563.7(MH+).
- 10 [01073] 4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-phenylpiperazine-1-carboxamide: ¹H-NMR(400MHz,d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.63 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.75 (d, 2H), 7.70 (d, 2H), 7.48 (d, 2H), 7.21-7.28 (m, 3H), 7.00 (d, 2H), 6.94 (t, 1H), 3.61 (m, 4H), 3.11 (m, 4H), 2.09 (s, 3H). MS (EI) for C₂₉H₂₉N₇O₂: 508.6 (MH+).
- 15 [01074] N-[4-(2-({3-(4-propanoylpiperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]-acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.68 (s, 1H), 7.34 (d, 1H), 7.26 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.62 (m, 4H), 3.13 (m, 4H), 2.35-2.39 (m, 2H), 2.09 (s, 3H), 1.02 (t, 3H). MS (EI) for C₂₅H₂₈N₆O₂: 445.4 (MH+).
- 20 [01075] N-{4-[2-({3-[4-(phenylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.21 (s, 1H), 9.48 (s, 1H), 8.47 (d, 1H), 8.11 (d, 2H), 7.72 (d, 2H), 7.63 (s, 1H), 7.41-7.47 (m, 5H), 7.31 (d, 1H), 7.26 (d, 1H), 7.14 (t, 1H), 6.56 (dd, 1H), 3.78 (m, 2H), 3.48 (m, 2H), 3.27 (m, 2H), 3.12 (m, 2H), 2.08 (s, 3H). MS (EI) for C₂₉H₂₈N₆O₂: 493.7 (MH+).
- 25 [01076] N-{4-[2-({3-[4-(2-phenylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.25 (s, 1H), 9.50 (s, 1H), 8.49 (d, 1H), 8.13 (d, 2H), 7.76 (d, 2H), 7.63 (s, 1H), 7.33 (d, 1H), 7.30 (d, 2H), 7.26 (m, 3H), 7.22 (m, 1H), 7.15 (t, 1H), 6.56 (dd, 1H), 3.79 (s, 2H), 3.66 (m, 4H), 3.11 (m, 2H), 3.05 (m, 2H), 2.09 (s, 3H). MS (EI) for C₃₀H₃₀N₆O₂: 507.7 (MH+).
- 30 [01077] N-{4-[2-({3-[4-(cyclopentylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.52 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.71 (s, 1H), 7.34 (d, 1H), 7.24 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.66 (m, 4H), 3.13 (m, 4H), 3.00-3.07 (m, 1H), 2.09 (s, 3H), 1.80 (m, 2H), 1.51-1.71 (m, 6H). MS (EI) for C₂₈H₃₂N₆O₂: 485.7 (MH+).
- 35 [01078] N-{4-[2-({3-[4-(2-pyridin-3-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]-phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.43-8.46 (m, 2H), 8.14 (d, 2H), 7.76 (d, 2H), 7.63-7.67 (m, 2H), 7.32-7.35 (m, 2H),

5 7.26 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.84 (s, 2H), 3.73 (m, 2H), 3.66 (m, 2H), 3.15 (m, 4H), 2.08 (s, 3H). MS (EI) for $C_{29}H_{29}N_7O_2$: 508.4 (MH⁺).

[01079] N-{4-[2-({3-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.70 (s, 1H), 7.34 (d, 1H), 7.25 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.61 (m, 4H), 3.12 (m, 4H), 2.39 (d, 2H), 2.11-2.19 (m, 1H), 2.09 (s, 3H), 1.72-7.78 (m, 2H), 1.52-1.59 (m, 2H), 1.47-1.52 (m, 2H), 1.09-1.18 (m, 2H). MS (EI) for $C_{29}H_{34}N_6O_2$: 499.4 (MH⁺).

[01080] N-(4-{2-[(3-{4-[(2-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.50 (s, 1H), 8.49 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.63 (s, 1H), 7.56 (d, 1H), 7.42-7.50 (m, 3H), 7.33 (d, 1H), 7.29 (d, 1H), 7.16 (t, 1H), 6.59 (dd, 1H), 3.79 (m, 2H), 3.50 (m, 2H), 3.24 (m, 2H), 3.15 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{27}ClN_6O_2$: 527.9 (MH⁺).

[01081] N-(4-{2-[(3-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.23 (s, 1H), 9.51 (s, 1H), 8.49 (d, 1H), 8.14 (d, 2H), 7.74 (d, 2H), 7.66 (s, 1H), 7.53 (d, 2H), 7.49 (d, 2H), 7.34 (d, 1H), 7.27 (d, 1H), 7.17 (t, 1H), 6.59 (dd, 1H), 3.79 (m, 2H), 3.50 (m, 2H), 3.24 (m, 2H), 3.15 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{27}ClN_6O_2$: 528.1 (MH⁺).

[01082] N-(4-{2-[(3-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.23 (s, 1H), 9.51 (s, 1H), 8.49 (d, 1H), 8.13 (d, 2H), 7.74 (m, 4H), 7.60 (s, 1H), 7.46 (dd, 1H), 7.34 (d, 1H), 7.27 (d, 1H), 7.17 (t, 1H), 6.59 (dd, 1H), 3.79 (m, 2H), 3.50 (m, 2H), 3.24 (m, 2H), 3.15 (m, 2H), 2.09 (s, 3H). MS (EI) for $C_{29}H_{26}Cl_2N_6O_2$: 562.7 (MH⁺).

[01083] N-(4-{2-[(3-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)acetamide: ¹H-NMR(400MHz,d6-DMSO): 10.23 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.75 (d, 2H), 7.67 (s, 1H), 7.34 (d, 1H), 7.28 (d, 1H), 7.17 (t, 1H), 6.92 (m, 1H), 6.59 (dd, 1H), 6.37 (dd, 1H), 6.05 (m, 1H), 3.80 (m, 4H), 3.69 (s, 3H), 3.19 (m, 4H), 2.09 (s, 3H). MS (EI) for $C_{28}H_{29}N_7O_2$: 450.7 (MH⁺).

[01084] N²,N²-dimethyl-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl]glycinamide: ¹H-NMR(400MHz,d6-DMSO): 10.0 (s, 1H), 9.48 (s, 1H), 8.48 (d, 1H), 8.15 (d, 2H), 7.84 (d, 2H), 7.65 (s, 1H), 7.29-7.33 (m, 2H), 6.86 (d, 1H), 3.81 (s, 3H), 3.72 (m, 4H), 3.11 (s, 2H), 2.92 (m, 4H), 2.29 (s, 6H). MS (EI) for $C_{25}H_{30}N_6O_3$: 463.8 (MH⁺).

- 5 **[01085] 3-(methyloxy)-N-[4-(2-{{3-(methyloxy)-4-morpholin-4-ylphenyl}amino}pyrimidin-4-yl)phenyl]propanamide:** ¹H-NMR(400MHz,d6-DMSO): 10.21 (s, 1H), 9.46 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.74 (d, 2H), 7.64 (s, 1H), 7.26-7.30 (m, 2H), 6.84 (d, 1H), 3.79 (s, 3H), 3.70 (m, 4H), 3.61 (t, 2H), 3.23 (s, 3H), 2.89 (m, 4H), 2.57 (t, 2H). MS (EI) for C₂₅H₂₉N₅O₄: 464.8 (MH⁺).
- 10 **[01086] N-(4-{2-[(4-{(2-chlorophenyl)sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.19 (s, 1H), 9.38 (s, 1H), 8.42 (d, 1H), 8.08 (d, 2H), 8.00 (dd, 1H), 7.68-7.74 (m, 4H), 7.65 (d, 2H), 7.58 (m, 1H), 7.25 (d, 1H), 6.91 (d, 2H), 3.15 (m, 4H), 3.02 (m, 4H), 2.07 (s, 3H). MS (EI) for C₂₈H₂₇ClN₆O₃S: 563.9 (MH⁺).
- 15 **[01087] N-{4-[2-({3-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.52 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.71 (s, 1H), 7.34 (d, 1H), 7.25 (d, 1H), 7.17 (t, 1H), 6.60 (dd, 1H), 3.56 (m, 2H), 3.65 (m, 2H), 3.20 (m, 2H), 3.13 (m, 2H), 2.09 (s, 3H), 2.04 (m, 1H), 0.77-0.49 (m, 4H). MS (EI) for C₂₆H₂₈N₆O₂: 457.5(MH⁺).
- 20 **[01088] N-{4-[2-({3-[4-(2-cyclopropylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.24 (s, 1H), 9.51 (s, 1H), 8.50 (d, 1H), 8.14 (d, 2H), 7.76 (d, 2H), 7.70 (s, 1H), 7.34 (d, 1H), 7.25 (d, 1H), 7.16 (t, 1H), 6.58 (dd, 1H), 3.62 (m, 4H), 3.13 (m, 4H), 2.32 (d, 2H), 2.09 (s, 3H), 0.99 (m, 1H), 0.45 (m, 2H), 0.14 (m, 2H). MS (EI) for C₂₇H₃₀N₆O₂: 477.5(MH⁺).
- 25 **[01089] N-[4-(2-{{3-(4-{{3-(methyloxy)phenyl}carbonyl}piperazin-1-yl)phenyl}amino}-pyrimidin-4-yl)phenyl]acetamide:** ¹H-NMR(400MHz,d6-DMSO): 10.22 (s, 1H), 9.48 (s, 1H), 8.47 (d, 1H), 9.11 (d, 2H), 7.72 (d, 2H), 7.62 (s, 1H), 7.35 (t, 1H), 7.31 (d, 1H), 7.26 (d, 1H), 7.14 (t, 1H), 7.01 (m, 1H), 6.97 (m, 2H), 6.56 (dd, 1H), 3.76 (s, 3H), 3.73 (m, 2H), 3.47 (m, 2H), 3.14 (m, 4H), 2.07 (s, 3H). MS (EI) for C₃₀H₃₀N₆O₃: 523.7 (MH⁺).
- 30 **[01090] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H-NMR (400MHz, d6-DMSO): 10.21 (s, 1H), 9.40 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.74 (d, 2H), 7.69 (d, 2H), 7.27 (d, 1H), 6.96 (d, 2H), 3.70 (m, 4H), 3.05 (m, 4H), 2.09 (s, 3H), 1.23 (s, 9H), MS(EI) for C₂₇H₃₂N₆O₂: 473.4(MH⁺).
- 35 **[01091] 2,6-dichloro-N-(3-(4-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)pyrimidin-2-ylamino)propyl)benzamide:** (400 MHz, CDCl₃): 8.16 (br, 1H), 8.0 – 8.8 (m, 2H), 7.26 (m,

- 5 4H), 6.82 (d, 1H), 6.75 (br, 1H), 5.4 (t, 1H), 4.29 (m, 4H), 3.68 (m, 2H), 3.56 (m, 2H), 1.92 (m, 2H). MS (EI): 459 (MH⁺).
- [01092] 1-(4-(4-(4-acetamidophenyl)pyrimidin-2-ylamino)phenyl)piperidine-3-carboxylic acid: MS (EI) for C₂₄H₂₅N₅O₃: 432 (MH⁺).
- [01093] tert-butyl methyl(2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylamino)-2-oxoethyl)carbamate: MS (EI) for C₂₈H₃₄N₆O₄: 519 (MH⁺).
- 10 [01094] tert-butyl 4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenylcarbamate: MS (EI) for C₂₇H₃₄N₆O₂: 475 (MH⁺).
- [01095] 2-(dimethylamino)-N-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide: NMR (400 MHz, d₆-DMSO): 10.0 (s, 1H), 9.37 (s, 1H), 8.41 (d, 1H), 8.11 (d, 1H), 7.85 (d, 2H), 7.63 (f, 2H), 7.46 (d, 1H), 7.25 (d, 1H), 6.83 – 6.92 (m, 2H), 3.11 (m, 4H), 2.51 (m, 4H), 2.37 (q, 2H), 2.36 (s, 6H), 2.26 (s, 2H), 1.05 (t, 3H). MS (EI): 460 (MH⁺).
- 15 [01096] 4-(4-aminophenyl)-N-(4-(4-ethylpiperazin-1-yl)phenyl)pyrimidin-2-amine: MS (EI) for C₂₂H₂₆N₆: 375 (MH⁺).
- 20 [01097] (S)-tert-butyl 1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylamino)-1-oxo-3-phenylpropan-2-ylcarbamate: MS (EI) for C₃₄H₃₈N₆O₄: 595 (MH⁺).
- [01098] (R)-tert-butyl 1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylamino)-1-oxo-3-phenylpropan-2-ylcarbamate: MS (EI) for C₃₄H₃₈N₆O₄: 595 (MH⁺).
- 25 [01099] (R)-2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-3-phenylpropanamide: NMR (400 MHz, d₆-DMSO): 11.40 (s, 1H), 10.20 (s, 1H), 8.43 – 8.62 (m, 3H), 8.17 (d, 2H), 7.91 (d, 2H), 7.89 (d, 2H), 7.84 (m, 2H), 7.20 – 7.38 (m, 4H), 4.10 (m, 4H), 3.63 (m, 2H), 3.40 – 3.57 (m, 6H), 3.20 (m, 1H), MS (EI): 495 (MH⁺).
- 30 [01100] (S)-2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-3-phenylpropanamide: NMR (400 MHz, d₆-DMSO): 11.40 (s, 1H), 10.20 (s, 1H), 8.43 – 8.62 (m, 3H), 8.17 (d, 2H), 7.91 (d, 2H), 7.89 (d, 2H), 7.84 (m, 2H), 7.20 – 7.38 (m, 4H), 4.10 (m, 4H), 3.63 (m, 2H), 3.40 – 3.57 (m, 6H), 3.20 (m, 1H), MS (EI): 495 (MH⁺).
- [01101] (S)-2-amino-N-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methylbutanamide: MS (EI) for C₂₇H₃₅N₇O: 474 (MH⁺).
- 35 [01102] (R)-2-amino-N-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methylbutanamide: MS (EI) for C₂₇H₃₅N₇O: 474 (MH⁺).

- 5 **[01103] 1-ethyl-3-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea:** NMR (400 MHz, d6-DMSO): 9.41 (s, 1H), 8.75 (s, 1H), 8.42 (d, 1H), 8.10 (d, 2H), 7.64 (s, 1H), 7.54 (d, 2H), 7.26 (m, 2H), 6.85 (d, 1H), 6.21 (br, 1H), 3.79 (s, 3H), 3.70 (m, 4H), 3.11 (q, 2H), 2.89 (m, 4H), 1.06 (t, 3H). MS (EI): 449 (MH⁺).
- [01104] (R)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperidine-2-**
10 **carboxamide:** NMR (400 MHz, d6-DMSO): 11.36 (s, 1H), 10.0 (s, 1H), 9.4 (d, 1H), 8.84 (m, 1H), 8.57 (d, 1H), 8.2 (d, 2H), 7.82 (m, 4H), 7.6 (br, 1H), 7.4 (d, 1H), 4.0 (m, 4H), 3.82 (m, 1H), 3.42 (m, 4H), 3.23 (m, 1H), 2.94 (m, 1H), 2.3 (m, 1H), 1.82 (m, 1H), 1.54 – 1.92 (m, 4H). MS (EI): 458 (MH⁺).
- [01105] N-{4-(2-([4-(4-ethylpiperazin-1-yl)phenyl]amino)-5-methylpyrimidin-4-**
15 **yl)phenyl]-acetamide:** ¹H NMR (400 MHz, d6-DMSO): 10.1 (s, 1H), 9.23 (s, 1H), 8.31 (s, 1H), 7.6-7.7 (m, 6H), 6.87 (d, 2H), 3.04 (m, 4H), 2.48 (m, 4H), 3.05 (m, 4H), 2.36 (q, 2H), 2.2 (s, 3H), 2.08 (s, 3H), 1.03 (s, 3H). MS (EI): 431 (MH⁺).
- [01106] 4-{4-[(4-{4-[(N,N-dimethylglycyl)amino]phenyl}pyrimidin-2-**
20 **yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.43 (d, 1H), 8.12 (d, 2H), 7.82 (d, 2H), 7.68 (d, 2H), 7.27 (d, 1H), 6.97 (d, 2H), 3.42 (m, 4H), 3.12 (s, 2H), 3.06 (q, 2H), 3.02 (m, 4H), 2.3 (s, 6H), 1.11 (t, 3H). MS (EI): 503 (MH⁺).
- [01107] N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-**
25 **yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N², N²-dimethylglycinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.44 (d, 1H), 8.11 (d, 2H), 7.83 (d, 2H), 7.69 (d, 2H), 7.29 (d, 1H), 6.96 (d, 2H), 3.7 (m, 4H), 3.16 (s, 2H), 3.05 (m, 4H), 2.31 (s, 6H), 1.2 (s, 9H). MS (EI): 516 (MH⁺).
- [01108] N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-**
30 **yl]phenyl}-N², N²-dimethylglycinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.67 (d, 2H), 7.30 (d, 1H), 6.95 (d, 2H), 3.56-3.62 (m, 4H), 3.1 (s, 2H), 2.99-3.05 (m, 4H), 2.31 (s, 6H), 1.8-2.25 (m, 7H). MS (EI): 514 (MH⁺).
- [01109] N², N²-dimethyl-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-**
35 **yl]phenyl}amino)-pyrimidin-4-yl]phenyl}glycinamide:** ¹H NMR (400 MHz, d6-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.84 (d, 2H), 7.68 (d, 2H), 7.29 (d, 1H), 6.95 (d, 2H), 3.58-3.67 (m, 4H), 3.11 (s, 2H), 2.99-3.10 (m, 4H), 2.92 (m, 1H), 2.29 (s, 6H), 1.02 (d, 6H). MS (EI): 502 (MH⁺).

- 5 **[01110] N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycinamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.43 (d, 1H), 8.14 (d, 2H), 7.68 (d, 2H), 7.26 (d, 1H), 6.97 (d, 2H), 3.8 (m, 2H), 3.6 (m, 2H), 3.18 (m, 4H), 3.07 (m, 2H), 2.28 (s, 6H), 2.02 (m, 1H), 0.78 (m, 4H). MS (EI): 501(MH⁺).
- 10 **[01111] N-[4-(2-{{4-(4-D-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl]phenyl]-N²,N²-dimethylglycinamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.55 (d, 1H), 8.23 (d, 2H), 7.84 (d, 2H), 7.71 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 3.8 (m, 2H), 3.84 (q, 1H), 3.62 (m, 4H), 3.12 (s, 2H), 3.05 (m, 4H), 2.31 (s, 6H), 1.12 (d, 3H). MS (EI): 504(MH⁺).
- 15 **[01112] N-[4-(2-{{4-(4-L-alanyl)piperazin-1-yl}phenyl}amino)pyrimidin-4-yl]phenyl]-N²,N²-dimethylglycinamide:** ¹H NMR (400 MHz, d₆-DMSO): 10.0 (s, 1H), 9.4 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.84 (d, 2H), 7.69 (d, 2H), 7.28 (d, 1H), 6.97 (d, 2H), 3.8 (m, 2H), 3.84 (q, 1H), 3.62 (m, 4H), 3.12 (s, 2H), 3.05 (m, 4H), 2.31 (s, 6H), 1.12 (d, 3H). MS (EI): 504(MH⁺).
- 20 **[01113] 2,6-dichloro-N-(3-{{4-(4-fluorophenyl)pyrimidin-2-yl}amino}propyl)benzamide:** ¹H NMR (400 MHz, DMSO): 8.705 (t, 1H), 8.354 (d, 1H), 7.136 (br s, 2H), 7.515 (d, 2H), 7.5 (d, 2H), 7.425 (m, 1H), 7.34 (t, 2H), 7.26 (t, 1H), 7.14 (d, 1H), 3.456 (br s, 2H), 3.355 (m, 2H), 1.827 (t, 2H). MS (EI): 420.1 (MH⁺).
- [01114] N-(4-{2-[(1-[(2,6-dichlorophenyl)carbonyl]azetidin-3-yl)methyl]amino}pyrimidin-4-yl}phenyl)acetamide:** ¹H NMR (400 MHz, DMSO): 11.031 (s, 1H), 8.344 (d, 1H), 8.137 (d, 2H), 7.839 (d, 2H), 7.621 (d, 2H), 7.531 (m, 1H), 7.45 (t, 1H), 7.146 (d, 1H), 4.157 (m, 1H), 3.867 (t, 2H), 3.626-3.556 (m, 3H), 2.842 (br s, 1H), 1.725 (s, 3H). MS (EI) for C₂₃H₂₁Cl₂N₅O₂: 470.2 (MH⁺).
- 25 **[01115] N-(4-{2-[(3-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide:** ¹H NMR (400 MHz, DMSO): 10.22 (s, 1H), 9.511 (s, 1H), 8.504 (d, 1H), 8.154 (d, 2H), 7.76 (d, 3H), 7.343 (d, 1H), 7.215-7.153 (m, 2H), 6.584 (d, 1H), 3.775 (t, 4H), 3.14 (t, 4H), 2.094 (s, 3H). MS (EI) for C₂₂H₂₃N₅O₂: 390.1 (MH⁺).
- 30 **[01116] N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)cyclohexyl]-2,6-dichloro-benzamide:** ¹H NMR (400 MHz, MeOD): 8.25 (d, 1H), 8.08 (d, 2H), 7.7 (d, 2H), 7.45-7.35 (m, 3H), 7.05 (d, 1H), 4.05 (m, 2H), 2.5 (m, 1H), 2.1 (m, 3H), 2.05 (s, 3H), 1.5 (m, 1H), 1.3 (m, 3H). MS (EI) for C₂₅H₂₅Cl₂N₅O₂: 498.3 (MH⁺).
- 35

- 5 **[01117] N-{4-[2-({4-(4-methylpiperazin-1-yl)phenyl)methyl}amino)pyrimidin-4-yl]phenyl}-acetamide:** ¹H NMR (400 MHz, DMSO): 10.192 (s, 1H), 8.294 (d, 1H), 8.062 (d, 2H), 7.713 (d, 2H), 7.653 (t, 1H), 7.285 (br d, 2H), 7.09 (d, 1H), 6.953 (d, 2H), 4.485 (d, 2H), 3.3 (br s, 8H), 2.827 (s, 3H), 2.079 (s, 3H). MS (EI) for C₂₄H₂₈N₆O: 417.4 (MH⁺).
- [01118] N-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichloro-**
10 **benzamide** ¹H NMR (400 MHz, DMSO): 10.66 (s, 1H), 10.324 (s, 1H), 9.638 (s, 1H), 8.501 (d, 1H), 8.144 (d, 2H), 7.782 (m, 4H), 7.65 (d, 2H), 7.597 (d, 2H), 7.498 (m, 1H), 7.349 (d, 1H), 2.11 (s, 3H). MS (EI) for C₂₅H₁₉Cl₂N₅O₂: 492 (MH⁺).
- [01119] N-{4-[2-(piperidin-4-ylamino)pyrimidin-4-yl]phenyl}acetamide:** ¹H NMR (400
15 MHz, DMSO): 10.184 (s, 1H), 8.288 (d, 1H), 8.044 (d, 2H), 7.71 (d, 2H), 7.049 (t, 2H), 3.8 (br s, 1H), 2.962 (d, 2H), 2.077 (s, 3H), 1.838 (br d, 2H), 1.372-1.334 (m, 2H). MS (EI) for C₁₇H₂₁N₅O: 312.3 (MH⁺).
- [01120] N-{4-[2-({1-[(2,6-dichlorophenyl)carbonyl]piperidin-4-yl}amino)pyrimidin-4-yl]-phenyl}acetamide:** ¹H NMR (400 MHz, DMSO): 10.171 (s, 1H), 8.312 (d, 1H), 8.067 (d, 2H), 7.71 9d, 2H), 7.584-7.546 (m, 2H), 7.461 (t, 1H), 7.246 (d, 1H), 7.093 (d, 1H), 4.468
20 (m, 1H), 4.1 (br s, 1H), 3.25-3.05 (m, 2H), 2.077 (s, 3H), 2.05 (m, 1H), 1.915 (br s, 1H), 1.58-1.532 (m, 2H). MS (EI) for C₂₄H₂₃Cl₂N₅O₂: 485.3 (MH⁺).
- [01121] N-{4-[2-({4-[(2-hydroxyethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide:** ¹H NMR (400 MHz, DMSO): 10.211 (s, 1H), 9.43 (s, 1H), 8.455 (d, 1H), 8.12 (d, 2H), 7.754-7.69 (m, 4H), 7.292 (d, 1H), 6.93 (m, 2H), 4.865 (t, 1H), 3.97 (t,
25 2H), 3.715 (q, 2H), 2.09 (s, 3H). MS (EI) for C₂₀H₂₀N₄O₃: 365.1 (MH⁺).
- [01122] 1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylurea:** ¹H NMR (400 MHz, DMSO): 9.371 (s, 1H), 8.986 (s, 1H), 8.795 9s, 1H), 8.436 (d, 1H), 8.121 (d, 2H), 7.697 (d, 2H), 7.633 (d, 2H), 7.49 (d, 2H), 7.321-7.265 (m, 3H), 7.014-6.928 (t d, 3H), 3.758 (t, 4H), 3.063 (t, 4H). MS (EI) for C₂₇H₂₆N₆O₂: 467.3 (MH⁺).
- 30 **[01123] N-[5-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-2-(4-ethylpiperazin-1-yl)phenyl]-2,6-dichlorobenzamide:** ¹H NMR (400 MHz, DMSO): 10.224 (s, 1H), 9.623 (d, 2H), 8.6 (br s, 1H), 8.48 (d, 1H), 8.237 (d, 2H), 7.735 (d, 2H), 7.598 (d, 2H), 7.521 (m, 2H), 7.35 (d, 1H), 7.181 (d, 1H), 3.36 (br s, 4H), 2.877 (t, 4H), 2.344 (q, 2H), 2.071 (s, 3H), 1.005 (t, 3H). MS (EI) for C₃₁H₃₁Cl₂N₇O₂: 604.3 (MH⁺).
- 35 **[01124] 1-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]-3-(phenylmethyl)urea:** ¹H NMR (400 MHz, MeOD): 8.082 (t, 3H), 7.53 (d, 2H), 7.422 (m, 3H), 7.236 (m, 4H), 7.159 (m, 1H), 7.037 (d, 2H), 4.32 (s, 2H), 3.971 (d, 2H), 3.608 (d, 2H),

5 3.195 (t, 2H), 3.126 (d, 2H), 3.045 (t, 2H), 1.3 (t, 3H). MS (EI) for $C_{30}H_{33}N_7O$: 508.4 (MH^+).

[01125] N^2,N^2 -dimethyl-N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}glycinamide: 1H NMR (400 MHz, DMSO): 9.994 (s, 1H), 9.418 (s, 1H), 8.667 (m, 2H), 8.454 (d, 1H), 8.127 (d, 2H), 7.907-7.827 (m d, 3H), 7.7 (d, 2H), 7.513 (m, 1H), 7.298 (d, 1H), 6.98 (d, 2H), 3.799 (br s, 2H), 3.489 (br s, 2H), 3.179 (br s, 2H), 3.113 (br s, 4H), 2.289 (s, 6H). MS (EI) for $C_{30}H_{32}N_8O_2$: 537.4 (MH^+).

[01126] N-(3-fluoro-4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-cyclopropanecarboxamide: 1H NMR (400 MHz, DMSO): 10.671 (s, 1H), 9.452 (s, 1H), 8.47 (d, 1H), 8.045 (t, 1H), 7.758 (d, 1H), 7.656 (d, 2H), 7.46 (d, 1H), 7.12 (q, 1H), 6.932 (d, 2H), 3.749 (t, 4H), 3.052 (t, 4H), 1.813 (m, 1H), 0.847 (d, 4H). MS (EI) for $C_{24}H_{24}FN_5O_2$: 434.3(MH^+).

[01127] N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]-pyrimidin-4-yl}phenyl)cyclopropanecarboxamide: 1H NMR (400 MHz, DMSO): 7.851 (d, 1H), 7.67 (d, 2H), 7.452 (br d, 2H), 7.285 (d, 2H), 7.091 (d, 1H), 7.763 (d, 1H), 6.688 (d, 2H), 6.291 (br s, 1H), 3.668 (s, 3H), 3.561 (s, 2H), 2.946 (br s, 4H), 2.5 (br s, 4H), 1.413 (m, 1H), 0.541 (m, 2H), 0.3 (m, 2H). MS (EI) for $C_{29}H_{32}N_8O$: 509.4(MH^+).

[01128] N-[4-(2-{[4-(4-L-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-acetamide: 1H NMR (400 MHz, DMSO): 10.212 (s, 1H), 9.407 (s, 1H), 8.449 (d, 1H), 8.121 (d, 2H), 7.732 (d, 2H), 7.698 (d, 2H), 7.282 (d, 1H), 6.978 (d, 2H), 3.804 (q, 1H), 3.621 (m, 4H), 3.037 (br m, 4H), 2.091 (s, 3H), 1.864 (br s, 2H), 1.10 (d, 3H). MS (EI) for $C_{25}H_{29}N_7O_2$: 460.4(MH^+).

[01129] N-[4-(2-{[4-(4-L-prolylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-acetamide: 1H NMR (400 MHz, DMSO): 10.234 (s, 1H), 9.409 (s, 1H), 8.449 (d, 1H), 8.121 (d, 2H), 7.757 (d, 2H), 7.699 (d, 2H), 7.282 (d, 1H), 6.979 (d, 2H), 3.849 (m, 1H), 3.619 (m, 4H), 2.992 (m, 6H), 2.625 (m, 1H), 2.092 (s, 3H), 1.986 (m, 1H), 1.685-1.536 (m, 3H). MS (EI) for $C_{27}H_{31}N_7O_2$: 486.2(MH^+).

[01130] N-[4-(2-{[4-(4-D-alanylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-acetamide: 1H NMR (400 MHz, DMSO): 10.22 (s, 1H), 9.406 (s, 1H), 8.449 (d, 1H), 8.12 (d, 2H), 7.755 (d, 2H), 7.697 (d, 2H), 7.282 (d, 1H), 6.978 (d, 2H), 3.791 (q, 1H), 3.621 (br s,

5 4H), 3.081 (br d, 4H), 2.091 (s, 3H), 1.709 (br s, 2H), 1.096 (d, 3H). MS (EI) for $C_{25}H_{29}N_7O_2$: 460.4(MH⁺).

[01131] N-[4-(2-([4-(4-D-prolylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-acetamide: ¹H NMR (400 MHz, DMSO): 10.211 (s, 1H), 9.407 (s, 1H), 8.449 (d, 1H), 8.121 (d, 2H), 7.754 (d, 2H), 7.698 (d, 2H), 7.282 (d, 1H), 6.979 (d, 2H), 3.872 (t, 1H), 3.621 (m, 4H), 3.082-2.979 (m, 6H), 2.656 (m, 1H), 2.091 (s, 3H), 2.013 (m, 2H), 1.676-1.522 (m, 3H). MS (EI) for $C_{27}H_{31}N_7O_2$: 486.4(MH⁺).

[01132] N-{4-[2-([4-(2-piperazin-1-ylacetyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}acetamide: ¹H NMR (400 MHz, DMSO): 10.219 (s, 1H), 9.401 (s, 1H), 8.448 (d, 1H), 8.121 (d, 2H), 7.754 (d, 2H), 7.694 (d, 2H), 7.281 (d, 1H), 6.977 (d, 2H), 3.707 (t, 2H), 3.59 (t, 2H), 3.319 (s, 2H), 3.1 (t, 2H), 3.018 (t, 2H), 2.702 (s, 4H), 2.336 (br s, 4H), 2.090 (s, 3H). MS (EI) for $C_{28}H_{34}N_8O_2$: 515.2(MH⁺).

[01133] N-[4-(2-([4-(4-L-alanylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, DMSO): 9.938 (s, 1H), 9.417 (s, 1H), 8.457 (d, 1H), 8.135 (d, 2H), 7.885 (d, 2H), 7.693 (d, 2H), 7.305 (d, 1H), 6.976 (d, 2H), 4.436 (q, 1H), 3.991 (q, 1H), 3.878-3.761 (q q, 4H), 3.622 (br s, 4H), 3.083 (br d, 4H), 2.222 (m, 1H), 2.019 (m, 1H), 1.913 (m, 2H), 1.843 (br s, 2H). MS (EI) for $C_{28}H_{33}N_7O_3$: 516.3(MH⁺).

[01134] N-[4-(2-([4-(4-L-prolylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, DMSO): 9.945 (s, 1H), 9.419 (s, 1H), 8.458 (d, 1H), 8.135 (d, 2H), 7.887 (d, 2H), 7.696 (d, 2H), 7.306 (d, 1H), 6.978 (d, 2H), 4.452 (q, 1H), 4.011 (q, 1H), 3.861 (q, 2H), 3.633 (m, 4H), 3.084-2.968 (m, 6H), 2.62 (m, 1H), 2.191 (m, 1H), 2.002 (m, 2H), 1.897 (m, 2H), 1.691-1.544 (m, 3H). MS (EI) for $C_{30}H_{35}N_7O_3$: 542.3(MH⁺).

[01135] N-[4-(2-([4-(4-D-alanylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, DMSO): 9.945 (s, 1H), 9.418 9s, 1H), 8.457 (d, 1H), 8.135 (d, 2H), 7.887 (d, 2H), 7.694 (d, 2H), 7.305 (d, 1H), 6.976 (d, 2H), 4.452 (q, 1H), 4.028 (q, 1H), 3.878-3.768 (m, 2H), 3.633 (m, 4H), 3.083 (br d, 4H), 2.209 (m, 1H), 2.002 (m, 1H), 1.862 (m, 3H), 1.1 (d, 3H). MS (EI) for $C_{28}H_{33}N_7O_3$: 516.3(MH⁺).

[01136] N-[4-(2-([4-(4-D-prolylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-tetrahydrofuran-2-carboxamide: ¹H NMR (400 MHz, DMSO). 9.954 (s, 1H), 9.421 (s, 1H), 8.458 (d, 1H), 8.136 (d, 2H), 7.889 (d, 2H), 7.697 (d, 2H), 7.306 (d, 1H), 6.978 (d, 2H), 4.454 (q, 1H), 4.028 (q, 1H), 3.889 (m, 2H), 3.645 (m, 4H), 3.083-2.985 (m, 6H), 2.669 (m,

- 5 1H), 2.209 (m, 1H), 2.002 (m, 2H), 1.879 (m, 2H), 1.681-1.548 (m, 3H), MS (EI) for $C_{30}H_{35}N_7O_3$: 542.3(MH⁺).
- [01137] (2,6-dichlorophenyl)(4-(4-(4-methylthiophen-2-yl)pyrimidin-2-ylamino)piperidin-1-yl)methanone:** ¹H-NMR (400MHz, d6-DMSO): 8.28 (d, 1H), 7.72 (m, 1H), 7.57 (m, 2H), 7.47 (m, 1H), 7.32 (s, 1H), 7.27 (m, 1H), 7.01 (m, 1H), 4.45 (m, 1H), 4.03
- 10 (m, 1H), 3.28-3.05 (m, 3H), 2.45 (s, 3H), 2.03-1.80 (m, 2H), 1.59-1.48 (m, 2H); MS (EI): 447 (MH⁺).
- [01138] (2,6-dichlorophenyl)(4-(4-(pyridin-3-yl)pyrimidin-2-ylamino)piperidin-1-yl)-methanone:** ¹H-NMR (400MHz, d6-DMSO): 9.28 (br. s, 1H), 8.69 (m, 1H), 8.41 (m, 2H), 7.59-7.52 (m, 2H), 7.46 (m, 2H), 7.25 (d, 1H), 4.46 (m, 1H), 4.14 (m, 1H), 3.32-3.10 (m,
- 15 3H), 2.06-1.89 (m, 2H), 1.63-1.54 (m, 3H); MS (EI): 428 (MH⁺).
- [01139] (2,6-dichlorophenyl)(4-(4-(5-methylthiophen-2-yl)pyrimidin-2-ylamino)piperidin-1-yl)methanone:** ¹H-NMR (400MHz, d6-DMSO): 8.24 (d, 1H), 7.70 (m, 1H), 7.57 (m, 2H), 7.47 (m, 1H), 7.24 (m, 1H), 7.00 (m, 1H), 6.88 (m, 1H), 4.45 (m, 1H), 4.02 (m, 1H), 3.28-3.05 (m, 3H), 2.47 (s, 3H), 2.03-1.80 (m, 2H), 1.57-1.50 (m, 2H); MS
- 20 (EI): 447 (MH⁺).
- [01140] 1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1H-pyrrole-2-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 9.99 (s, 1H), 9.39 (s, 1H), 8.45 (d, 1H), 8.13 (d, 2H), 7.90 (d, 2H), 7.68 (d, 2H), 7.30 (d, 1H), 7.09-7.08 (m, 1H), 7.05 (t, 1H), 6.97 (d, 2H), 6.13-6.11 (m, 1H), 3.90 (s, 3H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for
- 25 $C_{26}H_{26}N_6O_2$: 455 (MH⁺).
- [01141] 3-fluoro-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.96 (s, 1H), 9.43 (s, 1H), 8.79 (s, 1H), 8.62 (d, 1H), 8.47 (d, 1H), 8.20 (d, 2H), 7.88 (d, 2H), 7.76-7.67 (m, 2H), 7.32 (d, 1H), 6.94 (d, 2H), 6.56 (s, 1H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for $C_{26}H_{23}FN_6O_2$:
- 30 471 (MH⁺).
- [01142] 6-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.61 (s, 1H), 9.41 (s, 1H), 9.03 (d, 1H), 8.47 (d, 1H), 8.23 (dd, 1H), 8.19 (d, 2H), 7.95 (d, 2H), 7.68 (d, 2H), 7.45 (d, 1H), 7.31 (d, 1H), 6.94 (d, 2H), 3.74 (t, 4H), 3.05 (t, 4H), 2.57 (s, 3H). MS (EI) for
- 35 $C_{27}H_{26}N_6O_2$: 467 (MH⁺).
- [01143] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridazine-4-carboxamide:** ¹H NMR (400 MHz, d6-DMSO): 10.98 (s, 1H), 9.67 (s, 1H), 9.52 (d, 1H),

5 9.42 (s, 1H), 8.47 (d, 1H), 8.21 (d, 2H), 8.16-8.14 (m, 1H), 7.96 (d, 2H), 7.68 (d, 2H), 7.33 (d, 2H), 7.33 (d, 1H), 6.95 (d, 2H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for $C_{25}H_{23}N_7O_2$: 454 (MH⁺).

[01144] 2-cyclopropyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide: ¹H NMR (400 MHz, d₆-DMSO): 10.09 (s, 1H), 9.45 (s, 1H), 8.45 (d, 1H), 8.12 (d, 2H), 7.78 (d, 2H), 7.69 (d, 2H), 7.30 (d, 1H), 7.00 (s, 2H), 3.76 (s, 4H), 3.09 (s, 4H), 2.25 (d, 2H), 1.12-1.02 (m, 1H), 0.50-0.48 (m, 2H), 0.22-0.20 (m, 2H). MS (EI) for $C_{25}H_{27}N_5O_2$: 430 (MH⁺).

[01145] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)isoxazole-5-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.99 (s, 1H), 9.42 (s, 1H), 8.47 (d, 1H), 8.19 (d, 2H), 7.95 (d, 2H), 7.68 (d, 2H), 7.33-7.31 (m, 2H), 6.95 (d, 2H), 6.55 (s, 1H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for $C_{24}H_{22}N_6O_3$: 443 (MH⁺).

[01146] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.69 (s, 1H), 9.41 (s, 1H), 9.14 (s, 1H), 8.79 (d, 1H), 8.47 (d, 1H), 8.34-8.31 (m, 1H), 8.20 (d, 2H), 7.97 (d, 2H), 7.70 (d, 2H), 7.67-7.58 (m, 1H), 7.33 (d, 1H), 6.95 (d, 2H), 3.78 (t, 4H), 3.05 (t, 4H). MS (EI) for $C_{26}H_{24}N_6O_2$: 453 (MH⁺).

[01147] 4-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-benzamide: ¹H NMR (400 MHz, d₆-DMSO): 10.42 (s, 1H), 9.53 (s, 1H), 8.47 (s, 1H), 8.17 (d, 2H), 7.98 (d, 2H), 7.91 (d, 2H), 7.72 (s, 2H), 7.37 (d, 3H), 7.05 (s, 2H), 3.78 (s, 4H), 3.14 (s, 4H), 2.40 (s, 3H). MS (EI) for $C_{28}H_{27}N_5O_2$: 466 (MH⁺).

[01148] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-prolinamide: ¹H NMR (400 MHz, d₆-DMSO): 11.55 (s, 1H), 11.15 (s, 1H), 10.16 (s, 2H), 8.74 (s, 1H), 8.52 (s, 1H), 8.23 (d, 2H), 7.89 (d, 2H), 7.67 (d, 2H), 7.48 (s, 1H), 7.11 (d, 2H), 4.50 (s, br, 1H), 3.81 (d, 2H), 3.57 (d, 2H), 3.28-3.11 (m, 8H), 2.05-1.92 (m, 3H), 1.30 (t, 3H). MS (EI) for $C_{27}H_{33}N_7O$: 472 (MH⁺).

[01149] N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-butanamide: ¹H NMR (400 MHz, d₆-DMSO): 10.25 (s, 1H), 9.37 (s, 1H), 8.43 (s, 1H), 8.12 (d, 2H), 7.77 (d, 2H), 7.68 (d, 2H), 7.29 (s, 1H), 6.93 (d, 2H), 3.08 (s, 4H), 2.42-2.30 (m, 4H), 1.68-1.58 (m, 2H), 1.05 (t, 3H), 0.93 (t, 3H). MS (EI) for $C_{26}H_{32}N_6O$: 445 (MH⁺).

[01150] 1-ethyl-3-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]urea: ¹H NMR (400 MHz, d₆-DMSO): 9.32 (s, 1H), 8.85 (s, 1H), 8.40 (d, 1H), 8.05 (d, 2H), 7.68 (d, 2H), 7.54 (d, 2H), 7.23 (d, 1H), 6.92 (d, 2H), 6.36 (t, 1H), 3.18-3.05 (m,

5 6H), 2.54 (t, 4H), 2.46-2.38 (m, 2H), 1.09-1.02 (m, 6H). MS (EI) for C₂₅H₃₁N₇O: 446 (MH⁺).

[01151] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)furan-3-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.17 (s, 1H), 9.46 (s, 1H), 8.47-8.43 (m, 2H), 8.18 (d, 2H), 7.91 (d, 2H), 7.83 (d, 1H), 7.70 (s, 2H), 7.32 (s, 1H), 7.03-6.95 (m, 3H),
10 3.76 (s, 4H), 3.09 (s, 4H). MS (EI) for C₂₅H₂₃N₅O₃: 442 (MH⁺).

[01152] N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1,3-thiazole-4-carboxamide: ¹H NMR (400 MHz, d₆-DMSO): 10.61 (s, 1H), 9.40 (s, 1H), 9.30 (d, 1H), 8.56 (d, 1H), 8.46 (d, 1H), 8.16 (d, 2H), 8.06 (d, 2H), 7.69 (d, 2H), 7.32 (d, 1H), 6.96 (d, 2H), 6.56 (s, 1H), 3.74 (t, 4H), 3.05 (t, 4H). MS (EI) for C₂₄H₂₂N₆O₂S: 459 (MH⁺).

15 [01153] Based on the synthetic examples described hereinabove, the skilled artisan would be able to make the remainder of the compounds intended to be within the scope of the invention described in the appended claims.

Biological Assays

Assay Example 1

Measurement of JAK-2 Kinase Activity by ATP Hydrolysis

[01154] JAK-2 kinase activity was measured by monitoring peptide substrate dependent hydrolysis of ATP via quantitation of remaining ATP with luciferase based chemiluminescence. For compound evaluation, 0.5 µl of the compound dissolved in DMSO
25 was added to 10 µl of JAK-2 dissolved in assay buffer (20 mM HEPES pH 7.5, 10 mM MgCl₂, 0.03% Triton and 1mM DTT). After preincubation for 30 minutes at room temperature, the reaction was initiated by addition of 10 µl of ATP and the substrate peptide poly-Glu-Tyr in assay buffer. Final enzyme, ATP, and peptide concentrations were 3 nM, 1 µM, and 2 µM, respectively. After incubation for 60 minutes at room temperature, reaction
30 progress was quantitated by addition of 10 µl Kinase-Glo (Promega) and measurement of chemiluminescence in a Victor reader (Perkin Elmer). A reaction in which compound was omitted was used to determine maximum reaction progress. Omission of compound and enzyme from the reaction was used to determine zero reaction progress.

5

Assay Example 2*Measurement of JAK-3 Kinase Activity by ATP Hydrolysis*

[01155] JAK-3 was assayed similarly as JAK-2 (see Assay Example 1) except that the enzyme reaction was carried out for 180 minutes and enzyme, ATP, and peptide
10 concentrations were 30 nM, 2 μ M, and 4 μ M, respectively.

Biological Activity

[01156] Compounds in Table 1 were determined to have inhibitory activity for JAK-2 of less than 10 μ M. Other more preferred compounds of the invention have inhibitory activity
15 for JAK-2 of less than 100 nm. One of ordinary skill in the art can use the disclosures herein as well as what is known in the art to test the inhibitory activity of a particular compound.

Pharmaceutical Composition Examples

[01157] The following are representative pharmaceutical formulations containing a
20 compound of Formula I.

Tablet Formulation

[01158] The following ingredients are mixed intimately and pressed into single scored tablets.

Ingredient	Quantity per tablet, mg
compound of this invention	400
Cornstarch	50
croscarmellose sodium	25
Lactose	120
magnesium stearate	5

5

Capsule Formulation

[01159] The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

Ingredient	Quantity per tablet, mg
compound of this invention	200
lactose, spray-dried	148
magnesium stearate	2

10

Suspension Formulation

[01160] The following ingredients are mixed to form a suspension for oral administration.

Ingredient	Amount
compound of this invention	1.0 g
fumaric acid	0.5 g
sodium chloride	2.0 g
methyl paraben	0.15 g
propyl paraben	0.05 g
granulated sugar	25.5 g
sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
Flavoring	0.035 mL
Colorings	0.5 mg
distilled water	q.s. to 100 mL

Injectable Formulation

15 [01161] The following ingredients are mixed to form an injectable formulation.

Ingredient	Amount
compound of this invention	1.2 g
sodium acetate buffer solution	0.4 M 2.0 mL

Ingredient	Amount
HCl (1 N) or NaOH (1 M)	q.s. to suitable pH
water (distilled, sterile)	q.s.to 20 mL

5

[01162] All of the above ingredients, except water, are combined and heated to 60° - 70° with stirring. A sufficient quantity of water at 60° C. is then added with vigorous stirring to emulsify the ingredients, and water then added q.s. to 100 g.

10

Suppository Formulation

[01163] A suppository of total weight 2.5 g is prepared by mixing the compound of the invention with Witepsol® H-15 (triglycerides of saturated vegetable fatty acid; Riches-Nelson, Inc., New York), and has the following composition:


Ingredient	Quantity per tablet, mg
compound of this invention	500
Witepsol® H-15	balance

15 [01164] The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications can be made while remaining within the spirit and scope of the invention. It will be obvious to one of skill in the art that
20 changes and modifications can be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive.

[01165] The scope of the invention should, therefore, be determined not with reference to
25 the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

5 What is claimed is:

1. A compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein

10 D is hydrogen, halo, -CF₃, heterocycloalkyl or alkyl;

E is hydrogen, halo, -CF₃, heterocycloalkyl or alkyl; or

D and E, together with the carbon atoms to which they are attached, form a 5-7 membered heteroaryl or a 5-7 membered heterocycloalkyl, wherein the 5-7 membered heteroaryl or 5-7 membered heterocycloalkyl are each fused to the pyrimidinyl moiety to which D and E are attached;

15 L is a bond, -O- or -N(H)-;


Z is selected from alkoxy, cycloalkyl, heteroaryl optionally substituted with alkyl, halo, -C(O)OR²⁶, -C(=N-OH)alkyl, -C(O)R⁸, -C(O)NR³⁰R^{30a}, -CH₂R², -(CH₂)_{n5}NR²⁶R^{26a}, -CF₃, -CN, -SO₂R¹², -S-R^{12a}, -OR^{32a}, -NHC(O)R³², aryl, and heterocycloalkyl optionally substituted with 1 or 2 oxo, or

20 Z and R²⁵, together with the carbon atoms to which they are attached, join to form a 5 or 6 membered heterocycloalkyl, a 5 or 6 membered heteroaryl, or a 5 or 6 membered cycloalkyl ring, wherein the 5 or 6 membered heterocycloalkyl, 5 or 6 membered heteroaryl, and 5 or 6 membered cycloalkyl ring are fused to the phenyl moiety to which Z and R²⁵ are attached, and wherein the 5 or 6 membered heterocycloalkyl, 5 or 6 membered heteroaryl, or 5 or 6 membered cycloalkyl ring are each optionally substituted with 1, 2, or 3 groups independently selected from oxo, alkyl, alkoxy and halo;

n₁ is 0, 1, 2, 3, or 4, and each n₁ is independently selected when more than one n₁ is present;


30 n₂ is 0, 1, 2, 3, or 4, and each n₂ is independently selected when more than one n₂ is present;

- 5 n_3 is 0, 1, 2, or 3, and each n_3 is independently selected when more than one n_3 is present;
 n_4 is 0, 1, 2, 3 or 4, and each n_4 is independently selected when more than one n_4 is present;
 n_5 is 0, 1, 2, 3 or 4, and each n_5 is independently selected when more than one n_5 is present;
 p is 0-3;
 r is 1-3;
- 10 R^1 is hydrogen;
 R^2 is selected from one of the following groups:



5

or R^2 is selected from one of the following groups:



ring X in formula (d) of R^2 is a 5 or 6 membered unsaturated heterocyclic ring fused to the two carbon atoms of the phenyl moiety to which ring X is attached, wherein ring X contains 1 or 2 nitrogen atoms;

- 10 R^7 , R^7 , R^9 , R^{10} , R^{12} and R^{15} are each independently hydrogen, alkyl, alkoxy, or alkoxyalkyl; R^8 is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxylamino, hydroxyalkyl, alkoxyalkyl, dihydroxyalkyl, alkylamino, dialkylamino, aminoalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, alkylaminoalkyl, dialkylaminoalkyl, $-(\text{CH}_2)_r-\text{C}(\text{O})\text{OR}^7$, $-(\text{CH}_2)_r-\text{C}(\text{O})\text{NR}^7\text{R}^7$, aryl,
- 15 heteroaryl, cycloalkyl, arylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with one, two, three, four or five groups independently selected from alkyl, alkenyl,
- 20 lower alkynyl, halo, hydroxy, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl;
- each R^{11} , when R^{11} is present, is independently selected from alkyl, alkenyl, lower alkynyl, $-\text{CF}_3$, alkoxy, halo, haloalkoxy, haloalkyl, aminoalkyl, aminoalkoxy, alkylaminoalkyl, alkylaminoalkoxy, dialkylaminoalkyl, dialkylaminoalkoxy, oxo, thioalkyl,
- 25 alkylthioalkyl, $-(\text{CH}_2)_p-\text{OR}^{17}$, $-\text{CN}$, $-\text{O}-\text{CH}_2-\text{C}(\text{O})-\text{R}^{17}$, $-\text{C}(\text{O})\text{R}^{16}$, $-(\text{CH}_2)_p-\text{C}(\text{O})\text{OR}^{17}$, $-\text{S}(\text{O})_2\text{R}^{17}$, $-\text{S}(\text{O})_2\text{NR}^{15}\text{R}^{17}$, aryl, heteroaryl, cycloalkyl, arylalkyl, arylalkoxy, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally
- 30 substituted at any ring position with 1, 2, 3 or 4 R^{21} ;

5 R^{12} is hydrogen or alkyl;

R^{12a} is hydrogen or alkyl;

R^{13} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxylamino,

haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl,
aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, $-(CH_2)_r-$

10 $C(O)OR^7$,

$-(CH_2)_r-C(O)NR^7R^{7'}$, aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl,
heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein

the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl,
cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently

15 optionally substituted at the ring position with 1, 2, 3, 4 or 5 groups independently

selected from alkyl, alkenyl, lower alkynyl, halo, hydroxy, hydroxyalkyl,

alkoxycarbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl,

alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl; and wherein the alkyl

of cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, and heteroarylalkyl are

20 independently optionally substituted with 1, 2, 3, 4, or 5 groups selected from halo and

hydroxy;

R^{14} is a bond, heterocycloalkyl or cycloalkyl;

R^{16} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxyamino,

haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl,

25 aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl,

dialkylaminoalkyl,

$-(CH_2)_r-C(O)OR^7$, aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl,

heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein

the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl,

30 cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently

optionally substituted at the ring position with one, two, three, four or five groups

independently selected from alkyl, alkenyl, lower alkynyl, halo, hydroxy, hydroxyalkyl,

alkoxycarbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl,

alkylamino, dialkylamino, heterocyclalkoxy, oxo and haloalkyl; and wherein the alkyl

35 of cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, and heteroarylalkyl is optionally

substituted with 1, 2, 3, 4, or 5 groups selected from halo and hydroxy;

- 5 R^{17} is selected from hydrogen, hydroxy, alkyl, alkenyl, lower alkynyl, hydroxyamino, haloalkyl, alkyl substituted with halo and hydroxy, hydroxyalkyl, alkoxyalkyl, aminocarbonylalkyl, alkylaminocarbonylalkyl, dialkylaminocarbonylalkyl, dialkylaminoalkyl, $-(CH_2)_r-C(O)OR^7$, $-(CH_2)_r-C(O)NR^7R^7$, aryl, heteroaryl, cycloalkyl, arylalkyl, 10 diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, heteroaryl, cycloalkyl, arylalkyl, diarylalkyl, aryloxyalkyl, heteroarylalkyl, cycloalkylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at the ring position with one, two, three, four or five groups independently selected from alkyl, alkenyl, 15 lower alkynyl, halo, hydroxy, hydroxyalkyl, alkoxy, carbonyl, alkylcarbonyl, haloalkyl, haloalkoxy, lower alkoxy, amino, aryl, alkylamino, dialkylamino, heterocyclylalkoxy, oxo and haloalkyl; and wherein the alkyl of cycloalkylalkyl, heterocycloalkylalkyl, arylalkyl, and heteroarylalkyl is optionally substituted with 1, 2, 3, 4, or 5 groups selected from halo and hydroxy;
- 20 each R^{21} , when R^{21} is present, is independently selected from alkyl, alkenyl, lower alkynyl, cyano, halo, haloalkoxy, haloalkyl, hydroxyalkyl, amino, alkylamino, dialkylamino, dialkylaminoalkyl, dialkylaminoalkoxy, haloalkyl, oxo, $-OR^{13}$, $-NHS(O)_2R^{17}$, $-S(O)_2R^{17}$, $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)NR^{15}R^{17}$, $-NR^{15}C(O)R^{17}$, aryl, arylalkyl, heteroarylalkyl, aryloxy, and heteroaryl; wherein each of the aryl, arylalkyl, 25 heteroarylalkyl, aryloxy, and heteroaryl within R^{21} are optionally substituted at any ring position with 1, 2, or 3 groups selected from alkyl, lower alkoxy halo, phenyl, heteroaryl and alkylheteroalkyl;
- R^{25} is selected from alkyl, alkenyl, lower alkyl, halo, haloalkyl, haloalkoxy, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, $-OR^{12}$, cyano, 30 $-CH_2NHC(O)OR^7$, $-CH_2NHC(O)R^7$, $-SR^7$, $-S(O)_2R^7$, $-S(O)_2NR^7R^8$, $-C(O)OR^8$, $-C(O)NR^7R^8$, cycloalkyl, heterocycloalkyl, aryl and heteroaryl; wherein the cycloalkyl, heterocycloalkyl, aryl and heteroaryl are each optionally substituted with one, two or three groups independently selected from alkyl, alkenyl, halo, haloalkoxy, haloalkyl, amino, alkylamino, dialkylamino, aminoalkyl, 35 alkylaminoalkyl, $-OR^8$, $-NHS(O)_2R^8$, cyano, $-C(O)R^8$, $-CH_2NHC(O)OR^7$, $-CH_2NHC(O)R^7$, $-SR^7$, $-S(O)_2R^7$, $-S(O)_2NR^7R^8$, $-C(O)OR^8$, $-C(O)NR^7R^8$, $-NR^7C(O)-CHR^3-OR^8$, $-NR^7C(O)-CHR^3-NR^7-R^8$, and $-NR^7C(O)R^8$;

- 5 R^{26} is hydrogen, -C(O)-phenyl or alkyl, wherein the -C(O)-phenyl is optionally substituted at any ring position with 1, 2 or 3 halo;
- R^{26a} is hydrogen, alkyl, heteroaryl, -C(O) R^{32} , -C(O)NHR^{32a}, -S(O)₂R⁹, -SR⁹, -C(O)OR³², or -C(O)NR^{32a}R³²;
- 10 R^{27} and R^{28} are each independently selected from alkyl, alkenyl, hydroxy, alkoxy, and alkoxyalkyl;
- R^{27a} and R^{28a} are independently selected from hydrogen, alkyl, alkenyl, alkoxyalkyl, alkoxycarbonylalkyl, hydroxyalkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, dialkylaminoalkyl, arylcarbonylalkyl, aryloxyalkyl, dialkylaminoalkyl, alkyl-O-C(O)heterocycloalkyl, -(CH₂)_{n4}heterocycloalkyl, heterocycloalkylalkyl, heteroaryl,
- 15 heteroarylalkyl, -(CH₂)_{n4}-C(O) R^{29} , -(CH₂)_{n4}NR^{28a}R^{28a}, -(CH₂)_{n4}NHR^{28a}, -CH(phenyl)₂, -S(O)₂R²⁹, -C(O) R^{29} , -C(O)OR²⁹, and -C(O)NR^{29a}R²⁹, wherein the aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl groups within R^{27a} and R^{28a} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl,
- 20 alkoxy, alkylcarbonyl, phenyl, phenoxy, arylcarbonyl, -CF₃, oxo, -OCF₃, alkoxyphenyl, and heteroaryl optionally substituted with alkyl or halo;
- or R^{27} and R^{27a} , together with the nitrogen to which they are attached, form heterocycloalkylamino, heterocycloalkyl or heteroaryl, wherein the heterocycloalkylamino and heteroaryl are each independently optionally substituted
- 25 with 1, 2, 3, 4, or 5 R^{31} ;
- or R^{28} and R^{28a} together with the nitrogen to which they are attached form heterocycloalkyl or heteroaryl, wherein the heterocycloalkyl and heteroaryl are each optionally substituted with 1, 2, 3, 4, or 5 R^{31} ;
- R^{29a} is hydrogen or alkyl;
- 30 R^{29} is selected from alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl groups within R^{29} are each optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, alkoxy, alkylcarbonyl,
- 35 phenyl, phenoxy, arylcarbonyl, -CF₃, oxo, -OCF₃, alkoxyphenyl, and heteroaryl optionally substituted with alkyl or halo;
- R^{30a} is hydrogen or alkyl;

5 R^{30} is selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, alkoxyalkoxyalkyl, alkoxycarbonylalkyl, amino, alkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, aryl, arylalkyl, phenoxyalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, arylheteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the aryl, arylalkyl, phenoxyalkyl, cycloalkyl, arylheteroarylalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl groups within R^{30} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, alkoxy, alkoxyalkyl, $-C(O)OCH_3$, $-CF_3$, $-OCF_3$, alkylcarbonyl, phenyl, phenoxy, alkylphenoxy, dialkylaminoalkoxy and heteroaryl;


15 R^{31} is selected from alkyl, hydroxyalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, $-C(O)R^{30}$, $-C(O)NR^{30}R^{30a}$, $-C(O)OR^{30}$, $-S(O)_2R^{30}$, amino, dihydroxyalkyl, arylcarbonyl, alkylcarbonylamino, alkoxyphenyl, phenylalkoxyalkyl, arylheteroarylalkyl, alkylamino, $-O$ -dialkylamino, dialkylamino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, dialkylaminoalkoxy, oxo, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, spirocyclic cycloalkyl, spirocyclic heterocycloalkyl, and heterocycloalkylalkyl, wherein the aryl, arylalkyl, cycloalkyl, arylheteroarylalkyl, arylalkoxyalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl groups within R^{31} are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from halo, alkyl, $-CF_3$, $-OCF_3$, cyano, alkoxy, alkoxyalkyl, $-C(O)OCH_3$, alkylcarbonyl, phenyl optionally substituted at any ring position with halo, phenoxy, alkylphenoxy, arylalkoxyalkyl, dialkylaminoalkoxy and heteroaryl;

R^{32a} is hydrogen, $-OCF_3$, $-CF_3$, or alkyl;


30 R^{32} is selected from aryl, arylalkyl, arylalkoxy, arylcycloalkyl, alkoxycarbonylalkoxy, cycloalkyl, cycloalkylalkyl, cycloalkylhydroxyalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl, wherein the aryl, arylalkyl, cycloalkyl, arylcycloalkyl, cycloalkylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from hydroxy, oxo, alkyl, alkoxy, amino, hydroxyalkyl, alkylcarbonyl, alkoxycarbonyl, halo, $-CF_3$, $-OCF_3$, aminoalkyl, alkylaminoalkyl, aryl and dialkylaminoalkyl, and wherein the alkyl portion of the heteroarylalkyl can be substituted with amino;

- 5 or R^{32} is alkyl optionally substituted with 1, 2, 3, 4, or 5 groups independently selected from hydroxy, alkoxycarbonyl, alkoxy, $-\text{CF}_3$, halo, aminocarbonyl, alkylaminocarbonyl, alkoxycarbonylalkylamino, dialkylaminocarbonyl, $-\text{NR}^{34}\text{R}^{34a}$ and phenyl optionally substituted with 1, 2, or 3 halo;
- or R^{32} is alkylamino or arylalkylamino;
- 10 R^{34} is hydrogen or alkyl;
- R^{34a} is selected from hydrogen, alkyl, heteroaryl, aryl, aminoalkyl, aminocarbonylalkyl, heteroarylalkyl, arylalkoxy and arylalkyloxycarbonylalkyl; wherein the heteroaryl, aryl, heteroarylalkyl, arylalkoxy or arylalkyloxycarbonylalkyl are each independently optionally substituted at any ring position with 1, 2, 3, 4, or 5 groups selected from
- 15 hydroxy, oxo, alkyl, amino, hydroxyalkyl, alkylcarbonyl, alkoxycarbonyl, halo, aminoalkyl, alkylaminoalkyl, and dialkylaminoalkyl; and
- R^{35} is selected from halo, $-(\text{CH}_2)_p\text{C}(\text{O})\text{OR}_{17}$, cycloalkyl, heterocycloalkyl, and heterocycloalkylalkyl; wherein the heterocycloalkyl and heterocycloalkylalkyl are each optionally substituted with 1, 2, 3, 4, or 5 groups each independently selected from
- 20 alkyl, alkoxy, and halo.


2. The compound according to claim 1 having Formula II:




3. The compound according to claim 1 having Formula III:



- 5 4. The compound according to claim 1, wherein R^2 is




5. The compound according to claim 1, wherein R^2 is



10


wherein R^{28a} is arylalkyl or heteroarylalkyl, wherein the arylalkyl or heteroarylalkyl are each optionally substituted with 1, 2, 3, 4, or 5 substituents selected from halo and lower alkyl.

6. The compound according to claim 1, wherein R^2 is



15


7. The compound according to claim 1, wherein R^2 is




wherein R^{28a} is selected from lower alkyl, dialkylaminoalkyl, alkoxyalkyl, arylalkyl, heteroarylalkyl, and heterocycloalkylalkyl.

20


8. The compound according to claim 1, wherein R^2 is




- 5 wherein R^{11} and n_2 are as defined above for the compound of Formula I, and R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, join together to form a ring structure selected from thiazolidinyl, piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyrimidinyl, and pyridinyl, wherein the ring structure is optionally substituted with 1, 2, 3, 4 or 5 substituents selected from halo, lower alkyl or alkoxy.
- 10 9. The compound according to claim 1, wherein R^2 is



10. The compound according to claim 1, wherein L is a bond, and Z is




11. The compound according to claim 1, wherein L is a bond, Z is



- 15 hydrogen. and R^{25} is


12. The compound according to claim 1, wherein L is a bond, Z is



- hydrogen and E and D are hydrogen.


13. The compound according to claim 1, wherein R^{25} is on the 3 position.

14. The compound according to claim 1, wherein L is a bond, Z is




- C(O) R^{32} , and R^{26a} is


15. The compound according to claim 1, wherein L is a bond, Z is




- C(O) R^{32} , and R^{32} is selected from lower alkyl, cycloalkyl, diaminoalkyl, aminoalkyl, arylalkyl, heterocycloalkyl, alkoxyalkyl, alkylamino, and hydroxyalkyl optionally substituted with amino.




- 5 16. The compound according to claim 1, wherein L is a bond, Z is $-\text{C}(\text{O})\text{R}^{32}$, and R^{32} is cycloalkyl.




17. The compound according to claim 1, wherein L is a bond, Z is $-\text{C}(\text{O})\text{R}^{32}$, and R^{32} is lower alkyl.



- 10 18 The compound according to claim 1, wherein L is a bond, Z is $-\text{C}(\text{O})\text{R}^{32}$, R^{26} is hydrogen, wherein R^{32} selected from aryl, arylalkyl, cycloalkyl, alkoxy carbonylalkyl, heteroaryl, heteroarylalkyl, heterocycloalkyl, and heterocycloalkylalkyl, wherein R^{32} optionally substituted with 1, 2, 3, 4 or 5 groups selected from hydroxyl, oxo, alkyl, alkoxy, amino, hydroxyalkyl and halo.




- 15 19. The compound according to claim 1, wherein L is a bond, Z is $-\text{C}(\text{O})\text{R}^{32}$, R^{26} is hydrogen, wherein R^{32} is lower alkyl optionally substituted with 1, 2, 3, 4 or 5 groups selected from dialkylaminocarbonyl, hydroxy and $-\text{NR}^{34}\text{R}^{34a}$.



- 20 20. The compound according to claim 1, wherein Z is $-\text{C}(\text{O})\text{R}^{32}$, R^{26} is hydrogen, and R^{32} selected from tetrahydrofuran, pyrrolidinyl or pyrimidinyl, wherein R^{32} optionally substituted with 1, 2, 3, 4 or 5 groups selected from hydroxyl, oxo, alkyl, alkoxy, amino, hydroxyalkyl and halo.

- 25 21. The compound according to claim 1, wherein R^2 is



22. The compound according to claim 1, wherein R^{32} is methyl.

5

23 The compound according to claim 1, wherein R^{32} is alkyl substituted with $-NR^{34}R^{34a}$.

24. The compound according to claim 3, wherein R^{32} is methyl.

10

25. The compound according to claim 2, wherein R^{32} is methyl.

15

26. The compound according to claim 1, wherein R^{32} is U or $-\text{CH}_2\text{-U}$, and wherein U is selected from pyrrolidinyl, thiazolidinyl, morpholinyl, azetidiny, cyclobutyl, cyclopropyl, tetrahydrofuranyl, pyrazinyl, imidazolyl, piperazinyl, thienyl, thienylmethyl, furanyl, phenyl, prolinamidyl, pyridinyl, tetrahydronaphthalene, tetrazolyl, isoindolinyl, pyranyl, cyclopentyl, and octahydro-1H-indolyl.

20

27. The compound according to claim 1, wherein R^{11} , when present, is halo or lower alkyl.


28. The compound according to claim 1, wherein R^{11} , when present, is lower alkyl.

25


29. The compound according to claim 1, wherein R^{35} is heterocycloalkylalkyl, wherein the heterocycloalkyl is selected from piperazinyl, piperidinyl, morpholinyl and dioxanyl.

30. The compound according to claim 1, wherein n_2 is 0.


31. The compound according to claim 1, wherein R^2 is



30

32. The compound according to claim 1, wherein R^2 is , and wherein R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl.

33. The compound according to claim 1 having Formula IV:




wherein R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or two R^{31} .

34. The compound according to claim 33, wherein D, E and R^{25} are each hydrogen.

35. The compound according to claim 33, wherein R^{32} is heterocycloalkyl.

36. The compound according to claim 33, wherein R^{32} is alkyl optionally substituted with alkoxy, hydroxy, amino, alkylamino, or dialkylamino.

37. The compound according to claim 1 having Formula V:



wherein R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or two R^{31} .


38. The compound according to claim 37, wherein D, E and R^{25} are each hydrogen.

39. The compound according to claim 37, wherein R^{32} is heterocycloalkyl.

40. The compound according to claim 37, wherein R^{32} is alkyl optionally substituted with alkoxy, hydroxy, amino, alkylamino, or dialkylamino.

5

41. The compound according to claim 1 having Formula VI:



10 wherein R^{28} and R^{28a} , together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or two R^{31} .


42. The compound according to claim 41, wherein D, E and R^{25} are each hydrogen.








15 43. The compound according to claim 41, wherein R^{32} is heterocycloalkyl.

44. The compound according to claim 41, wherein R^{32} is alkyl optionally substituted with alkoxy, hydroxy, amino, alkylamino, or dialkylamino.

20 45. A compound according to claim 1 selected from








N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide










2,6-dichloro- <i>N</i> -(3-([4-(2,4-dichlorophenyl)pyrimidin-2-yl]amino)propyl)benzamide	
2,6-dichloro- <i>N</i> -[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]benzamide	
2,6-dichloro- <i>N</i> -(3-([4-(2,3-dihydro-1-benzofuran-6-yl)pyrimidin-2-yl]amino)propyl)benzamide	
<i>N</i> -[3-([4-(4-(acetylamino)phenyl)pyrimidin-2-yl]amino)propyl]-2-fluoro-6-iodobenzamide	
<i>N</i> -(3-([4-(4-aminophenyl)pyrimidin-2-yl]amino)propyl)-2,6-dichlorobenzamide	
<i>N</i> -[4-([4-(4-(acetylamino)phenyl)pyrimidin-2-yl]amino)phenyl]-2,6-dichlorobenzamide	
<i>N</i> -{4-[2-([3-([4-ethylpiperazin-1-yl]carbonyl)phenyl]amino)pyrimidin-4-yl]phenyl}acetamide	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-[2-(dimethylamino)ethyl]benzamide	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluorobenzamide	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-fluoro-6-iodobenzamide	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyridine-4-carboxamide	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,3,4,5,6-pentafluorobenzamide	
4-(4-chlorophenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide	

<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
4-(2,4-dichlorophenyl)- <i>N</i> -{3-[(2-piperidin-1-ylethyl)oxy]phenyl}pyrimidin-2-amine	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chlorobenzamide	
<i>N</i> -(4-{2-[(3-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-{2-[(3-piperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-bromobenzamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3-fluorobenzamide	








<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide</p>	
<p><i>N</i>-(4-{2-[(3-{[(2,6-dichlorophenyl)sulfonyl]amino}phenyl)amino]-5-methylpyrimidin-4-yl}phenyl)acetamide</p>	
<p>2,6-dichloro-<i>N</i>-(3-{[4-(1<i>H</i>-indol-5-yl)pyrimidin-2-yl]amino}phenyl)benzamide</p>	
<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]-5-fluoropyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide</p>	
<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-methylbenzamide</p>	
<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,4-dichlorobenzamide</p>	
<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,3-dichlorobenzamide</p>	








<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,5-dichlorobenzamide	
<i>N</i> -[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -(4-{2-[(4-piperidin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide	
<i>N</i> -(4-{2-[(2-methyl-4-piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide	
<i>N</i> -(4-{5-methyl-2-[(3-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide	
<i>N</i> -(4-{6-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide	
<i>N</i> -[3-({4-[4-(acetylamino)-2-chlorophenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide	








<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-difluorobenzamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,4,5-trifluorobenzamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]benzamide	
<i>N</i> -(4-{6-morpholin-4-yl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3,5-difluorobenzamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-fluoro-3-(methoxy)benzamide	
<i>N</i> -[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2-chloro-6-fluoro-4-methylbenzamide	








<p><i>N</i>-(4-{2-[(3-{[(2,6-dimethylphenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>4-(2,4-dichlorophenyl)-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p>4-(2,4-dichlorophenyl)-<i>N</i>-{3-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}pyrimidin-2-amine</p>	
<p><i>N</i>-(3-{[4-(4-aminophenyl)pyrimidin-2-yl]amino}phenyl)-2,6-dichlorobenzamide</p>	
<p>4-(4-aminophenyl)-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p>4-[4-(ethylamino)phenyl]-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p><i>N</i>-[5-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-2-morpholin-4-ylphenyl]-2,6-dichlorobenzamide</p>	








<i>N</i> -(4-{5-fluoro-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{[2-(4-ethylpiperazin-1-yl)-2-oxoethyl]oxy}phenyl)amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -[4-(2-{[3-(morpholin-4-ylcarbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -(3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-5-[(4-ethylpiperazin-1-yl)carbonyl]phenyl)-2,6-dichlorobenzamide	
4-[4-(dimethylamino)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
2,6-dichloro- <i>N</i> -(3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)phenyl)benzamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)phenyl)acetamide	

<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-1-methylpiperidine-4-carboxamide</p>	
<p><i>N</i>-{4-[2-({3-[(phenylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-(2-ethylphenyl)benzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-(phenylmethyl)benzamide</p>	
<p><i>N</i>-{4-[2-({3-[(4-cyclopentylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-{4-[2-({3-[(4-phenylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-(3-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	








<p><i>N</i>-(2-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenylacetamide</p>	
<p><i>N</i>-{4-[2-({3-[(4-pyrazin-2-yl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenylacetamide</p>	
<p><i>N</i>-(4-{2-[(3-{[4-(3-chlorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>3-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-[(1-methyl-1<i>H</i>-benzimidazol-2-yl)methyl]benzamide</p>	
<p>3-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-propylbenzamide</p>	
<p>3-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-cyclopropylbenzamide</p>	
<p>3-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-[(3-fluorophenyl)methyl]benzamide</p>	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(naphthalen-1-ylmethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(dimethylamino)ethyl]- <i>N</i> -methylbenzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(2-methylphenyl)methyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(3-chlorophenyl)methyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(2-phenylethyl)benzamide	
<i>N</i> -{4-[2-({3-[(4-methylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(tetrahydrofuran-2-ylmethyl)benzamide	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[3-(2-oxopyrrolidin-1-yl)propyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(3 <i>s</i> ,5 <i>s</i> ,7 <i>s</i>)-tricyclo[3.3.1.1~3,7~]dec-1-yl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(methyloxy)ethyl]benzamide	
<i>N</i> -[4-(2-({[3-(1,3-thiazolidin-3-yl)carbonyl]phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -{4-[2-({3-[(4-pyridin-2-yl)piperazin-1-yl]carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[2-(methyloxy)phenyl]methyl}benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[3-(methyloxy)phenyl]methyl}benzamide	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(2-fluorophenyl)methyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(4-fluorophenyl)methyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(3,3-dimethylbutyl)benzamide	
<i>N</i> -[4-(2-{[3-(thiomorpholin-4-ylcarbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(2-thienylmethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[3-(dimethylamino)propyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[{2-(trifluoromethyl)phenyl}methyl]benzamide	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[3-(trifluoromethyl)phenyl]methyl}benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[4-(trifluoromethyl)phenyl]methyl}benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[2,4-difluorophenyl]methyl}benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -ethyl- <i>N</i> -methylbenzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[4-((trifluoromethyl)oxy)phenyl]methyl}benzamide	
<i>N</i> -{4-[2-({3-[(4-acetylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(cyclopropylmethyl)benzamide	

3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[2-(2-fluorophenyl)ethyl]benzamide	
<i>N</i> -[4-(2-({3-(pyrrolidin-1-ylcarbonyl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -{4-[2-({3-[(4-pyrimidin-2-yl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide	
<i>N</i> -methyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)benzamide	
<i>N</i> -[4-(2-({3-(1,3-dioxan-2-yl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -[4-(2-({3-(morpholin-4-ylmethyl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	








<p><i>N</i>-{4-[2-({3-[(4-ethylpiperazin-1-yl)methyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-[3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)phenyl]-3-[(2-morpholin-4-ylethyl)oxy]benzamide</p>	
<p>4-[4-(methylamino)phenyl]-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p><i>N</i>-[4-(2-({4-[4-(acetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[(3-({4-(2-fluorophenyl)piperazin-1-yl}carbonyl)phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-[2-(phenyloxy)ethyl]benzamide</p>	
<p>methyl 1-{{3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)phenyl}carbonyl}piperidine-4-carboxylate</p>	

<p><i>N</i>-[4-(2-{[3-({4-[3-(methyloxy)phenyl]piperazin-1-yl}carbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-{2-[2-(methyloxy)phenyl]ethyl}benzamide</p>	
<p><i>N</i>-[4-(2-{[3-(1,3-dihydro-2<i>H</i>-isoindol-2-ylcarbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-(biphenyl-4-ylmethyl)benzamide</p>	
<p><i>N</i>-(4-{2-([3-({4-(phenylcarbonyl)piperazin-1-yl}carbonyl)phenyl]amino}pyrimidin-4-yl)phenyl}acetamide</p>	
<p><i>N</i>-[4-(2-{[3-({4-[4-(methyloxy)phenyl]piperazin-1-yl}carbonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>3-({4-[4-(acetilamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-methyl-<i>N</i>-{[2-(methyloxy)phenyl]methyl}benzamide</p>	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -[(2-fluorophenyl)methyl]- <i>N</i> -methylbenzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -(diphenylmethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -(2-pyridin-2-ylethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -(pyridin-2-ylmethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl} amino)- <i>N</i> -[2-(2-chlorophenyl)ethyl]benzamide	
<i>N</i> -[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-fluoropyrimidin-4-yl)phenyl]acetamide	
<i>N</i> ² -[3-(1 <i>H</i> -imidazol-1-yl)propyl]- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	








<p><i>N</i>-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-<i>N</i>²-(2-pyridin-3-ylethyl)glycinamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-(pyridin-3-ylmethyl)benzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-(pyridin-4-ylmethyl)benzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-methyl-<i>N</i>-(phenylmethyl)benzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-cyclopentylbenzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-[(2-chlorophenyl)methyl]benzamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-[(4-chlorophenyl)methyl]benzamide</p>	








3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -(furan-2-ylmethyl)benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -{[4-(methyloxy)phenyl]methyl}benzamide	
<i>N</i> -[4-(2-{{3-({4-[2-(methyloxy)phenyl]piperazin-1-yl}carbonyl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[3-(methyloxy)propyl]benzamide	
<i>N</i> -(4-{2-[(3-{{(2 <i>R</i> ,6 <i>S</i>)-2,6-dimethylmorpholin-4-yl}carbonyl)phenyl}amino]pyrimidin-4-yl}phenyl)acetamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -[(6-chloropyridin-3-yl)methyl]benzamide	
3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)- <i>N</i> -butylbenzamide	








<p><i>N</i>-(4-{2-[(3-{[4-(2-chlorophenyl)piperazin-1-yl]carbonyl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-<i>N</i>-ethyl-<i>N</i>-[2-(methyloxy)ethyl]benzamide</p>	
<p><i>N</i>-(4-(2-(3-(3-morpholinopropoxy)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p><i>N</i>-(4-(2-(3-(2-(dimethylamino)ethoxy)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p><i>N</i>-[3-({4-[4-(acetylamino)phenyl]-5-methylpyrimidin-2-yl}amino)phenyl]-2,6-dimethylbenzamide</p>	
<p><i>N</i>-[4-(2-{[4-(phenyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>4-(4-aminophenyl)-<i>N</i>-[4-(phenyloxy)phenyl]pyrimidin-2-amine</p>	

<i>N</i> -{4-[2-({4-[(phenylmethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
4-(4-aminophenyl)- <i>N</i> -[3-(morpholin-4-ylsulfonyl)phenyl]pyrimidin-2-amine	
<i>N</i> -(4-{2-[(3,5-dimorpholin-4-yl)phenyl]amino}-5-fluoropyrimidin-4-yl)phenylacetamide	
<i>N</i> -{4-[2-({4-[4-(phenylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
<i>N</i> -(4-{2-[(4-{4-[(5-methyl-3-phenylisoxazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{4-[(5-methyl-1-phenyl-1H-pyrazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{4-[(2-phenyl-1,3-thiazol-4-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	

<p><i>N</i>-[4-(2-{{4-(4-{{6-(phenyloxy)pyridin-3-yl}methyl)piperazin-1-yl}phenyl)amino}pyrimidin-4-yl}phenyl]acetamide</p>	
<p><i>N</i>-(4-[2-{{4-[4-(cyclohexylmethyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl]phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[[4-{4-[(1<i>S</i>,4<i>S</i>)-bicyclo[2.2.1]hept-5-en-2-ylmethyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-[4-(2-{{4-(4-pentylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[[4-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-[4-(2-{{4-(4-{{3,5-bis(methyloxy)phenyl}methyl)piperazin-1-yl}phenyl)amino}pyrimidin-4-yl}phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[[4-{4-[(4-fluorophenyl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	








<p><i>N</i>-(4-{2-[(4-{4-[(1-methyl-1<i>H</i>-pyrrol-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(2,4-dichlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-(9<i>H</i>-fluoren-2-ylmethyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(3-methyl-2-thienyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(5-ethylfuran-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(3-{[4-(1,1-dimethylethyl)phenyl]oxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-(3-thienylmethyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	








methyl 4-({4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazin-1-yl}methyl)benzoate	
<i>N</i> -(4-{2-[(4-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{4-[(4-{[3-(dimethylamino)propyl]oxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -[4-(2-{[4-(4-{2-[(phenylmethyl)oxy]ethyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -(4-{2-[(4-{4-[(2-chloroquinolin-3-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{4-[(4-chloro-2,6-dimethylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -{1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]pyrrolidin-3-yl}acetamide	








<i>N</i> ² -[3-(4-methylpiperazin-1-yl)propyl]- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	
<i>N</i> ² -(1-methylpiperidin-4-yl)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	
<i>N</i> -(4-{2-[(4-{(pyridin-4-yl)methyl}oxy)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-{2-[(4-{[2-(methyloxy)ethyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
2-(dimethylamino)- <i>N</i> -(4-{2-(4-morpholinophenylamino)pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)furan-2-carboxamide	
2-(methyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	

<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclobutanecarboxamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-3-carboxamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-2-carboxamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-3-carboxamide	
<i>N</i> -[4-(2-{[4-(dimethylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -(4-{2-[(4-chlorophenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(3-{[(2-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	

<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)piperidine-4-carboxamide	
2-amino- <i>N</i> -(4-{2-(4-morpholinophenylamino)pyrimidin-4-yl}phenyl)propanamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)morpholine-2-carboxamide	
<i>N</i> ² -methyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-beta-alaninamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)phenylalaninamide	








<p><i>N</i>-[4-(2-{[4-(3-oxopiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-[4-(2-{[4-(4-{[5-(3-chlorophenyl)furan-2-yl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-[4-(2-{[4-(4-{[4-fluoro-2-(trifluoromethyl)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-[4-(2-{[4-(4-{[4-(1<i>H</i>-imidazol-1-yl)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-[4-(2-{[4-(4-{[2,5-bis(trifluoromethyl)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-(2,6-dimethylphenyl)methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-(2,3-dimethylphenyl)methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	








<p><i>N</i>-[4-(2-([4-(4-{[2,4-bis(ethyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-[4-(2-([4-(4-{[3-(ethyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-{4-[2-([4-[4-(2,2'-bithien-5-ylmethyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-[4-(2-([4-(4-{[4-(2-thienyl)phenyl]methyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(4-cyanophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-[4-(2-([4-(4-{[2,5-bis(methyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-{4-[2-([4-[4-(2,2-diphenylethyl)piperazin-1-yl]phenyl]amino)pyrimidin-4-yl]phenyl}acetamide</p>	








<p><i>N</i>-{4-[2-({4-[4-(1<i>H</i>-pyrrol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-{4-[2-(1<i>H</i>-indazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-{4-[2-(1<i>H</i>-indol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-[4-(2-{[4-(morpholin-4-ylmethyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-(4-{2-[(3-{[(3-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-{2-[(3-{[(4-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>4-[4-({4-[4-(acetamidino)phenyl]pyrimidin-2-yl}amino)phenyl]-<i>N</i>-ethylpiperazine-1-carboxamide</p>	

<i>N</i> -{4-[2-({4-[4-(ethylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
<i>N</i> -{4-[2-(1 <i>H</i> -indazol-5-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide	
<i>N</i> -[4-(2-({4-(4-propylpiperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -[4-(2-({4-(4-butylpiperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -{4-[2-({4-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
4-[4-(methylsulfonyl)phenyl]- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
ethyl <i>N</i> -[4-({4-[4-(acetlamino)phenyl]pyrimidin-2-yl}amino)phenyl]- <i>N</i> -methylglycinate	








4-[3-(methylsulfonyl)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
4-[4-(methylthio)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
N-(4-{2-[(4-cyclohexylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
N-{4-[2-({4-[(tetrahydrofuran-2-ylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
N-{4-[2-({4-[(phenylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
N-[4-(2-{[4-(acetylamino)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
methyl (4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)carbamate	








1-ethyl-3-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)urea	
ethyl 1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-3-carboxylate	
ethyl [4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]acetate	
4-[4-(methyloxy)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
4-[3-(methyloxy)phenyl]-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
4-(1H-indol-5-yl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
2-[(2-amino-2-oxoethyl)amino]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	








2-morpholin-4-yl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
2,6-dichloro- <i>N</i> -{3-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}benzamide	
<i>N</i> ² -(2-aminoethyl)- <i>N</i> ² -methyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)glycinamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)- <i>N</i> ² -1 <i>H</i> -pyrazol-5-ylglycinamide	
phenylmethyl <i>N</i> -{2-[(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]amino}-2-oxoethyl}- <i>L</i> -alaninate	
4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)benzamide	
1,1-dimethylethyl [(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)methyl]carbamate	








<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylacetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylpropanamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-2-carboxamide	
5-methyl- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrazine-2-carboxamide	
2-(ethyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-(phenyloxy)acetamide	








<i>N</i> -[4-(2-{[4-(1 <i>H</i> -pyrrol-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -[4-(2-{[4-(2,6-dimethylmorpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
ethyl 1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-4-carboxylate	
2-cyclopentyl- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-3-pyridin-3-ylpropanamide	
6-(methyloxy)- <i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)pyridine-3-carboxamide	
methyl 4-[(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl]amino]-4-oxobutanoate	








<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide	
<i>N</i> -(4-{2-[(4-bis[2-(methoxy)ethyl]amino)phenyl]amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -[4-(2-{[4-(morpholin-4-ylsulfonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
4-(4-(aminomethyl)phenyl)- <i>N</i> -(4-morpholinophenyl)pyrimidin-2-amine	
<i>N</i> -[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)methyl]acetamide	
<i>N</i> -(4-morpholin-4-ylphenyl)-4-{4-[(propylamino)methyl]phenyl}pyrimidin-2-amine	
<i>N</i> -(4-{2-[(4-piperidin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	








<i>N</i> -(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
2-(2-methylphenyl)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)cyclopentanecarboxamide	
<i>N,N</i> -dimethyl- <i>N'</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanediamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)- <i>N</i> ² -pyrimidin-4-ylglycinamide	
3-chloro- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-4-carboxamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-piperidin-1-ylacetamide	








<i>N</i> ² -ethyl- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)glycinamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyrrolidin-1-ylacetamide	
2-(1 <i>H</i> -imidazol-1-yl)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-piperazin-1-ylacetamide	
<i>N</i> -[4-(2-{[4-(4-phenylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
<i>N</i> -(4-{2-[(3-chloro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-[(4-piperazin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide	

'N-[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyridin-2-yl]-2,6-dichlorobenzamide	
'N-[6-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)pyrimidin-4-yl]-2,6-dichlorobenzamide	
'5-fluoro-N ⁴ -[2-(methyloxy)phenyl]-N ² -[3-(methyloxy)phenyl]pyrimidine-2,4-diamine	
'2,6-dichloro-N-{3-[(4-{[3-chloro-4-(methyloxy)phenyl]oxy}pyrimidin-2-yl)amino]phenyl}benzamide	
'4-{[2-chloro-4-(methyloxy)phenyl]oxy}-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
'N-[4-({2-[(4-morpholin-4-ylphenyl)amino]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}amino)phenyl]acetamide	
'N-[4-({2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}oxy)phenyl]acetamide	








<p><i>N</i>-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-{4-[2-({4-[(2<i>R</i>,6<i>S</i>)-2,6-dimethylmorpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-(4-{2-[(4-{4-[(2-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-{4-[2-({4-[4-(1<i>H</i>-pyrazol-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p><i>N</i>-(4-{2-[(3-piperazin-1-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-[4-(2-{[3-(4-{[2-(methyloxy)phenyl]methyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p><i>N</i>-{4-[2-({3-[4-(1,3-thiazol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	

<p><i>N</i>-(4-{2-[(3-bromo-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-[4-(2-{[4-{[2-(diethylamino)ethyl]oxy}-3-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzoic acid</p>	
<p>4-(4-furan-2-ylphenyl)-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p>4-[4-(3-methyl-1,2,4-oxadiazol-5-yl)phenyl]-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p>4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzonitrile</p>	
<p>methyl 4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzoate</p>	

4-(4-fluorophenyl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine	
<i>N</i> -[3-({2-[(4-morpholin-4-ylphenyl)amino]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl} amino)phenyl]acetamide	
<i>N</i> -(4-morpholin-4-ylphenyl)-4-[4-(pyridin-3-ylamino)phenyl]pyrimidin-2-amine	
<i>N</i> -(4-morpholin-4-ylphenyl)-4-[4-(pyridin-2-ylamino)phenyl]pyrimidin-2-amine	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)methanesulfonamide	
1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-(phenylmethyl)urea	
4-(2,3-dihydro-1,4-benzodioxin-6-yl)- <i>N</i> -(4-morpholin-4-ylphenyl)pyrimidin-2-amine	








<p><i>N</i>-(4-{2-[(4-morpholin-4-ylphenyl)amino]-7<i>H</i>-pyrrolo[2,3-<i>d</i>]pyrimidin-4-yl}phenyl)acetamide</p>	
<p><i>N</i>-(4-morpholin-4-ylphenyl)-4-quinolin-6-ylpyrimidin-2-amine</p>	
<p>4-[4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl]-<i>N</i>-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	
<p><i>N</i>-(4-morpholin-4-ylphenyl)-4-(4-pyrimidin-5-ylphenyl)pyrimidin-2-amine</p>	
<p><i>N</i>-(4-morpholin-4-ylphenyl)-4-quinoxalin-6-ylpyrimidin-2-amine</p>	
<p>2-chloro-<i>N</i>-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide</p>	
<p>2-(2-fluorophenyl)-<i>N</i>-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	








<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrimidine-5-carboxamide	
(2 <i>S</i>)- <i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-2-carboxamide	
<i>N</i> -(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)- <i>N</i> ² -phenylglycinamide	
<i>N</i> -(4-{2-[(4-ethylpiperazin-1-yl)carbonyl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
<i>N</i> -(4-{2-(3-methoxy-4-morpholino-phenylamino)pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)-pyrimidin-4-yl}phenyl)acetamide	
<i>N</i> -(4-{2-(4-(4-(3-methylbutanoyl)piperazin-1-yl)phenylamino)-pyrimidin-4-yl}phenyl)acetamide	








<i>N</i> -(4-(2-(4-(4-(cyclopropanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(cyclopentanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(2-methoxybenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-pentanoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-picolinoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-isonicotinoylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	








<i>N</i> -(4-(2-(4-(4-(1-acetylpiperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(2-cyclopropylacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(2-(2-methoxyethoxy)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(2-(pyridin-3-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-(4-(3-(pyridin-3-yl)propanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide	








<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyridin-3-yl)acetamide	
<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)isonicotinamide	
<i>N</i> -(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-prolinamide	
<i>N</i> -[4-(2-{[3-(methyloxy)-4-morpholin-4-yl]phenyl}amino)pyrimidin-4-yl)phenyl]-D-prolinamide	
O-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-serinamide	
(<i>S</i>)-3-hydroxy- <i>N</i> -(4-(2-(4-morpholino-phenylamino)-pyrimidin-4-yl)-phenyl)-butanamide	
(<i>R</i>)-3-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide	








<i>(R)</i> -2-amino-3-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide	
2-Hydroxy-2-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide	
2-methyl- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
<i>(R)</i> - <i>N</i> -(4-(2-(4-((<i>R</i>)-3-(dimethylamino)pyrrolidin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
4-amino-1,1-dioxo- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydro-2 <i>H</i> -thiopyran-4-carboxamide	
<i>(R)</i> -4-(4-aminophenyl)- <i>N</i> -(4-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)-pyrimidin-2-amine	
<i>(R)</i> - <i>N</i> -(4-(2-(4-(3-(dimethylamino)pyrrolidin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxy-propanamide	








<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperazine-2-carboxamide	
2-amino- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1,2,3,4-tetrahydronaphthalene-2-carboxamide	
4-(4-(1,1-dioxo-isothiazolidin-2-yl)phenyl)- <i>N</i> -(4-morpholinophenyl)-pyrimidin-2-amine	
4-(4-(1 <i>H</i> -tetrazol-1-yl)phenyl)- <i>N</i> -(4-morpholinophenyl)-pyrimidin-2-amine	
(<i>R</i>)- <i>N</i> -(4-(2-(3-(benzyloxy)-4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide	
(<i>S</i>)-2-amino-3-hydroxy- <i>N</i> -(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide	
<i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(1 <i>H</i> -tetrazol-1-yl)acetamide	







<p>(R)-N-(4-(2-(3-ethoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(1,2,3,4-tetrahydroquinolin-6-ylamino)pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide</p>	
<p>3-hydroxy-N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-3-methylbutanamide</p>	
<p>(3S,7S)-7-(hydroxymethyl)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)quinuclidine-3-carboxamide</p>	
<p>1-hydroxy-N-(4-(2-(4-morpholino-phenylamino)-pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p>(S)-2-amino-N-(4-(2-(3-methyl-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(R)-N-(4-(2-(3-methyl-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	








<p>(R)-N-(4-(2-(4-morpholino-3-(trifluoromethyl)-phenylamino)pyrimidin-4-yl)-phenyl)-pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(4-((S)-tetrahydrofuran-2-carbonyl)-piperazin-1-yl)-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(4-((R)-tetrahydrofuran-2-carbonyl)-piperazin-1-yl)-phenylamino)-pyrimidin-4-yl)phenyl)-pyrrolidine-2-carboxamide</p>	
<p>4-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperazine-1-carboxamide</p>	
<p>3-methoxy-N-(4-(2-(4-morpholino-3-(trifluoromethyl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>3-methoxy-N-(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-propane-1-sulfonamide</p>	
<p>2-methoxy-N-(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-ethanesulfonamide</p>	

<p>)-3-hydroxy-<i>N</i>-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-phenyl)butanamide</p>	
<p>1-(3-hydroxy-<i>N</i>-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-phenyl)butanamide</p>	
<p>4-(2-(4-morpholinophenylamino)pyrimidin-4-phenyl)-2,5-dihydro-1<i>H</i>-pyrrole-2-carboxamide</p>	
<p>1-(dimethylamino)propyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-phenyl)urea</p>	
<p>1-(4-(2-(4-(4-(<i>S</i>)-pyrrolidin-2-ethyl)piperazin-1-yl)phenylamino)pyrimidin-4-phenyl)pyrrolidine-2-carboxamide</p>	
<p>1-amino-<i>N</i>-(4-(2-(4-(4-ethylpiperazin-1-enylamino)-5-methylpyrimidin-4-enyl)propanamide</p>	
<p>1-methoxypropyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-phenyl)urea</p>	








<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-ethylpiperazin-1-yl)phenylamino)-5-methylpyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)-3-fluorophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(3-chloro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>1-(2-morpholinoethyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea</p>	
<p>1-(2-(dimethylamino)ethyl)-3-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(pyrrolidin-2-yl)acetamide</p>	
<p>2,3-dihydroxy-<i>N</i>-(4-(2-(4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)-propanamide</p>	

<p>(<i>S</i>)-2-amino-4-methyl-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide</p>	
<p>(<i>R</i>)-2-amino-4-methyl-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide</p>	
<p><i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)isoindoline-1-carboxamide</p>	
<p><i>N</i>-ethyl-4-(4-(4-(4-(tetrahydrofuran-2-carboxamido)phenyl)pyrimidin-2-ylamino)phenyl)piperazine-1-carboxamide</p>	
<p><i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-((<i>R</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-((<i>S</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	








<p><i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide</p>	
<p>3-methoxy-<i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p><i>N</i>-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1-methylpyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-4-(4-(4-(4-(2-aminopropanamido)phenyl)-pyrimidin-2-ylamino)phenyl)-<i>N</i>-ethylpiperazine-1-carboxamide</p>	








<p>(<i>R</i>)-2-amino-<i>N</i>-(4-(2-(4-(4-((<i>R</i>)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(<i>R</i>)-2-amino-<i>N</i>-(4-(2-(4-(4-((<i>S</i>)-pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(<i>R</i>)-2-amino-<i>N</i>-(4-(2-(4-(4-((<i>S</i>)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p><i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	








1-ethyl-3-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)urea	
(S)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
(R)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide	
(S)-3-methoxy-N-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
(R)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
N-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
N-(4-(2-(4-(4-(isobutyryl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	








<p><i>N</i>-(4-(2-(4-(1-butyl-1,2,4-triazin-4-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	
<p>1-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea</p>	
<p><i>N</i>-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide</p>	
<p><i>N</i>-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p><i>N</i>-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	
<p>1-ethyl-3-(4-(2-(4-(4-pivaloylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea</p>	
<p>1-(4-(2-(4-(4-(cyclobutanecarbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea</p>	

1-ethyl-3-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea	
N-ethyl-4-(4-(4-(4-(3-ethylureido)phenyl)pyrimidin-2-ylamino)phenyl)piperazine-1-carboxamide	
(S)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(R)-1-(2-hydroxyethyl)-N-(4-(2-(4-morpholinobenzyl)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(R)-1-isopropyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(S)-2-(dimethylamino)-N-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
1-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea	

(R)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
4-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)tetrahydro-2H-pyran-4-carboxamide	
(R)-2-amino-N-(4-(2-(4-(4-isobutyrylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
(R)-2-amino-N-(4-(2-(4-(4-((R)-2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
(R)-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(R)-2-amino-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
(R)-2-(dimethylamino)-N-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	

<p>(R)-N-(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(S)-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-2-amino-N-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(2R)-N-(4-(2-(4-(4-(tetrahydrofuran-3-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(S)-1-ethyl-3-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea</p>	
<p>(S)-1-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea</p>	
<p>N-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	

<p>(S)-N-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	
<p>3-methoxy-N-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(R)-2-amino-N-(4-(5-methyl-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>2-(dimethylamino)-N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>1-ethyl-3-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea</p>	
<p>3-methoxy-N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(R)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide</p>	

<p>(S)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-5-oxopyrrolidine-2-carboxamide</p>	
<p>(S)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide</p>	
<p>(2R,3S)-2-amino-3-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide</p>	
<p>(R)-2-amino-N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p>N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	
<p>N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide</p>	








<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
(<i>R</i>)-2-amino- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
(2 <i>S</i> ,3 <i>R</i>)-2-amino-3-hydroxy- <i>N</i> -(4-(2-(4-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-((2 <i>S</i> ,6 <i>R</i>)-2,6-dimethylmorpholino)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(3-hydroxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
1-ethyl-3-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea	
(<i>R</i>)-1-ethyl-3-(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea	








3,3,3-trifluoro-2-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide	
<i>(R)</i> -1-(4-(2-(4-(4-(2-aminopropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-ethylurea	
2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
<i>(R)</i> - <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
<i>(R)</i> -2-amino- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	
<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)-3-methoxypropanamide	








<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide	
<i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide	
<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide	
2-(dimethylamino)- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
3-methoxy- <i>N</i> -(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide	








1-ethyl-3-(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)urea	
<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropanoyl)piperazin-1-yl)benzyl)pyrimidin-4-yl)phenyl)acetamide	
(<i>S</i>)- <i>N</i> -(4-(2-(4-(4-(2-(dimethylamino)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide	
(4 <i>S</i>)-4-hydroxy- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(<i>R</i>)- <i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide	
(<i>S</i>)- <i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide	








<i>N</i> -(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
<i>N</i> -(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide	
(<i>S</i>)- <i>N</i> -(4-(2-(4-(4-(pyrrolidine-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)butyramide	
(2 <i>R</i>)- <i>N</i> -(4-(2-(4-(4-(tetrahydrofuran-2-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(<i>R</i>)- <i>N</i> -(4-(5-chloro-2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
<i>N</i> -(4-(2-(4-(4-(3-(diethylamino)propanoyl)piperazin-1-yl)benzyl)pyrimidin-4-yl)phenyl)acetamide	
(<i>S</i>)-1-(2-hydroxyethyl)- <i>N</i> -(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide	








<p>(S)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanediamide</p>	
<p>(R)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanediamide</p>	
<p>(R)-2-amino-N1-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)succinamide</p>	
<p>(R)-N-(4-(2-(4-(4-((4-chloro-1-methyl-1H-pyrazol-3-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(S)-1-ethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(4-(2-ethoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(4-(2-(pyrrolidin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	








<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-(2-morpholinoacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p><i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1<i>H</i>-imidazole-4-carboxamide</p>	
<p>2-(dimethylamino)-<i>N</i>-(4-(2-(4-(4-(2-(piperazin-1-yl)acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(4-(4-(3-(dimethylamino)-2,2-dimethylpropyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-2-carboxamide</p>	
<p>(<i>R</i>)-2-hydroxy-2-methyl-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide</p>	
<p>(<i>S</i>)-2-hydroxy-2-methyl-<i>N</i>-(4-(2-(4-morpholinobenzyl)pyrimidin-4-yl)phenyl)butanamide</p>	
<p>(<i>R</i>)-2-methoxy-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	








<p>(S)-2-methoxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(R)-N-(4-(2-(4-(4-(2-methoxyacetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(4-acetyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(S)-2-amino-N-(4-(5-methyl-2-(4-(4-((1-methyl-1H-imidazol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>N-(4-(2-(4-(4-(piperidine-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)tetrahydrofuran-3-carboxamide</p>	
<p>(2R,4S)-4-hydroxy-N-(4-(2-(4-(4-(3-methoxypropanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopentanecarboxamide</p>	








<p>(R)-N-(4-(2-(4-(4-formylpiperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-1-(2-hydroxyethyl)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-3-carboxamide</p>	
<p>1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	
<p>N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1H-pyrrole-2-carboxamide</p>	
<p>N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-1H-imidazole-2-carboxamide</p>	
<p>(S)-2-hydroxy-3,3-dimethyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide</p>	
<p>(R)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	








(S)-2-cyclohexyl-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide	
(S)-2-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide	
1-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclobutanecarboxamide	
(R)-N-(4-(2-(6-morpholinopyridin-3-ylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(S)-N-(4-(2-(3-chloro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide	
(2R,3R)-2-amino-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide	
1-hydroxy-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)cyclopentanecarboxamide	






<p>(R)-N-(4-(2-(4-(4-(4-(dimethylamino)butanoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-N-(4-(2-(4-(2-methoxyethyl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-7-ylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(R)-2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)butanamide</p>	
<p>(R)-2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide</p>	
<p>(R)-2-amino-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)hexanamide</p>	
<p>(R)-2-amino-3-methoxy-N-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(2S,3R)-2-amino-3-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pentanamide</p>	






<p>(<i>R</i>)-<i>N</i>-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p><i>N</i>-(4-(2-(3-methoxy-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)-2-(1<i>H</i>-tetrazol-1-yl)acetamide</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)indoline-2-carboxamide</p>	
<p>(<i>R</i>)-<i>tert</i>-butyl 2-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenylcarbamoyl)pyrrolidine-1-carboxylate</p>	
<p>1-acetyl-4-amino-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)piperidine-4-carboxamide</p>	
<p>(<i>R</i>)-2-amino-3-methoxy-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>(<i>S</i>)-<i>N</i>-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	






<p>(<i>R</i>)-2-amino-<i>N</i>-(4-(2-(3-fluoro-4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide</p>	
<p>2-hydroxy-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-(2-hydroxyethyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-((1-methyl-1<i>H</i>-pyrrol-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(<i>R</i>)-<i>N</i>-(4-(2-(4-(4-((<i>R</i>)-pyrrolidin-2-yl)methyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)pyrrolidine-2-carboxamide</p>	
<p>(2<i>S</i>,3<i>aS</i>,7<i>aS</i>)-<i>N</i>-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)octahydro-1<i>H</i>-indole-2-carboxamide</p>	
<p><i>N</i>-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)cyclopropanecarboxamide</p>	






N-(4-{5-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)cyclopropanecarboxamide	
N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)valinamide	
N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl)acetamide	
N-(4-{2-[(3,5-dimorpholin-4-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl)acetamide	
N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-alaninamide	
N-{4-[2-(4-{4-(2-methylpropanoyl)piperazin-1-yl}phenyl)amino}pyrimidin-4-yl]phenyl}acetamide	
2-amino-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-2-phenylacetamide	






N-(4-{5-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
3-(methyloxy)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)propanamide	
N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)prolinamide	
N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-alaninamide	
N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl)phenyl)acetamide	
N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl)phenyl)acetamide	
N-(4-{2-[(2-methyl-4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	





N-[4-(2-[(4-pyrrolidin-1-yl)phenyl]amino)pyrimidin-4-yl]phenyl]acetamide	
N-[4-(2-[[4-(diethylamino)phenyl]amino]pyrimidin-4-yl)phenyl]acetamide	
N-[4-(2-[(4-azepan-1-yl)phenyl]amino)pyrimidin-4-yl]phenyl]acetamide	
N-{4-[2-({4-[methyl(2-phenylethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
N-[4-(2-{{4-(1,4-dioxo-8-azaspiro[4.5]dec-8-yl)phenyl}amino}pyrimidin-4-yl)phenyl]acetamide	






<p>N-[4-(2-{[4-(2-oxopiperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-{[4-(2-methylpiperidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-valinamide</p>	<p>Chiral</p> 
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-valinamide</p>	<p>Chiral</p> 
<p>2-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)alaninamide</p>	






<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tryptophanamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1,2,3,4-tetrahydroisoquinoline-1-carboxamide</p>	
<p>O-(1,1-dimethylethyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide</p>	<p>Chiral</p> 
<p>3-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-carboxamide</p>	
<p>bis(1,1-dimethylethyl) (2R)-2-[[4-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]carbonyl]piperazine-1,4-dicarboxylate</p>	<p>Chiral</p> 





<p>N-(4-{2-[(4-{4-[2-(2-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[2-(2-methylphenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[2-(3-fluorophenyl)acetyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-[(4-{4-(3-thienylcarbonyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-(4-{2-[(4-{4-[(6-chloropyridin-3-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	






<p>N-(4-{2-[(4-{4-[(3-methylfuran-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-(2-(4-(4-(3-fluoro-2-methylbenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-(4-(2-(4-(4-(1H-imidazole-4-carbonyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-(4-(2-(4-(4-(2-methoxynicotinoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-(4-(2-(4-(4-(4-fluoro-3-methylbenzoyl)piperazin-1-yl)phenylamino)pyrimidin-4-yl)phenyl)acetamide</p>	







<p>N-{4-[2-({4-[4-(naphthalen-2-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({4-[4-(quinolin-8-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-[4-(2-{[4-(4-{[4-(1,1-dimethylethyl)phenyl]sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-{[4-(4-{[5-bromo-2-(methyloxy)phenyl]sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	






<p>N-(4-{2-[(4-{4-[(phenylmethyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-[4-(2-{[4-(4-{3-(trifluoromethyl)phenyl)sulfonyl]piperazin-1-yl}phenyl)amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-(4-{2-[(4-{4-[(2-methylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(3-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(2,4-difluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	





<p>N-{4-[2-({3-[4-({4-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)amino)pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-(4-{2-[(3-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-({3-[4-({2-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)amino)pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-(4-{2-[(3-{4-[(3-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-({3-[4-(2,3-dihydroxypropyl)piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	





<p>N-{4-[2-({3-[4-(1,3-benzodioxol-5-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(pyridin-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(pyridin-3-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(1H-pyrrol-2-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	





<p>4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-(phenylmethyl)piperazine-1-carboxamide</p>	
<p>N-[4-(2-({3-(4-({2-(methyloxy)phenyl}carbonyl)piperazin-1-yl)phenyl}amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-{4-[2-({3-[4-(1H-pyrazol-4-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(3-pyridin-3-ylpropanoyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-(4-{2-[(3-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	





<p>N-[4-(2-([3-(4-{2-[(4-fluorophenyl)oxy]acetyl)piperazin-1-yl]phenyl)amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-{4-[2-([3-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-([3-[4-(pyridin-4-ylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-([3-[4-(pyridin-2-ylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-{2-([3-{4-[(2-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-([3-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}acetamide</p>	





<p>N-{4-[2-({3-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-[4-(2-({3-(methyloxy)-4-morpholin-4-ylphenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide</p>	
<p>(2R)-N-[4-(2-({3-(methyloxy)-4-morpholin-4-ylphenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	
<p>(2S)-N-[4-(2-({3-(methyloxy)-4-morpholin-4-ylphenyl}amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	





<p>N-(4-{2-[(4-{(2-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(3-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>ethyl 3-[(4-{2-[(4-morpholin-4-yl)phenyl]amino]pyrimidin-4-yl}phenyl)amino]-3-oxopropanoate</p>	
<p>N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide</p>	





<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide</p>	
<p>N-ethyl-4-{4-[(4-{4-[(tetrahydrofuran-3-ylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}piperazine-1-carboxamide</p>	
<p>N-[4-(2-{[4-(4-D-alanyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide</p>	
<p>N-[4-(2-{[4-(4-L-alanyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-3-carboxamide</p>	





<p>N-[4-(2-{{[4-(4-D-protyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide</p>	
<p>N-[4-(2-{{[4-(4-L-protyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide</p>	
<p>N-{4-[2-(1H-benzimidazol-6-ylamino)-5-methylpyrimidin-4-yl]phenyl}acetamide</p>	
<p>4-(4-furan-2-ylphenyl)-N-(4-morpholin-4-ylphenyl)pyrimidin-2-amine</p>	





<p>N-(4-morpholin-4-ylphenyl)-4-[4-(pyrimidin-2-ylamino)phenyl]pyrimidin-2-amine</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl]cyclopropanecarboxamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]cyclopropanecarboxamide</p>	
<p>N-(4-{2-[(3,5-dimorpholin-4-ylphenyl)amino]-5-methylpyrimidin-4-yl}phenyl)-N²,N²-dimethylglycinamide</p>	





<p>N²,N²-dimethyl-N-(4-{5-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)glycinamide</p>	
<p>N-(4-{5-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)-D-prolinamide</p>	
<p>N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide</p>	





<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide</p>	
<p>N-ethyl-4-[4-({4-[4-(D-prolylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperazine-1-carboxamide</p>	
<p>N-[4-(2-{{4-(4-D-prolylpiperazin-1-yl)phenyl}amino})pyrimidin-4-yl]phenyl]-D-prolinamide</p>	
<p>N-[4-(2-{{4-(4-L-prolylpiperazin-1-yl)phenyl}amino})pyrimidin-4-yl]phenyl]-D-prolinamide</p>	

<p>1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)piperidine-2-carboxamide</p>	
<p>N-{4-[2-({4-[4-(piperidin-4-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-4-ylacetamide</p>	





<p>2-(3-fluorophenyl)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p>3-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)propanamide</p>	
<p>2-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p>2-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)-3-phenylpropanamide</p>	





<p>(1R,2R)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylcyclopropanecarboxamide</p>	
<p>2-(4-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>3-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide</p>	
<p>3-(3-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide</p>	





<p>3-(2-fluorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide</p>	
<p>Nalpha,Nalpha-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-phenylalaninamide</p>	
<p>2-(2-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-pyridin-2-ylacetamide</p>	






<p>2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-{4-[(trifluoromethoxy)phenyl]}acetamide</p>	
<p>2-[2-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>2-[3-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	





<p>2-[4-(methyloxy)phenyl]-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-alaninamide</p>	
<p>N-{4-[2-({4-[4-(N,N-dimethylglycyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-3-(methyloxy)propanamide</p>	





<p>(2R)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide</p>	
<p>N²,N²-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-alaninamide</p>	
<p>1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide</p>	
<p>N²,N²-dimethyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide</p>	





<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-1-phenylcyclopropanecarboxamide</p>	
<p>2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide</p>	
<p>(2S)-1-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)azetidine-2-carboxamide</p>	
<p>2,4,6-trichloro-N-(3-{[4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino}propyl)benzamide</p>	





<p>N-[3-({4-[3,4-bis(methoxy)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide</p>	
<p>2,6-dichloro-N-[3-({4-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-2-yl)amino)propyl]benzamide</p>	
<p>2,6-dichloro-N-(3-{[4-(2,3-dihydro-1,4-benzodioxin-6-yl)-5-fluoropyrimidin-2-yl]amino}propyl)benzamide</p>	
<p>2,6-dichloro-N-{3-[(4-{3-[(dimethylamino)methyl]phenyl}pyrimidin-2-yl)amino]propyl}benzamide</p>	





<p>2,6-dichloro-N-[3-({4-[3-(1-methylethyl)phenyl]pyrimidin-2-yl}amino)propyl]benzamide</p>	
<p>2,6-dichloro-N-{3-[(4-{4-[(1-methylethyl)oxy]phenyl}pyrimidin-2-yl)amino]propyl}benzamide</p>	
<p>N-[3-({4-[3-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2,6-dichlorobenzamide</p>	
<p>2,6-dichloro-N-[3-({4-[(E)-2-phenylethenyl]pyrimidin-2-yl}amino)propyl]benzamide</p>	
<p>phenyl (4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)carbamate</p>	

<p>phenylmethyl (4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)carbamate</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methoxy)propanamide</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide</p>	
<p>4-{4-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide</p>	






<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-3-(methoxy)propanamide</p>	
<p>3-(methoxy)-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}propanamide</p>	
<p>N-ethyl-4-(4-{[4-(4-{3-(methoxy)propanoyl}amino)phenyl]pyrimidin-2-yl}amino)phenyl)piperazine-1-carboxamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-2-phenylacetamide</p>	




<p>1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyrrolidin-2-one</p>	
<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide</p>	
<p>(2S)-2-hydroxy-3-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide</p>	





<p>(2R)-2-hydroxy-3-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide</p>	
<p>N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-alaninamide</p>	
<p>(2S)-2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylethanamide</p>	
<p>2-amino-2-(4-chlorophenyl)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	





<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)morpholine-3-carboxamide</p>	
<p>1-ethyl-3-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]urea</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]-D-prolinamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)-3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	





1-(2,6-dichlorophenyl)-3-(3-([4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino)propyl)urea	
1-[2-fluoro-5-(trifluoromethyl)phenyl]-3-(3-([4-(4-methyl-2-thienyl)pyrimidin-2-yl]amino)propyl)urea	
2,6-dichloro-N-[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]benzenesulfonamide	
N-[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]-2,6-difluorobenzenesulfonamide	
N-[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]naphthalene-2-sulfonamide	
N-[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]-3,4-bis(methoxy)benzenesulfonamide	
3-chloro-N-[3-([4-(4-(dimethylamino)phenyl)pyrimidin-2-yl]amino)propyl]propane-1-sulfonamide	




N-[3-({4-[4-(dimethylamino)phenyl]pyrimidin-2-yl}amino)propyl]propane-1-sulfonamide	
methyl (3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)propyl)carbamate	
1-methylethyl (3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)propyl)carbamate	
phenylmethyl (3-({4-(2,4-dichlorophenyl)pyrimidin-2-yl}amino)propyl)carbamate	
N-{4-[2-({3-(3-chlorophenyl)isoxazol-5-yl}methyl)amino]pyrimidin-4-yl}phenyl}acetamide	




<p>ethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)piperidine-1-carboxylate</p>	
<p>1,1-dimethylethyl 4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)piperidine-1-carboxylate</p>	
<p>N-(4-{2-[(4-cyanophenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	




<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyridin-4-yl}phenyl)acetamide</p>	
<p>1,1-dimethylethyl {1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidin-4-yl} carbamate</p>	
<p>N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide</p>	
<p>N-{1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidin-4-yl} acetamide</p>	




<p>4-(4-aminophenyl)-N-[4-(4-aminopiperidin-1-yl)phenyl]pyrimidin-2-amine</p>	
<p>N-[4-(2-({[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl)-3-(methoxy)propanamide</p>	
<p>N-{4-[2-({[4-(2-methylpropanoyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide</p>	
<p>N-{4-[2-({[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide</p>	





<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}tetrahydrofuran-2-carboxamide</p>	
<p>N-cyclopropyl-4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzamide</p>	
<p>N-[2-(methoxy)ethyl]-4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}benzamide</p>	
<p>2,6-dichloro-N-{3-[(4-pyridin-3-ylpyrimidin-2-yl)amino]propyl}benzamide</p>	





<p>2,6-dichloro-N-(3-{{4-(4-methyl-3,4-dihydro-2H-1,4-benzoxazin-7-yl)pyrimidin-2-yl}amino}propyl)benzamide</p>	
<p>2,6-dichloro-N-(3-{{4-(2,3-dihydro-1,4-benzodioxin-6-yl)-6-methylpyrimidin-2-yl}amino}propyl)benzamide</p>	
<p>N-(4-{2-[(3-{{(2,6-dichlorophenyl)carbonyl}amino}propyl)amino]pyrimidin-4-yl}phenyl)morpholine-4-carboxamide</p>	





<p>2,6-dichloro-N-{3-[(4-{4-[(cyclopropylcarbonyl)amino]phenyl}pyrimidin-2-yl)amino]propyl}benzamide</p>	
<p>N-(4-{2-[(3-{[(2,6-dichlorophenyl)carbonyl]amino}propyl)amino]pyrimidin-4-yl}phenyl)thiophene-2-carboxamide</p>	
<p>2,6-dichloro-N-(3-{[4-(4-{[N-(2-morpholin-4-ylethyl)glycyl]amino}phenyl)pyrimidin-2-yl]amino}propyl)benzamide</p>	





<p>1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)ethanone</p>	
<p>(1E)-1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)ethanone oxime</p>	
<p>N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-2-phenylacetamide</p>	





<p>N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-bromobenzamide</p>	
<p>N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-fluorobenzamide</p>	
<p>N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)propyl]-2-chlorobenzamide</p>	




<p>N-[4-(2-{[3-(morpholin-4-ylsulfonyl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-{4-[2-({3-[(cyclohexylmethyl)amino]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-{2-[(3-{[(5-bromo-2-fluorophenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(3-{[(2,5-dimethylphenyl)methyl]amino}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	




<p>N-(4-{2-[(3,4-dimorpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}cyclopropanecarboxamide</p>	
<p>N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide</p>	





<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}butanamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-2-carboxamide</p>	
<p>2-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide</p>	
<p>3-(methoxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide</p>	






<p>4-(methoxy)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)benzamide</p>	
<p>4-chloro-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)benzamide</p>	
<p>(2R)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	
<p>(2S)-N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	





<p>1-(2-hydroxyethyl)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-prolinamide</p>	<p>Chiral</p> 
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)thiophene-2-carboxamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-3-carboxamide</p>	
<p>2-phenyl-N-{4-[2-({4-[4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	





<p>3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-N-(diphenylmethyl)benzamide</p>	
<p>N-[4-(2-({4-[4-(4-methylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-{4-[2-({4-[4-(phenylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	






<p>N-{4-[2-({4-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({4-[4-(cyclohexylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-{2-[(4-{4-[(2-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	







<p>N-(4-{2-[(4-{4-[(3-fluorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(3-fluoro-4-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(3,5-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	







<p>N-[4-(2-([4-(4-{[3-(methoxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-(4-{2-[(4-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(4-methylphenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-({4-[4-(furan-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	





<p>N-(4-{2-[(4-{4-[(3-methylphenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-({4-[4-(phenylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({4-[4-(2-thienylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-{2-[(4-{4-[(4-fluorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	






<p>N-[4-(2-([4-(4-{4-(methoxy)phenyl)sulfonyl}piperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-(4-{2-[(4-{4-(4-chlorophenyl)sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-(4-{2-[(4-{4-(3-chlorophenyl)sulfonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	
<p>N-{4-[2-((4-[4-(biphenyl-4-ylsulfonyl)piperazin-1-yl]phenyl)amino)pyrimidin-4-yl]phenyl}acetamide</p>	





<p>N-{4-[2-({4-[4-(naphthalen-1-ylsulfonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-{2-[(3-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-[4-(2-{[3-(4-{3-(methyloxy)phenyl}methyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl]acetamide</p>	
<p>N-{4-[2-({3-[4-(3-methylbutyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2,3-dihydro-1,4-benzodioxin-6-ylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	






N-{4-[2-({3-[4-(cyclopropylmethyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	
N-(4-{2-[(3-{4-[3-(methylthio)propyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
N-(4-{2-[(3-{4-[(4-{[3-(dimethylamino)propyl]oxy}phenyl)methyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide	
N-{4-[2-({3-[4-({3-[(trifluoromethyl)oxy]phenyl)methyl]piperazin-1-yl}phenyl)amino)pyrimidin-4-yl]phenyl}acetamide	
4-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-N-phenylpiperazine-1-carboxamide	
N-[4-(2-{[3-(4-propanoylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	






<p>N-{4-[2-({3-[4-(phenylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2-phenylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(cyclopentylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2-pyridin-3-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2-cyclopentylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-(4-[2-({3-[4-[(2-chlorophenyl)carbonyl]piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl)acetamide</p>	





<p>N-(4-{2-[(3-{4-[(4-chlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(3-{4-[(3,4-dichlorophenyl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(3-{4-[(1-methyl-1H-pyrrol-2-yl)carbonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N²,N²-dimethyl-N-[4-(2-{[3-(methyloxy)-4-morpholin-4-yl]phenyl}amino)pyrimidin-4-yl]phenyl]glycinamide</p>	





<p>3-(methoxy)-N-[4-(2-{[3-(methoxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]propanamide</p>	
<p>N-(4-{2-[(4-{4-[(2-chlorophenyl)sulfonyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-{4-[2-({3-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-{4-[2-({3-[4-(2-cyclopropylacetyl)piperazin-1-yl]phenyl)amino]pyrimidin-4-yl}phenyl}acetamide</p>	
<p>N-[4-(2-{[3-(4-{[3-(methoxy)phenyl]carbonyl}piperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide</p>	




<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>1-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]piperidine-3-carboxylic acid</p>	
<p>1,1-dimethylethyl methyl{2-[(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)amino]-2-oxoethyl}carbamate</p>	
<p>1,1-dimethylethyl [4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]carbamate</p>	




<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-N²,N²-dimethylglycinamide</p>	
<p>4-(4-aminophenyl)-N-[4-(4-ethylpiperazin-1-yl)phenyl]pyrimidin-2-amine</p>	
<p>Nalpa-{[(1,1-dimethylethyl)oxy]carbonyl}-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-phenylalaninamide</p>	<p>Chiral</p> 
<p>Nalpa-{[(1,1-dimethylethyl)oxy]carbonyl}-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-phenylalaninamide</p>	<p>Chiral</p> 
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-phenylalaninamide</p>	<p>Chiral</p> 





<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-L-phenylalaninamide</p>	<p>Chiral</p> 
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-L-valinamide</p>	<p>Chiral</p> 
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-valinamide</p>	<p>Chiral</p> 
<p>1-ethyl-3-[4-(2-{[3-(methoxy)-4-morpholin-4-yl]phenyl}amino}pyrimidin-4-yl)phenyl]urea</p>	
<p>(2R)-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)piperidine-2-carboxamide</p>	<p>Chiral</p> 





<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl]acetamide</p>	
<p>4-{4-[(4-{4-[(N,N-dimethylglycyl)amino]phenyl}pyrimidin-2-yl)amino]phenyl}-N-ethylpiperazine-1-carboxamide</p>	
<p>N-{4-[2-({4-[4-(2,2-dimethylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycinamide</p>	
<p>N-{4-[2-({4-[4-(cyclobutylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycinamide</p>	





<p>N²,N²-dimethyl-N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl} amino)pyrimidin-4-yl]phenyl} glycineamide</p>	
<p>N-{4-[2-({4-[4-(cyclopropylcarbonyl)piperazin-1-yl]phenyl} amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycineamide</p>	
<p>N-{4-[2-({4-[4-(D-alanyl)piperazin-1-yl]phenyl} amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycineamide</p>	
<p>N-{4-[2-({4-[4-(L-alanyl)piperazin-1-yl]phenyl} amino)pyrimidin-4-yl]phenyl}-N²,N²-dimethylglycineamide</p>	





<p>N-(4-{2-[(1-{(2,6-dichlorophenyl)carbonyl}azetidin-3-yl)methyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-(4-{2-[(3-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl}phenyl)acetamide</p>	
<p>N-[3-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)cyclohexyl]-2,6-dichlorobenzamide</p>	







<p>N-{4-[2-({[4-(4-methylpiperazin-1-yl)phenyl]methyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-[4-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)phenyl]-2,6-dichlorobenzamide</p>	
<p>N-{4-[2-(piperidin-4-ylamino)pyrimidin-4-yl]phenyl}acetamide</p>	






<p>N-{4-[2-({1-[(2,6-dichlorophenyl)carbonyl]piperidin-4-yl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>N-{4-[2-({4-[(2-hydroxyethyl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide</p>	
<p>1-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-3-phenylurea</p>	
<p>N-[5-({4-[4-(acetylamino)phenyl]pyrimidin-2-yl}amino)-2-(4-ethylpiperazin-1-yl)phenyl]-2,6-dichlorobenzamide</p>	






<p>1-[4-(2-([4-(4-ethylpiperazin-1-yl)phenyl]amino)pyrimidin-4-yl)phenyl]-3-(phenylmethyl)urea</p>	
<p>N²,N²-dimethyl-N-{4-[2-([4-(4-(pyridin-3-ylcarbonyl)piperazin-1-yl]phenyl)amino)pyrimidin-4-yl]phenyl}glycinamide</p>	
<p>N-(3-fluoro-4-{2-([4-(4-morpholin-4-yl)phenyl]amino)pyrimidin-4-yl}phenyl)cyclopropanecarboxamide</p>	
<p>N-(4-{2-([4-(4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl)phenyl]amino)pyrimidin-4-yl}phenyl)cyclopropanecarboxamide</p>	





<p>N-[4-(2-{{[4-(4-L-alanyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-{{[4-(4-L-prolyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-{{[4-(4-D-alanyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-{{[4-(4-D-prolyl)piperazin-1-yl]phenyl}amino}pyrimidin-4-yl)phenyl]acetamide</p>	

<p>N-[4-(2-({4-[4-(2-piperazin-1-ylacetyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl)phenyl]acetamide</p>	
<p>N-[4-(2-({4-(4-L-alanyl)piperazin-1-yl}phenyl)amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	
<p>N-[4-(2-({4-(4-L-prolyl)piperazin-1-yl}phenyl)amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	
<p>N-[4-(2-({4-(4-D-alanyl)piperazin-1-yl}phenyl)amino)pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	

<p>N-[4-(2-{[4-(4-D-prolylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]tetrahydrofuran-2-carboxamide</p>	
<p>1-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-1H-pyrrole-2-carboxamide</p>	
<p>3-fluoro-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)pyridine-4-carboxamide</p>	
<p>6-methyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)pyridine-3-carboxamide</p>	
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)pyridazine-4-carboxamide</p>	
<p>2-cyclopropyl-N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide</p>	

<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)isoxazole-5-carboxamide</p>	
<p>N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)pyridine-3-carboxamide</p>	
<p>4-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)benzamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]-D-prolinamide</p>	
<p>N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]butanamide</p>	

<p>1-ethyl-3-[4-(2-{{4-(4-ethylpiperazin-1-yl)phenyl}amino}pyrimidin-4-yl)phenyl]urea</p>	
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)furan-3-carboxamide</p>	
<p>N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-1,3-thiazole-4-carboxamide</p>	
<p>2,6-dichloro-N-(3-{[4-(2,3-dihydro-1,4-benzodioxin-6-yl)pyrimidin-2-yl]amino}propyl)benzamide</p>	
<p>2,6-dichloro-N-(3-{[4-(4-fluorophenyl)pyrimidin-2-yl]amino}propyl)benzamide</p>	

N-(4-{2-[(2-methyl-4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
N-(4-{2-[(4-pyrrolidin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl)acetamide	
N-[4-(2-{[4-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	
N-[4-(2-{[3-(methyloxy)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide	

5

46. A compound according to claim 1 selected from

N-[4-(2-{[4-(4-ethylpiperazin-1-yl)phenyl]amino}pyrimidin-4-yl)phenyl]acetamide;

N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)cyclopropanecarboxamide;

N-(4-{5-methyl-2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-

10 yl)phenyl)cyclopropanecarboxamide;

N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)valinamide;

N-(4-{2-[(4-{4-[(1-methyl-1H-imidazol-2-yl)methyl]piperazin-1-yl}phenyl]amino}pyrimidin-4-yl)phenyl)acetamide;

2-(dimethylamino)-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)acetamide;

15 N-(4-{2-[(3,5-dimorpholin-4-yl)phenyl]amino}-5-methylpyrimidin-4-yl)phenyl)acetamide;

N-(4-{2-[(4-morpholin-4-yl)phenyl]amino}pyrimidin-4-yl)phenyl)-D-alaninamide;

- 5 N-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide;
2-amino-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-2-phenylacetamide;
N-(4-{5-methyl-2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)acetamide;
- 10 3-(methyloxy)-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-alaninamide;
N-(4-{2-[(4-{4-[3-(dimethylamino)-2,2-dimethylpropyl]piperazin-1-
- 15 yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide;
N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)acetamide;
2-Hydroxy-2-methyl-N-(4-(2-(4-morpholinophenylamino)pyrimidin-4-yl)phenyl)propanamide;
- 20 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide;
N-[4-(2-{[3-(methyloxy)-4-morpholin-4-ylphenyl]amino}pyrimidin-4-yl)phenyl]-D-prolinamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)tetrahydrofuran-3-carboxamide;
- 25 O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide;
1-ethyl-3-{4-[2-({4-[4-(2-methylpropanoyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}urea;
N-ethyl-4-(4-{[4-(4-{[(ethylamino)carbonyl]amino}phenyl)pyrimidin-2-yl]amino}phenyl)piperazine-1-carboxamide;
- 30 N²,N²-dimethyl-N-(4-{2-[(4-{4-[3-(methyloxy)propanoyl]piperazin-1-yl}phenyl)amino]pyrimidin-4-yl}phenyl)glycinamide;
N-(4-(2-(3-methoxy-4-morpholino-phenylamino)pyrimidin-4-yl)phenyl)acetamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
(3R)-3-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-
- 35 yl}phenyl)butanamide;
(3S)-3-hydroxy-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)butanamide;

- 5 N-{4-[2-({4-[4-(tetrahydrofuran-2-ylcarbonyl)piperazin-1-yl]phenyl}amino)pyrimidin-4-yl]phenyl}-D-prolinamide;
2-hydroxy-2-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)propanamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)prolinamide;
10 N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-prolinamide;
N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide;
O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-L-serinamide;
O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
O-methyl-N-(4-{2-[(4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-serinamide;
15 and
N-(4-{2-[(3-fluoro-4-morpholin-4-ylphenyl)amino]pyrimidin-4-yl}phenyl)-D-prolinamide.

47. A pharmaceutical composition comprising a compound according to claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, excipient,
20 or diluent.

48. A method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a compound according to claim 1, or a pharmaceutically salt thereof.
25

49. A method of inhibiting JAK-2 in a cell, comprising contacting the cell, in which inhibition of JAK-2 is desired, with a pharmaceutical composition comprising a compound according to claim 1, or a pharmaceutically acceptable salt thereof.

30 50. A method for treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a therapeutically effective amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof.

35 51. A method for treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said

5 treatment a pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof.

52. A pharmaceutical composition comprising a compound according to claim 45, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, excipient,
10 or diluent.

53. A pharmaceutical composition comprising a compound according to claim 46, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, excipient, or diluent.

15 54. A method for treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal in need of said treatment a pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof.

20 55. The method according to claim 54, wherein the disease, disorder or syndrome being treated is a myeloproliferative disorder, cancer, cardiovascular disease, and/or hematopoietic abnormality where hyperactivation of JAK-STAT signaling is present.

25 56. The method according to claim 55, wherein the myeloproliferative disorders are selected from myelofibrosis, thrombocythemia, polycythemia vera, essential thrombocythemia, agnogenic myeloid metaplasia, and chronic myelogenous leukemia.

57. The method according to claim 55, wherein the cancers are selected from
30 leukemias, lymphomas, multiple myeloma, prostate cancers, lung cancers, breast cancers, and ovarian cancers.

58. The method according to claim 55, wherein the cancers are selected from congestive heart failure and hypertension.

35 59. The method according to claim 55, wherein hematopoietic abnormality is thrombocytosis.

5

60. A method of treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal a compound according to claim 1, or a pharmaceutically acceptable salt thereof, in combination with one or more treatment(s) selected from surgery, one or more therapeutic agent(s), plateletpheresis, and
10 radiation.

61. A method of treating a disease, disorder, or syndrome mediated, at least in part, by inhibiting JAK-2, which method comprises administering to an animal a pharmaceutical composition comprising a therapeutically effective amount of a compound according to
15 claim1, or a pharmaceutically acceptable salt thereof, in combination with one or more treatment(s) selected from surgery, one or more therapeutic agent(s), plateletpheresis, and radiation.